Working references compiled primarily by the graduate students on the RAPP project: Elizabeth Beck, Ffion Cassidy & Sarina Loots.

Aarif K, P P (2014) Injured migratory shorebirds and gulls in the Kadalundi-community reserve. J Env Biol 35: 243–246.

Able K, AM (1995) Manipulations of polarized skylight calibrate magnetic orientation in a migratory bird. J Comp Physiol A Neuroethol Sens Neural Behav Physiol 177: 351–356.

ABMI (2013) The status of biodiversity in the Athabasca oil sands area. Preliminary Assessment 2013. The Alberta Biodiversity Monitoring Institute.

Acevedo MA, Villanueva-Rivera LJ (2006) Using automated digital recording systems as effective tools for the monitoring of birds and amphibians. Wildlife Soc B 34: 211–214.

Ackerman JT, Eadie JM, Moore TG (2006) Does life history predict risk-taking behavior of wintering dabbling ducks? Condor 108: 530–546.

Adams NJ (1994) Patterns and impacts of oiling of African penguins *Spheniscus demersus* - 1981-1991. Biol Cons 68: 35–41.

Agler B, Seiser P, Kendall S, Irons D (1994) Marine bird and sea otter population abundance of Prince William Sound, Alaska: trends following the T/V Exxon Valdez oil spill, 1989-93. Exxon Valdez Oil Spill Restoration Project Final Report (Restoration Project 93045). US Fish and Wildlife Service, Anchorage, Alaska, U.S.A.

AGRA Earth and Environmental Ltd (1996) Athabasca river bridge to steepbank mine river hydraulics and ice study. AGRA Earth and Environmental Ltd. Submitted to H. A. Simons for Suncor Inc. Oil Sands Group. Golder Project Report No. 952-2307

AGRA Earth and Environmental Limited (1996) Erosion resistance of Suncor's reclaimed sand structures. Prepared for Suncor Inc. Calgary, Alberta: AGRA Earth & Environmental Limited.

AGRA Earth and Environmental Ltd (2002) Preliminary results from long-term soil and vegetation plots established in the Oil Sands region. Prepared for Oil Sands Soil and Vegetation Working Group. Calgary, Alberta: AMEC Earth & Environmental Limited and Edmonton, Alberta: Can-Ag Enterprises Ltd.

AGRA Earth and Environmental Ltd (2002) Preliminary results from the land capability calibration sites in the Oil Sands region. Prepared for Oil Sands Soil and Vegetation Working Group. Calgary, Alberta: AMEC Earth & Environmental Limited and Edmonton, Alberta: Can-Ag Enterprises Ltd.

Aguilera E, Knight RL, Cummings JL (1991) An evaluation of two hazing methods for urban Canada geese. Wildlife Soc B 19: 32–35.

Ainley DG, Grau CR, Roudybush TE, Morrell SH, Utts JM (1981) Petroleum ingestion reduces reproduction in Cassin's auklet. Mar Pollut Bull 12: 314–317.

Ajie BC, Pintor LM, Watters J, Kerby JL, Hammond JI, et al. (2007) A framework for determining the fitness consequences of antipredator behavior. Behav Ecol 18: 267–270.

Akre CJ, Headley JV, Conly FM, Peru KM, Dickson LC (2004) Spatial patterns of natural polycyclic aromatic hydrocarbons in sediment in the lower Athabasca River. J Environ Sci Health A 39: 1163–1176.

Alabaster C, Hughes E (2010) Examination of the effect of array weighting function on radar target detectability. IEEE Trans Aerosp Electron Syst 46: 1364–1375.

Albers P, Szaro R (1978) Effects of no. 2 fuel oil on common eider eggs. Mar Pollut Bull 9: 138–139.

Albers PH (1977) Effects of external applications of fuel oil on hatchability of mallard eggs. In DA Wolfe, editor. Fate and effects of petroleum hydrocarbons in marine organisms and ecosystems. Pergammon Press. pp.158–163.

Albers PH (1977) Effects of oil on aquatic birds. In: Paul L. Fore, editor. Proceedings of the 1977 Oil Spill Response Workshop. Washington, DC. pp. 61–68.

Albers PH (1978) The effects of petroleum of different stages of incubation in bird eggs. Bull Environ Contam Toxicol 19: 624–630.

Albers PH (1980) Transfer of crude oil from contaminated water to bird eggs. Environ Res 22: 307–314.

Albers PH (1983) Effects of oil on avian reproduction: a review and discussion. In The Effects of Oil on Birds. A Multi-discipline Symposium. Tri-State Bird Rescue & Research, Inc., Newark, DE. pp. 78–96.

Albers PH (1987) Petroleum and individual polycyclic aromatic hydrocarbons. In: Hoffman, David J., editor. Handbook of ecotoxicology. Boca Raton, Florida: Lewis Publishers. pp. 330–355.

Albers PH (1990) Oil spills and the environment: a review of chemical fate and biological effects of petroleum. The Effects of Oil on Wildlife: Research, Rehabilitation, and General Concerns. In: Proceedings from the Oil Symposium, Herdon.

Albers PH (1995) Oil, biological communities and contingency planning. In: Frink L, Ball-Weir K, and Smith C, editors. Wildlife and Oil Spills: Response, Research, and Contingency Planning. Newark, DE. pp. 1–9.

Albers PH (2006) Birds and polycyclic aromatic hydrocarbons. Avian Poultry Biol Rev 17: 125–140.

Albers PH (2007) An Annotated Bibliography on Petroleum Pollution. Version 2007; USGS Patuxent Wildlife Research Center, Laurel, Maryland, USA.

Albers PH, Belisle AA, Swineford DM, Hall RJ (1985) Environmental contamination in the oil fields of western Pennsylvania. Oil Petrochemical Pollution 2: 265–280.

Albers PH, Gay ML (1982) Effects of a chemical dispersant and crude oil on breeding ducks. Bull Environ Contam Toxicol 29: 404–411.

Albers PH, Gay ML (1982) Unweathered and weathered aviation kerosine: Chemical characterization and effects on hatching success of duck eggs. Bull Environ Contam Toxicol 28: 430–434.

Albers PH, Heinz GH (1983) FLIT-MLO and No. 2 fuel oil: effects of aerosol applications to mallard eggs on hatchability and behavior of ducklings. Environ Res 30: 381–388.

Albers PH, Service TAE (1991) Oil spills and living organisms. Texas Agricultural Extension Service. College Station, Texas. pp. 9

Alberta Environment (2008) Guideline for wetland establishment on reclaimed oil sands leases (2nd edition). Prepared by Harris, M.L. of Lorax Environmental for the Wetlands and Aquatics Subgroup of the Reclamation Working Group of the Cumulative Environmental Management Association, Fort McMurray, AB. Canada.

Alberta Environmental Centre (1986). Acute and subacute toxicity of different fractions of Athabasca bitumen to fish. Alberta Environmental Centre, Vegreville, AB, Canada. 35 pp, 1986. to fish. AECV86-RS.

Alberta Government (2013) Facts and Statistics. Available: http://www.energy.alberta.ca/OilSands/791.asp.

Alberta Government. Environmental Protection and Enhancement Act. Chapter E-12. 2010; Available: http://www.qp.alberta.ca/documents/acts/e12.pdf

Alberta_Government (2012) 2011-2012 Environment and Water Annual Report. pp. 1-118.

Alberta_Government (2013) Oil Sands Information Portal. Available: http://www.environment.alberta.ca/apps/osip/. Accessed 27 November 2013.

Aldhous P (2010) Let there be night, for wildlife's sake. New Sci: 8–9.

Alerstam T, Chapman JW, Bäckman J, Smith AD, Karlsson H, et al. (2011) Convergent patterns of long-distance nocturnal migration in noctuid moths and passerine birds. Proc R Soc B Biol Sci 278: 3074–3080.

Allan J (2006) A heuristic risk assessment technique for birdstrike management at airports. Risk Anal 26: 723–729.

Alldredge MW, Pacifici K, Simons TR, Pollock KH (2008) A novel field evaluation of the effectiveness of distance and independent observer sampling to estimate aural avian detection probabilities. J Appl Ecol 45: 1349–1356.

Alldredge MW, Pollock KH, Simons TR, Johnson D (2006) Estimating detection probabilities from multiple-observer point counts. Auk 123: 1172–1182.

Allen C (1990) Mitigating impacts to wildlife at FMC Gold Company's Paradise Peak mine. In Proceedings of the Nevada wildlife/mining workshop. Nevada Mining Association: Reno, NV, U.S.A. pp. 27–29.

Allen EW (2008) Process water treatment in Canada's oil sands industry: I. Target pollutants and treatment objectives. J Environ Eng Sci 7: 123–138.

Al-Muhareb E, Morgan T, Herod A, Kandiyoti R (2007) Characterization of petroleum asphaltenes by size exclusion chromatography, UV-fluorescence and mass spectrometry. Pet Sci Technol 25: 81–91.

Alonso-Alvarez C, Munilla I, López-Alonso M, Velando A (2007) Sublethal toxicity of the Prestige oil spill on yellow-legged gulls. Environ Int 33: 773–781.

Alonso-Alvarez C, Pérez C, Velando A (2007) Effects of acute exposure to heavy fuel oil from the Prestige spill on a seabird. Aquat Toxicol 84: 103–110.

Altwegg R, Crawford RJ, Underhill LG, Williams ATJ (2008) Long-term survival of de-oiled Cape gannets *Morus capensis* after the Castillo de Bellver oil spill of 1983. Biol Cons 141: 1924–1929.

Álvarez F (2004) The conspicuous gape of the nestling common cuckoo *Cuculus canorus* as a supernormal stimulus for rufous bush chat *Cercotrichas galactotes* hosts. Ardea 92: 63–68.

Amcoff M, Lindqvist C, Kolm N (2013) Sensory exploitation and plasticity in female mate choice in the swordtail characin. Anim Behav 85: 891–898.

AMEC Earth and Environmental (2005) Air Quality Assessment of the Deer Creek Energy Limited Joselyn North Mine Project. Submitted to Deer Creek Energy Limited. Calgary, Alberta, Canada.452 p.

Amirault-Langlais DL, Thomas PW, McKnight J (2007) Oiled piping plovers (*Charadrius melodus melodus*) in eastern Canada. Waterbirds 30: 271–274.

Amo L, Galvan I, Tomas G, Sanz J (2008) Predator odour recognition and avoidance in a songbird. Funct Ecol 22: 289–293.

Andelt WF, Hopper SN (1996) Effectiveness of alarm-distress calls for frightening herons from a fish rearing facility. The Progressive fish-culturist 58: 258–262.

Anderies JM, Katti M, Shochat E (2007) Living in the city: resource availability, predation, and bird population dynamics in urban areas. J Theor Biol 247: 36–49.

Anderson DW, Gress F, Fry DM (1996) Survival and dispersal of oiled brown pelicans after rehabilitation and release. Mar Pollut Bull 32: 711–718.

Anderson DW, Newman SH, Kelly PR, Herzog SK, Lewis KP (2000) An experimental soft-release of oil-spill rehabilitated American coots (*Fulica americana*): I. Lingering effects on survival, condition and behavior. Environ Pollut 107: 285–294.

Anderson J (2014) Organic matter accumulation in reclaimed soils beneath different vegetation types in the Athabasca oil sands. Master of Science, The University of British Columbia, Vancouver, British Columbia, Canada.

Anderson J editor (1975) Laboratory studies on the effects of oil on marine organisms: an overview. API Publication No. 4249. Submitted to American Petroleum Institute Division of Environmental Affairs.

Anderson JC (2012) Acute and chronic toxicity of untreated, aged, and ozonated oil sands process-affected water in *Chironomus dilutus* larvae.Master of Science, University of Saskatchewan, Saskatchewan, Canada.

Andres BA (1997) The Exxon Valdez oil spill disrupted the breeding of black oystercatchers. The J Widlife Manag 61: 1322–1328.

Andres BA (1999) Effects of persistent shoreline oil on breeding success and chick growth in Black Oystercatchers. Auk 116: 640–650.

Arnold TW, Zink RM (2011) Collision mortality has no discernible effect on population trends of North American birds. PLoS ONE 6: e24708.

Avery M (2002) Birds in pest management. In: Pimental D, editor. Encyclopedia of pest management. USDA National Wildlife Research Center-Staff Publications. pp. 104-106.

Avery M, Springer PF, Cassel JF (1976) The effects of a tall tower on nocturnal bird migration: a portable ceilometer study. Auk 93: 281–291.

Avery ML, Daneke DE, Decker DG, Lefebvre PW, Matteson RE, et al. (1988) Flight pen evaluation of eyespot balloons to protect citrus from bird depredations. In: Proceedings of the Thirteenth Vertebrate Pest Conference. Paper 56.

Avery ML, Genchi AC (2004) Avian perching deterrents on ultrasonic sensors at airport windshear alert systems. Wildlife Soc B 32: 718–725.

Avery ML, Humphrey JS, Tillman EA, Phares KO, Hatcher JE (2002) Dispersing vulture roosts on communication towers. J Raptor Res 36: 45–50.

Avery, M. L. 2003. Avian repellants. In: Plimmer JR, Gammon DW, Ragsdale NN editors. Encyclopedia of Agrochemicals. John Wiley & Sons, Hoboken, New Jersey, USA.

Avilés JM, Soler JJ, Pérez-Contreras T (2006) Dark nests and egg colour in birds: a possible functional role of ultraviolet reflectance in egg detectability. Proc R Soc B Biol Sci 273: 2821–2829.

Avilés JM, Soler JJ, Pérez-Contreras T, Soler M, Møller AP (2006) Ultraviolet reflectance of great spotted cuckoo eggs and egg discrimination by magpies. Behav Ecol 17: 310–314.

Ayles GB, Dubé M, Rosenberg D (2004) Oil sands regional aquatic monitoring program (RAMP) scientific peer review of the five year report (1997-2001). Submitted to the RAMP Stering Committee.

Baar L, Matlack RS, Johnson WP, Barron RB (2008) Migration chronology of waterfowl in the Southern High Plains of Texas. Waterbirds 31: 394–401.

Bäckman J, Alerstam T (2003) Orientation scatter of free-flying nocturnal passerine migrants: components and causes. Anim Behav 65: 987–996.

Baddaloo EG (1986) In: Proceedings of Alberta oil sands tailings wastewater treatment technology workshop, Fort McMurray.

Baddaloo EG, Mah-Paulson MH, Verbeek AG, Niimi AJ (1999) In: Proceedings of the 26th Annual Aquatic Toxicity Workshop: October 4-6, 1999, Edmonton, Alberta.

Baek S, Field R, Goldstone M, Kirk P, Lester J, et al. (1991) A review of atmospheric polycyclic aromatic hydrocarbons: sources, fate and behavior. Water Air Soil Pollut 60: 279–300.

Baillie SR, Mead C (1982) The effect of severe oil pollution during the winter of 1980-81 on British and Irish Auks. Ringing & Migration 4: 33–44.

Baird PH (1990) Concentrations of seabirds at oil-drilling rigs. Condor 92: 768–771.

Bairlein F (2003) The study of bird migrations-some future perspectives: capsule routes and destinations have been unveiled but modern techniques offer the chance to explore much more. Bird Study 50: 243–253.

Bajzak D, Piatt JF (1990) Computer-aided procedure for counting waterfowl on aerial photographs. Wildlife Soc B 18: 125–129.

Ball JR, Bayne EM (2012) Using video monitoring to assess the accuracy of nest fate and nest productivity estimates by field observation. Auk 129: 438–448.

Ball SA (2009) Suspending vulture effigies from roosts to reduce bird strikes. Hum–Wildl Confl 3: 257–259.

Ballachey BE, Bodkin JL, Monson DH (2013) Quantifying long-term risks to sea otters from the 1989 "Exxon Valdez" oil spill: Reply to Harwell & Gentile (2013). Mar Ecol Prog Ser 488: 297–301.

Balseiro A, Espí A, Márquez I, Pérez V, Ferreras M, et al. (2005) Pathological features in marine birds affected by the Prestige's oil spill in the north of Spain. J Wildl Dis 41: 371–378.

Banks A, Sanderson W, Hughes B, Cranswick P, Smith L, Whitehead S, Musgrove A, Haycock B, Fairney N (2008) The Sea Empress oil spill (Wales, UK): Effects on common scoter *Melanitta nigra* in Carmarthen Bay and status ten years later. Mar Pollut Bull 56: 895–902.

Banks RC (1979) Human related mortality of birds in the United States. US Department of the Interior, Fish and Wildlife Service Washington, DC, USA.

Bardeli R, Wolff D, Kurth F, Koch M, Tauchert K-H, et al. (2010) Detecting bird sounds in a complex acoustic environment and application to bioacoustic monitoring. Pattern Recognit Lett 31: 1524–1534.

Barham PJ, Crawford RJ, Underhill LG, Wolfaardt AC, Barham BJ, et al. (2006) Return to Robben Island of African Pengui Sci Total Environ ns that were rehabilitated, relocated or reared in captivity following the Treasure oil spill of 2000. Ostrich-J Afr Ornithol 77: 202–209.

Bari M, Kindzierski W, Cho S (2014) A wintertime investigation of atmospheric deposition of metals and polycyclic aromatic hydrocarbons in the Athabasca oil sands region, Canada. Sci Total Environ 485: 180–192.

Barjaktarovic L, Bishay FS, Nix PG (1995) Ceriodaphnia dubia and Chironomus tentans in situ toxicity tests to assess oil sands wastewater treatment in constructed wetlands. Society of Environmental Toxicology and Chemistry 2nd World Congress, Oil Sands Session, November 5-9, 1995, Vancouver, BC, Canada.

Barrett RT (1979) Small oil spill kills 10-20 000 seabirds in north Norway. Mar Pollut Bull 10: 253–255.

Barron MG (2012) Ecological impacts of the Deepwater Horizon oil spill: implications for immunotoxicity. Toxicol Pathol 40: 315–320.

Bateman P, Fleming P (2011) Who are you looking at? Hadeda ibises use direction of gaze, head orientation and approach speed in their risk assessment of a potential predator. J Zool 285: 316–323.

Bateson P, Laland KN (2013) Tinbergen's four questions: an appreciation and an update. Trends Ecol Evol 28: 712–718.

Battersby JE, Greenwood JJ (2004) Monitoring terrestrial mammals in the UK: past, present and future, using lessons from the bird world. Mamm Rev34: 3–29.

Bauer S, Van Dinther M, Høgda KA, Klaassen M, Madsen J (2008) The consequences of climate-driven stop-over sites changes on migration schedules and fitness of Arctic geese. J Anim Ecol 77: 654–660.

Baxter A (2007) Laser dispersal of gulls from reservoirs near airports. 2007 Bird Strike Committee USA/Canada, 9th Annual Meeting, Kingston, Ontario, Canada.Paper 2. http://digitalcommons.unl.edu/birdstrike2007/2

Baxter A, Bell J, Allan J, Fairclough J (1999) The interspecificity of distress calls. 1999 Bird Strike Committee-USA/ Canada, First Joint Annual Meeting, Vancouver, BC, Canada.Paper 8. http://digitalcommons.unl.edu/birdstrike1999/8

Bayer RD (1988) Oiled birds: how to search for and capture oiled birds at Oregon intertidal areas. Newport, Oregon: Gahmken Press. 30 p.

Bayne E, Nielsen B (2011) Temporal trends in bird abundance in response to thinning of lodgepole pine (*Pinus contorta*). Can J Forest Res 41: 1917–1927.

Bayne EM, Campbell J, Haché S (2012) Is a picture worth a thousand species? Evaluating human perception of biodiversity intactness using images of cumulative effects. Ecol Indic 20: 9–16.

Bayne EM, Habib L, Boutin S (2008) Impacts of chronic anthropogenic noise from energy-sector activity on abundance of songbirds in the boreal forest. Conserv Biol 22: 1186–1193.

Bayne EM, Scobie CA, Rawson-Clark M (2012) Factors influencing the annual risk of birdwindow collisions at residential structures in Alberta, Canada. Wildlife Res 39: 583–592.

Beak Associates Consulting Ltd. (1988) 1983 Trace element concentrations in benthic invertebrates and sediments in the Athabasca River near the Suncor tar island plant site. Report prepared for Suncor Limited. 23 p.

Beason R, Nohara TJ, Weber P (2013) Beware the Boojum: caveats and strengths of avian radar. Hum-Wildl Interact 7: 16–46.

Beason RC (1978) The influences of weather and topography on water bird migration in the southwestern United States. Oecologia 32: 153–169.

Beason RC, Nohara TJ, Weber P (2010) Color vision as a model for precise altitude determination for avian radar. Proceedings of the 29th International Bird Strike Committee. Paper 12. Cairns, Australia. http://www.int-birdstrike.org/Cairns_Papers/IBSC29%20WP12.pdf

Beauchamp G (2010) Determinants of false alarms in staging flocks of semipalmated sandpipers. Behav Ecol 21: 584–587.

Beck E (2014) Health of domestic mallards (*Anas platyrhynchos domestica*) following exposure to oil sands process-affected water. Master of Science, University of Alberta, Edmonton, Alberta, Canada.

Becker DA, Wood PB, Keyser PD, Wigley TB, Dellinger R, et al. (2011) Threshold responses of songbirds to long-term timber management on an active industrial forest. Forest Ecol Manag 262: 449–460.

Begg G, Reid J, Tasker M, Webb A (1997) Assessing the vulnerability of seabirds to oil pollution: sensitivity to spatial scale. Colon Waterbird 20: 339–352.

Beier N, Sego D (2008) The oil sands tailings research facility. Geotechnical News 26: 72–77.

Belant JL, Gabrey SW, Dolbeer RA, Seamans TW (1995) Methyl anthranilate formulations repel gulls and mallards from water. Crop Prot 14: 171–175.

Belant JL, Woronecki PP, Dolbeer RA, Seamans TW (1998) Ineffectiveness of five commercial deterrents for nesting starlings. Wildlife Soc B: 264–268.

Belisle AA, Gay ML, Coon NC (1981) Comparison of two extraction methods for the analysis of petroleum hydrocarbon residues in mallard duck eggs by GC and GC-MS. Chemosphere 10: 1197–1203.

Bellrose FC (1976) Ducks, Geese and Swans of North America. Harrisburg, Pennsylvania: Stackpole Books. 568 p.

Bendell-Young L (1997) Assessing the ecological characteristics of wetlands receiving oil sands effluent. Prepared for Suncor Inc. Oil Sands by Simon Fraser University

Bendell-Young L, Bennett K, Crowe A, Kennedy C, Kermode A, et al. (2000) Ecological characteristics of wetlands receiving an industrial effluent. Ecol Appl 10: 310–322.

Bendell-Young L, Farrell T, Moore M, Plant A, Kermode A, et al. (1996) Determining the ecological viability of constructed wetlands for the treatment of oil sands wastewater. Prepared for Suncor Inc. Oil Sands Group and Syncrude Canada Limited, by Simon Fraser University. Burnaby, B.C. Canada.

Bendell-Young LI (1997) Assessing the Ecological Viability of Wetlands for Treatment of Oil Sands Wastewaters; A Feasibility Study. Prepared for Suncor Inc. Oil Sands Group, by Simon Fraser University. Burnaby, B.C. Canada.

Bendell-Young, L. I., Farrell A. P., Kennedy C. J., Kermonde A., Moore M. M., & Plant A. L. Ecological viability of wetlands receiving oil sands effluent. In: Proceedings of the 23rd Annual Aquatic Toxicity Workshop, October 7-9, 1996, Calgary, Alberta, editors, J. S. Goudey, S. M. Swanson, M. D. Treissman and A. J. Niimi Canadian Technical Report of Fisheries and Aquatic Sciences 2144

Bennett A, Cuthill I (1994) Ultraviolet vision in birds: what is its function? Vis Res34: 1471–1478.

Bennett ATD, Théry M (2007) Avian color vision and coloration: multidisciplinary evolutionary biology. Am Nat 169: S1–S6.

Bennett KE (1997) Assessing the health of juvenile mallards held in situ on an industrially-based wetland using physiological endpoints: implications for populations. Master of Science, Simon Fraser University.

Bennun L, Matiku P, Mulwa R, Mwangi S, Buckley P (2005) Monitoring important bird areas in Africa: towards a sustainable and scaleable system. Biodivers Conserv 14: 2575–2590.

Berg C (2003) Best practices for migratory bird care during oil spill response. Anchorage, Alaska: US Fish and Wildlife Service, Ecological Services Field Office.

Berge A, Delwiche M, Gorenzel W, Salmon T (2007) Sonic broadcast unit for bird control in vineyards. Appl Eng Agric 23: 819-825.

Berge A, Delwiche M, Gorenzel WP, Salmon T (2007) Bird Control in vineyards using alarm and distress calls. Am J Enol Vitic 58: 135–143.

Berkes F, Colding J, Folke C (2000) Rediscovery of traditional ecological knowledge as adaptive management. Ecol Appl 10: 1251–1262.

Bernáth B, Szedenics G, Molnár G, Horváth G (2001) Visual ecological impact of a peculiar oil lake on the avifauna: dual choice field experiments with water-seeking birds using huge shiny black and white plastic sheets. Arch Nat Conserv Landsc Res 40: 1–28.

Bernhardt GE, Blackwell BF, DeVault TL, Kutschbach-Brohl L (2010) Fatal injuries to birds from collisions with aircraft reveal anti-predator behaviours. Ibis 152: 830–834.

Berthold P (1993) Bird migration: a general survey. Oxford University Press, Oxford, UK.

Berthold P (2001) Bird migration: a general survey. Oxford University Pres, Oxford, UK.

Berthold P, Fiedler W, Querner P (2000) Migratory restlessness or Zugunruhe in birds-a description based on video recordings under infrared illumination. Journal fur Ornithologie 141: 285–299.

Bevanger K (1994) Bird interactions with utility structures: collision and electrocution, causes and mitigating measures. Ibis 136: 412–425.

Bevanger K (1995) Estimates and population consequences of tetraonid mortality caused by collisions with high tension power lines in Norway. J Appl Ecol: 745–753.

BGC Engineering Inc., 2010. Review of reclamation options for oil sands tailings substrates. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-2. 59 pp.

BGC_Engineering_Inc (2010) Oil sands tailings technology review. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-1. 136 pp.

Bibby CJ (1981) An experiment on the recovery of dead birds from the North Sea. Ornis Scandinavica 12: 261–265.

Biber E (2011) Problem of environmental monitoring. U Colo L Rev 83: 1. Available at: http://scholarship.law.berkeley.edu/facpubs/1606

Bigger D, Peery MZ, Chinnici S, Courtney SP (2006) Efficacy of audiovisual and radar surveys for studying marbled murrelets in inland habitats. J Widlife Manag 70: 505–516.

Birkholz D, Goudey S, Nelson R, Gulley J, MacKinnon M (1995) The effects of oil sands wastewaters on fish exposed to sub-lethal concentrations: field and laboratory studies. Edmonton, Alberta: Enviro-Test Laboratories.

Birks SJ, Yi Y, Cho S, Gibson JJ, Hazewinkel R (2013) Characterizing the organic composition of snow and surface water in the Athabasca region. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-40. 62 pp.

Bishay FS, Bayley SE, Nix PG (1995) Ammonia removal in constructed wetlands receiving oil sands waste water, Ft. McMurray, AB. Society of Environmental Toxicology and Chemistry 2nd World Congress, Oil Sands Session, November 5-9, 1995, Vancouver, B.C., Canada.

Blackwell BF (2002) Understanding avian vision: the key to using light in bird management. USDA National Wildlife Res Center-Staff Publications: 465.Available: http://digitalcommons.unl.edu/icwdm usdanwrc/465

Blackwell BF, Bernhardt GE (2004) Efficacy of aircraft landing lights in stimulating avoidance behavior in birds. J Widlife Manag 68: 725–732.

Blackwell BF, Bernhardt GE, Cepek JD, Dolbeer RA (2002) Lasers as non-lethal avian repellents: potential applications in the airport environment. USDA National Wildlife Research Center - Staff Publications. Available: http://digitalcommons.unl.edu/icwdm usdanwrc/147

Blackwell BF, Bernhardt GE, Dolbeer RA (2002) Lasers as nonlethal avian repellents. J Widlife Manag 66: 250–258.

Blackwell BF, DeVault TL, Seamans TW, Lima SL, Baumhardt P, et al. (2012) Exploiting avian vision with aircraft lighting to reduce bird strikes. J Appl Ecol 49: 758–766.

Blackwell BF, Fernández-Juricic E, Seamans TW, Dolan T (2009) Avian visual system configuration and behavioural response to object approach. Anim Behav77: 673–684.

Blackwell BF, Seamans TW, Schmidt PM, Vault TL, Belant JL, et al. (2013) A framework for managing airport grasslands and birds amidst conflicting priorities. Ibis 155: 199–203.

Blackwell BF, Wright SE (2006) Collisions of red-tailed hawks (*Buteo jamaicensis*), turkey vultures (*Cathartes aura*), and black vultures (*Coragyps atratus*) with aircraft: implications for bird strike reduction. J Raptor Res 40: 76–80.

Blaich CF, Miller DB (1988) Alarm call responsivity of mallard ducklings (*Anas platyrhynchos*): VI. Effects of sibling and self-produced auditory stimulation. J Comp Psychol 102: 56-60.

Blaise C, Gagné F (2013) Aquatic ecotoxicology: what has been accomplished and what lies ahead? An Eastern Canada historical perspective. J Xenobiotics 3:e8

Blake B (1983) A comparative study of the diet of auks killed during an oil incident in the Skagerrak in January 1981. J Zool 201: 1–12.

Blesdoe EK, Blumstein DT (2014) What is the sound of fear? Behavioral responses of white-crowned sparrows *Zonotrichia leucophrys* to synthesized nonlinear acoustic phenomena. Curr Zool. In press.

Bley T, Bessei W (2008) Recording of individual feed intake and feeding behavior of Pekin ducks kept in groups. Poult Sci 87: 215–221.

Blokpoel H (1973) Bird migration forecasts for military air operations. Occasional Paper Number 16. Canadian Wildlife Service, Environment Canada, Ottawa, ON, Canada.

Blokpoel H, Burton J (1973) Weather and height of nocturnal migration in east central Alberta: a radar study. Bird Banding 46: 311-328.

Bluff LA, Rutz C (2008) A quick guide to video-tracking birds. Biol Lett 4: 319–322.

Blumstein DT (2003) Flight-initiation distance in birds is dependent on intruder starting distance. J Widlife Manag: 852–857.

Blumstein DT (2010) Flush early and avoid the rush: a general rule of antipredator behavior? Behav Ecol 21: 440–442.

Blumstein DT, Anthony LL, Harcourt R, Ross G (2003) Testing a key assumption of wildlife buffer zones: is flight initiation distance a species-specific trait? Biol Cons 110: 97–100.

Blumstein DT, Daniel JC, Griffin AS, Evans CS (2000) Insular tammar wallabies (*Macropus eugenii*) respond to visual but not acoustic cues from predators. Behav Ecol 11: 528–535.

Boag D, Lewin V (1980) Effectiveness of three waterfowl deterrents on natural and polluted ponds. J Widlife Manag 44: 145–154.

Bocetti CI (2011) Cruise ships as a source of avian mortality during fall migration. Wilson J Ornithol 123: 176–178.

Boersma P (1986) Ingestion of petroleum by seabirds can serve as a monitor of water quality. Science 231: 373–376.

Boersma PD (1987) Penguins oiled in Argentina. Science 236: 135–135.

Boersma PD, Davies EM, Reid WV (1988) Weathered crude oil effects on chicks of fork-tailed storm-petrels (*Oceanodroma furcata*). Arch Environ Contam Toxicol 17: 527–531.

Bomford M, O'Brien PH (1990) Sonic deterrents in animal damage control: a review of device tests and effectiveness. Wildlife Soc B: 411–422.

Bonebrake TC, Christensen J, Boggs CL, Ehrlich PR (2010) Population decline assessment, historical baselines, and conservation. Conserv Lett 3: 371–378.

Booth AL, Muir BR (2013) "How far do you have to walk to find peace again?": A case study of First Nations' operational values for a community forest in Northeast British Columbia, Canada. Natural Resources Forum. Vol. 37. pp. 153–166.

Booth AL, Skelton NW (2011) Improving First Nations' participation in environmental assessment processes: recommendations from the field. Impact Assessment and Project Appraisal 29: 49–58.

Boreas Environmental Consulting Services Ltd. (1987) Suncor avian research program: ponds 1, 1a, 2, 3 (April - October 1987). Prepared for Suncor Inc.

Boreas Environmental Consulting Services Ltd. (1989) Suncor avian research program: observation data from 1989. Prepared for Suncor Inc.

Bott RD (2012) Evolution of Canada's Oil and Gas Industry. Calgary, Alberta: Canadian Centre for Energy Information.

Bourne W (1978) Amoco Cadiz seems likely to exterminate the French auks. Mar Pollut Bull 9: 145.

Bourne W (1979) The Christos Bitas affair. Mar Pollut Bull 10: 122–123.

Bourne W, Bibby C (1975) Temperature and the seasonal and geographical occurrence of oiled birds on west European beaches. Mar Pollut Bull 6: 77–80.

Bourne W, Johnston L (1971) The threat of oil pollution to north Scottish seabird colonies. Mar Pollut Bull 2: 117–119.

Bourne WRP (1968) Observation of an encounter between birds and floating oil. Nature 219: 632.

Bourne WRP, Parrack JD, Potts GR (1967) Birds killed in the Torrey Canyon disaster. Nature 215: 1123-1125.

Boutin S, Boyce MS, Hebblewhite M, Hervieux D, Knopff KH, et al. (2012) Why are caribou declining in the oil sands? Front Ecol Environ 10: 65–67.

Boutin S, Haughland DL, Schieck J, Herbers J, Bayne E (2009) A new approach to forest biodiversity monitoring in Canada. Forest Ecol Manag 258: S168–S175.

Bowman TD, Schempf PF, Bernatowicz JA (1995) Bald eagle survival and population-dynamics in Alaska after the Exxon Valdez oil-spill. The J Widlife Manag 59: 317–324.

Bowman TD, Schempf PF, Hodges JI (1997) Bald eagle population in Prince William Sound after the Exxon Valdez oil spill. The J Widlife Manag: 962–967.

Boyce MS, Vernier PR, Nielsen SE, Schmiegelow FKA (2002) Evaluating resource selection functions. Ecol Modell 157: 281–300.

Boyle WA, Norris DR, Guglielmo CG (2010) Storms drive altitudinal migration in a tropical bird. Proc R Soc B Biol Sci 277: 2511–2519.

Bradley BA, Olsson AD, Wang O, Dickson BG, Pelech L, et al. (2012) Species detection vs. habitat suitability: Are we biasing habitat suitability models with remotely sensed data? Ecol Modell 244: 57–64.

Brand M, Key G, Nohara TJ, Beason R, Herricks E, et al. (2010) Integration and validation of avian radars (IVAR), Final Report. ESTCP Project RC-200723.

Brandes TS (2008) Automated sound recording and analysis techniques for bird surveys and conservation. Bird Conservation International 18: S163–S173.

Brausch JM, Blackwell BR, Beall BN, Caudillo C, Kolli V, et al. (2010) Effects of polycyclic aromatic hydrocarbons in Northern Bobwhite Quail (*Colinus virginianus*). J Toxicol Environ Health A 73: 540–551.

Bridge ES, Thorup K, Bowlin MS, Chilson PB, Diehl RH, et al. (2011) Technology on the move: recent and forthcoming innovations for tracking migratory birds. Biosci 61: 689–698.

Briggs KT, Gershwin ME, Anderson DW (1997) Consequences of petrochemical ingestion and stress on the immune system of seabirds. ICES J Mar Sci 54: 718–725.

Briggs KT, Yoshida SH, Gershwin ME (1996) The influence of petrochemicals and stress on the immune system of seabirds. Regul Toxicol Pharm 23: 145–155.

Briot J, Bataille P (2003) A new laser equipment designed for avian dispersal in airport environment. International Birdstrike Committee. IBSC26/WP-AE7. Warsaw, Poland.

Briot JL (2005) Last experiments with a laser equipment designed for avian dispersal in airport environment. IBSC27/WP V-1. Athens, Greece.

Brough T (1969) The dispersal of starlings from woodland roosts and the use of bio-acoustics. J Appl Ecol: 403–410.

Brough T, Bridgman C (1980) An evaluation of long grass as a bird deterrent on British airfields. J Appl Ecol: 243–253.

Brown KM, Erwin RM, Richmond ME, Buckley P, Tanacredi JT, et al. (2001) Managing birds and controlling aircraft in the Kennedy Airport-Jamaica Bay wildlife refuge complex: The need for hard data and soft opinions. Environ Manag 28: 207–224.

Brown R, Gillespie D, Lock A, Pearce P, Watson G (1973) Bird mortality from oil slicks off eastern Canada, February-April 1 970. Can Field Nat 87: 225–234.

Brown WK, Hall WK, Linton LR, Huenefeld RE, Shipley LA (2000) Repellency of three compounds to caribou. Wildlife Soc B: 365–371.

Brownlee BG, Lait S, Madill R, Bunce NJ, Fedorak PM, et al. (1996) Environmental dynamics of base/neutral compounds from oil sands fine tailings. Phase One (1995/1996) Progress Report. NWRI Contribution No. 96-168. 32 p.

Brua RB, Cash KJ, Culp JM (2003) Final Summary Report Task 5: hydrocarbon/oil sands and heavy oil research and development: assessment of natural and anthropogenic impacts of oil sands contaminants within the northern river basins.

Bruder J, Cavo V, Wicks M (1997) Bird hazard detection with airport surveillance radar. International Conference on Radar 1997. pp. 160–163.

Bruderer B (1997) The study of bird migration by radar part 1: The Technical Basis. Naturwissenschaften 84: 1–8.

Bruderer B (1997) The study of bird migration by radar part 2: Major achievements. Naturwissenschaften 84: 45–54.

Bruderer B (1999) Three decades of tracking radar studies on bird migration in Europe and the Middle East. Migrating Birds Know No Boundaries: Proceedings International Seminar on Birds and Flight Safety in the Middle East, Israel. pp. 107–141.

Bruderer B (2007) An Introduction to radar ornithology. International Conference and Workshop on Radar Ornithology and Entomology. Heligoland, 25-28 June 2007.

Bruderer B, Peter D, Boldt A, Liechti F (2010) Wing-beat characteristics of birds recorded with tracking radar and cine camera. Ibis 152: 272–291.

Bruderer B, Peter D, Steuri T (1999) Behaviour of migrating birds exposed to X-band radar and a bright light beam. J Exp Biol 202: 1015–1022.

Bruderer B, Popa-Lisseanu AG (2005) Radar data on wing-beat frequencies and flight speeds of two bat species. Acta Chiropt 7: 73–82.

Bruderer B, Steuri T, Baumgartner M (1995) Short-range high-precision surveillance of nocturnal migration and tracking of single targets. Israel J Zool 41: 207–220.

Brunström B, Broman D, Näf C (1990) Embryotoxicity of polycyclic aromatic hydrocarbons (PAHs) in three domestic avian species, and of PAHs and coplanar polychlorinated biphenyls (PCBs) in the common eider. Environ Pollut 67: 133–143.

Brunström B, Broman D, Näf C (1991) Toxicity and EROD-inducing potency of 24 polycyclic aromatic hydrocarbons (PAHs) in chick embryos. Arch Toxicol 65: 485–489.

Buckland ST, Handel C (2006) Point-transect surveys for songbirds: robust methodologies. Auk 123: 345–357.

Bugnyar T, Stöwe M, Heinrich B (2004) Ravens, Corvus corax, follow gaze direction of humans around obstacles. Roc R Soc Lond B Biol Sci 271: 1331–1336.

Buler JJ, Diehl RH (2009) Quantifying bird density during migratory stopover using weather surveillance radar. IEEE Trans Geosci Remote Sens 47: 2741–2751.

Buler JJ, Moore FR (2011) Migrant-habitat relationships during stopover along an ecological barrier: extrinsic constraints and conservation implications. J Ornithol152: 101–112.

Buler JJ, Moore FR, Woltmann S (2007) A multi-scale examination of stopover habitat use by birds. Ecology 88: 1789–1802.

Burger AE (1993) Estimating the mortality of seabirds following oil spills: effects of spill volume. Mar Pollut Bull 26: 140–143.

Burger AE (1997) Behavior and numbers of marbled murrelets measured with radar. J Field Ornithol: 208–223.

Burger AE, Fry DM (1993) The status, ecology, and conservation of marine birds of the North Pacific. In: Vermeer K, Briggs KT, Morgan KH, Siegel-Causey D, editors. Ottawa: Canadian Wildlife Service Special Publication. pp. 254–263.

Burger J (1997) Effects of oiling on feeding behavior of sanderlings and semipalmated plovers in New Jersey. Condor 99: 290–298.

Burger J, Gochfeld M (1981) Discrimination of the threat of direct versus tangential approach to the nest by incubating herring and great black-backed gulls. J Comp Physiol Psychol 95: 676-684.

Burger J, Tsipoura N (1998) Experimental oiling of sanderlings (*Calidris alba*): Behavior and weight changes. Env Toxicol Chem 17: 1154–1158.

Burgman M, Lindenmayer D, Elith J (2005) Managing landscapes for conservation under uncertainty. Ecology 86: 2007–2017.

Burke C, Montevecchi W, Wiese F (2012) Inadequate environmental monitoring around offshore oil and gas platforms on the Grand Bank of Eastern Canada: Are risks to marine birds known? J Environ Manag 104: 121–126.

Burkhardt D (1982) Birds, berries and UV. Naturwissenschaften 69: 153–157.

Burkhardt D (1989) UV vision: a bird's eye view of feathers. J Comp Physiol A 164: 787–796.

Burrowes A, Marsh R, Teare M, Evans C, Ramos S, et al. (2010) Alberta's Energy Reserves 2009 Supply/Demand Outlook 2010-2019. Energy Resources Conservation Board: Calgary, AB, Canada.

Burrowes, A. et al. 2011. ST98-2011: Alberta's Energy Reserves 2010 and Supply/Demand Outlook 2011-2020. Energy Resources Conservation Board: Calgary, AB, Canada.

Burton AC, Huggard D, Bayne E, Schieck J, Sólymos P, et al. (2014) A framework for adaptive monitoring of the cumulative effects of human footprint on biodiversity. Environ Monit Assess: 1–13.

Butler RG (1988) Field studies on the reproductive effects of oil and emulsion on marine birds. Report. API PUBLICATION NO. 4466. Health and Environmental Sciences Department. Washington, DC, U.S.A.105 p.

Butler RG, Borthwick J, Harmon RS, Leighton FA, Peakall DB (1986) Effects of crude oil exposure on standard metabolic rate of Leach's storm-petrel. Condor 88: 248.

Butler RG, Harfenist A, Leighton FA, Peakall DB (1988) Impact of sublethal oil and emulsion exposure on the reproductive success of Leach's Storm-Petrels: short and long-term effects. J Appl Ecol 25: 125–143.

Butler RG, Lukasiewicz P (1979) A field study of the effect of crude oil on Herring gull (*Larus argentatus*) chick growth. Auk 96: 809–812.

Butt N, Beyer H, Bennett J, Biggs D, Maggini R, et al. (2013) Biodiversity Risks from Fossil Fuel Extraction. Science 342: 425–426.

Butterworth E, Leach A, Gendron M, Pollard B, and Stewart GR (2002) Peace-Athabasca Delta waterbird inventory program: 1998–2001 final report. Ducks Unlimited Canada. Edmonton, Alberta, Canada.

Buxton RT, Major HL, Jones IL, Williams JC (2013) Examining patterns in nocturnal seabird activity and recovery across the Western Aleutian Islands, Alaska, using automated acoustic recording. Auk 130: 331–341.

Cairns D, Elliot R (1987) Oil spill impact assessment for seabirds: the role of refugia and growth centres. Biol Cons 40: 1–9.

Calvert AM, Bishop CA, Elliot RD, Krebs EA, Kydd TM, et al. (2013) A synthesis of human-related avian mortality in Canada. Avian Conserv Ecol 8: 11.

Calvert AM, Taylor PD, Walde S (2009) Cross-scale environmental influences on migratory stopover behaviour. Glob Change Biol 15: 744–759.

Calvert AM, Walde SJ, Taylor PD (2009) Nonbreeding-season drivers of population dynamics in seasonal migrants: conservation parallels across taxa. Avian Conserv Ecol 4: Available: http://www.ace-eco.org/vol4/iss2/art5/

Campbell M, Francis CM (2011) Using stereo-microphones to evaluate observer variation in North American breeding bird survey point counts. Auk 128: 303–312.

Camphuysen C, Fox A, Leopold M, Petersen IK (2004) Towards standardised seabirds at sea census techniques in connection with environmental impact assessments for offshore wind farms in the UK A comparison of ship and aerial sampling methods for marine birds, and their applicability to offshore wind farm assessments. Koninklijk Nederlands Instituut voor Onderzoek der Zee Report commissioned by COWRIE.

Camphuysen C, Heubeck M (2001) Marine oil pollution and beached bird surveys: the development of a sensitive monitoring instrument. Environ Pollut112: 443–461.

Camphuysen CJ (2013) Monitoring and assessment of the proportion of oiled Common Guillemots from beached bird surveys in The Netherlands update winter 2012/13. Report commissioned by the Ministry of Infrastructure and the Environment (I&M), RWS Water, Traffic and Living Environment (RWS-WVL) and DG Mobility and Transport (DGB)

Camphuysen KC (2010) Declines in oil-rates of stranded birds in the North Sea highlight spatial patterns in reductions of chronic oil pollution. Mar Pollut Bull 60: 1299–1306.

Camphuysen KCJ (1998) Beached bird surveys indicate decline in chronic oil pollution in the North Sea. Mar Pollut Bull 36: 519–526.

Candusso JP, Crochet M (2001) Effect of beam broadening on the VHF Doppler mini-radar simple method for correcting wind velocity errors. J Atmos Sol-Terr Phys 63: 275–284.

Canterbury GE, Martin TE, Petit DR, Petit LJ, Bradford DF (2000) Bird communities and habitat as ecological indicators of forest condition in regional monitoring. Conserv Biol 14: 544–558.

Cantox Environmental Inc. (2005) Human Health Risk Assessment. Prepared For: JDEL Associates Ltd. For the Deer Creek energy limited Joslyn North mine project. Project No. 85591.

CAPP (2011) The Facts on Oil Sands. Calgary, Alberta: Canadian Association of Petroleum Producers.

CAPP (2012) Land use in Canada's oil sands. Calgary, Alberta: Canadian Association of Petroleum Producers.

Carlson M, Bayne E, Stelfox B (2008) Assessing the future wildlife impacts of conservation and development in the Mackenzie watershed. Proceedings of the Fourth International Partners in Flight Conference. pp. 13–16.

Carr JM, Lima SL (2010) High wind speeds decrease the responsiveness of birds to potentially threatening moving stimuli. Anim Behav80: 215–220.

Carr JM, Lima SL (2013) Wintering birds avoid warm sunshine: predation and the costs of foraging in sunlight. Oecologia: DOI 10.1007/s00442-013-2804-7

Carter HR (2003) Oil and California's seabirds: an overview. Mar Ornithol 31: 1–7.

Carter J, Lyons NJ, Cole HL, Goldsmith AR (2008) Subtle cues of predation risk: starlings respond to a predator's direction of eye-gaze. Proc R Soc B Biol Sci 275: 1709–1715.

Cassady F (2014 expected) Behavioral responses of birds to deterrents on water bodies and adjacent shores. Master of Science, University of Alberta, Edmonton, Alberta, Canada.

Castege I, Lalanne Y, Gouriou V, Hemery G, Girin M, et al. (2007) Estimating actual seabirds mortality at sea and relationship with oil spills: lesson from the "Prestige" oil spill in Aquitaine (France). Ardeola 54: 289–307.

Castelli PM, Sleggs SE (2000) Efficacy of border collies to control nuisance Canada geese. Wildlife Soc B: 385–392.

Catry P, Lemos R, Brickle P, Phillips R, Matias R, et al. (2013) Predicting the distribution of a threatened albatross: The importance of competition, fisheries and annual variability. Progress in Oceanography 110: 1–10.

Cavanaugh K, Goldsmith A, Holmes W, Follett B (1983) Effects of ingested petroleum on the plasma prolactin levels during incubation and on the breeding success of paired mallard ducks. Arch Environ Contam Toxicol 12: 335–341.

Cavanaugh K, Holmes W (1982) Effects of ingested petroleum on plasma levels of ovarian steroid hormones in photostimulated mallard ducks. Arch Environ Contam Toxicol 11: 503–508.

Cavanaugh K, Holmes W (1987) Effects of ingested petroleum on the development of ovarian endocrine function in photostimulated mallard ducks (*Anas platyrhynchos*). Arch Environ Contam Toxicol 16: 247–253.

CEATAG (1998) Naphthenic Acids Background Information - Discussion Report. Edmonton, Alberta: CONRAD Environmental Aquatics Technical Advisory Group.

Chabanet P, Loiseau N, Join J-L, Ponton D (2012) VideoSolo, an autonomous video system for high-frequency monitoring of aquatic biota, applied to coral reef fishes in the Glorioso Islands (SWIO). J Exp Mar Bio Ecol 430: 10–16.

Chalaturnyk RJ, Scott JD, Özüm B (2002) Management of oil sands tailings. Pet Sci Technol 20: 1025–1046.

Chambers CN (2007) "Well its remote, I suppose, innit?" The relational politics of bird-watching through the CCTV lens. Scottish Geogr J 123: 122–134.

Chen W, Ning H, Li J (2011) Flying bird detection and hazard assessment for avian radar system. J Aerosp Eng 25: 246–255.

Cheskey ED, Wells JV, Casey-Lefkowitz S (2011) Birds at Risk: The Importance of Canada's Boreal Wetlands and Waterways. New York City, New York: Natural Resources Defense Council.

Chevallier D, Handrich Y, Georges JY, Baillon F, Brossault P, et al. (2010) Influence of weather conditions on the flight of migrating black storks. Proc R Soc B Biol Sci 277: 2755–2764.

Chilson PB, Frick WF, Kelly JF, Howard KW, Larkin RP, et al. (2012) Partly cloudy with a chance of migration: weather, radars, and aeroecology. Bull Am Meteorol Soc 93: 669–686.

Chipman RB, Dollbeer RA, Pruesser KJ, Sullivan DP, Losito ED, et al. (2004) Emergency wildlife management response to protect evidence associated with the terrorist attack on the World Trade Center, New York City. USDA National Wildlife Res Center-Staff Publications: 86.

Ciborowski JJH, Whelly MP (1997) Chironomidae as Indicators of Wetland Viability: Report on Field Work in Wetlands of the Fort McMurray, Alberta Area (June 15 - July 23, 1997)).

Clark L (1998) Review of bird repellents. Proceedings of the Eighteenth Vertebrate Pest Conference (1998). p. 6.

Clark R (1970) Oiled seabird research unit. Mar Pollut Bull 1: 22–24.

Clark R (1984) Impact of oil pollution on seabirds. Environ Pollut A Ecol Biol 33: 1–22.

Clark R, Dunnet G, Addy J (1984) Seabirds and north sea oil. Mar Pollut Bull 15: 272–274.

Clark WS, Gorney E (1987) Oil contamination of raptors migrating along the Red Sea. Environ Pollut 46: 307–313.

Cleaning and Rehabilitation of Oiled Wildlife (2007). Workshop Proceedings. Algarve, 21-24 October 2006. 24 p.

Clemente JS, Fedorak PM (2005) A review of the occurrence, analyses, toxicity, and biodegradation of naphthenic acids. Chemosphere 60: 585–600.

Clemente JS, Prasad N, MacKinnon MD, Fedorak PM (2003) A statistical comparison of naphthenic acids characterized by gas chromatography-mass spectrometry. Chemosphere 50: 1265–1274.

Clive Welham Forr (2013) Factors affecting ecological resilience of reclaimed oil sands uplands. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-34. 44 pp.

Cobb JLS (1976) Seabird mortality. Bird Study 23: 299–300.

Cochran WW, Graber RR (1958) Attraction of nocturnal migrants by lights on a television tower. Wilson Bull 70: 378–380.

Cochran WW, Mouritsen H, Wikelski M (2004) Migrating songbirds recalibrate their magnetic compass daily from twilight cues. Science 304: 405–408.

Cohen B, Williams TC (1980) Short-range corrections for migrant bird tracks on search radars. J Field Ornithol 51: 248–253.

Colavecchia M (2006) Toxicity of oil sands to the early life stages of fish. Doctor of Philosophy. Queen's University, Kingston, ON, Canada.

Colavecchia MV, Backus SM, Hodson PV, Parrott JL (2004) Toxicity of oil sands to early life stages of fathead minnows (*Pimephales promelas*). Environ Toxicol Chem 23: 1709–1718.

Colavecchia MV, Hodson PV, Parrott JL (2006) CYP1A induction and blue sac disease in early life stages of white suckers (*Catostomus commersoni*) exposed to oil sands. J Toxicol Environ Health A 69: 967–994.

Colavecchia MV, Hodson PV, Parrott JL (2007) The relationships among CYP1A induction, toxicity, and eye pathology in early life stages of fish exposed to oil sands. J Toxicol Environ Health A 70: 1542–1555.

Collinge WE (1921) The effects of oil from ships on certain sea-birds. Nature 106: 830.

Collins R, Brazier H, Whelan J (1994) Rehabilitating a herd of oiled mute swans *Cygnus olor*. Biol Environ: Proc R Ir Acad. pp. 83–89.

Colwell RR, Walker JD, Cooney JJ (1977) Ecological aspects of microbial degradation of petroleum in the marine environment. Crit Rev Microbiol5: 423–445.

Committee TOSVR (1998) Guidelines for Reclamation to Forest Vegetation in the Alberta Oil Sands Region. Oil Sands Vegetation Reclamation Committee. Edmonton, AB, Canada. 57 p.

Conover M (1999) Can waterfowl be taught to avoid food handouts through conditioned food aversions? Wildlife Soc B: 160–166.

Conover MR (2010) Fear-provoking stimuli. In: Resolving human-wildlife conflicts: the science of wildlife damage management. Boca Raton: CRC press. pp. 229-247.

Conover MR, Chasko GG (1985) Nuisance Canada goose problems in the eastern United States. Wildlife Soc B 13: 228–233.

Cook A, Rushton S, Allan J, Baxter A (2008) An evaluation of techniques to control problem bird species on landfill sites. Environ Manag 41: 834–843.

Coon NC, Albers PH, Szaro RC (1979) No. 2 fuel oil decreases embryonic survival of great black-backed gulls. Bull Environ Contam Toxicol 21: 152–156.

Coon NC, Dieter MP (1981) Responses of adult mallard ducks to ingested South Louisiana crude oil. Environ Res 24: 309–314.

Cooper BA, Day RH, Ritchie RJ, Cranor CL (1991) An improved marine radar system for studies of bird migration. J Field Ornithol 62: 367–377.

Cooper BA, Ritchie RJ (1995) The altitude of bird migration in east-central Alaska: a radar and visual study. J Field Ornithol 66: 590–608.

Cooper Jr WE (2008) Visual monitoring of predators: occurrence, cost and benefit for escape. Anim Behav 76: 1365–1372.

Cooper Jr WE, Frederick WG (2007) Optimal flight initiation distance. J Theor Biol 244: 59–67.

Cooper WE (2006) Dynamic risk assessment: prey rapidly adjust flight initiation distance to changes in predator approach speed. Ethology 112: 858–864.

Cooper WE (2009) Fleeing and hiding under simultaneous risks and costs. Behav Ecol 20: 665–671.

Copeland HE, Pocewicz A, Kiesecker JM (2011) Geography of energy development in western North America: potential impacts on terrestrial ecosystems. Energy Development and Wildlife Conservation in Western North America. Springer. pp. 7–22.

Corilo YE, Podgorski DC, McKenna AM, Lemkau KL, Reddy CM, et al. (2013) Oil spill source identification by principal component analysis of electrospray ionization Fourier transform ion cyclotron resonance mass spectra. Anal Chem 85: 9064–9069.

Corre ML, Ollivier A, Ribes S, Jouventin P (2002) Light-induced mortality of petrels: a 4-year study from Reunion Island (Indian Ocean). Biol Cons 105: 93–102.

Costanza R, d' Arge R, De Groot R, Farber S, Grasso M, et al. (1997) The value of the world's ecosystem services and natural capital. Nature 387: 253–260.

Couillard C, Leighton F (1990) Sequential study of the pathology of Prudhoe Bay crude oil in chicken embryos. Ecotoxicol Environ Saf 19: 17–23.

Couillard C, Leighton F (1990) The toxicopathology of Prudhoe Bay crude oil in chicken embryos. Fundam Appl Toxicol14: 30–39.

Couillard CM, Leighton FA (1991) Bioassays for the toxicity of petroleum oils in chicken embryos. Environ Toxicol Chem 10: 533–538.

Couillard CM, Leighton FA (2009) Critical period of sensitivity to petroleum toxicity in the chicken embryo. Environ Toxicol Chem 10: 249–253.

Coulton DW, Virgl JA, English C (2013) Falcon nest occupancy and hatch success near two diamond mines in the Southern Arctic, Northwest Territories. Avian Conserv Ecol 8: 14.

Crawford RJM, Davis SA, Harding RT, Jackson LF, Leshoro TM, et al. (2000) Initial impact of the Treasure oil spill on seabirds off western South Africa. S Afr J Mar Sci 22: 157–176.

Crawford RL, Engstrom RT (2001) Characteristics of avian mortality at a north Florida television tower: a 29-year study. J Field Ornithol 72: 380–388.

Creasey R (2012) Professional judgment in mineable oil sands reclamation certification: workshop summary. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-25. 52 pp.

Cresswell W (1993) Escape responses by redshanks, *Tringa totanus*, on attack by avian predators. Anim Behav 46: 609–611.

Cresswell W, Hilton GM, Ruxton GD (2000) Evidence for a rule governing the avoidance of superfluous escape flights. Roc R Soc Lond B Biol Sci 267: 733–737.

Crocker AD, Cronshaw J, Holmes W (1974) The effect of a crude oil on intestinal absorption in ducklings (*Anas platyrhynchos*). Environ Pollut 7: 165–177.

Cruz-Martinez L, Smits JEG (2012) Potential to use animals as monitors of ecosystem health in the oil sands region. Report No. TR-18; OSRIN: University of Alberta, Edmonton, Alberta, Canada.

Culhane PJ (1987) The precision and accuracy of US environmental impact statements. Environ Monit Assess 8: 217–238.

Culik BM, Wilson RP, Woakes AT, Sanudo FW (1991) Oil pollution of Antarctic penguins: effects on energy metabolism and physiology. Mar Pollut Bull 22: 388–391.

Cullen S, Robertson GJ (1999) Temporal changes in the anti-predator flocking behavior of wintering shorebirds. Waterbirds 22: 156–159.

Cumming GS (2001) Using between-model comparisons to fine-tune linear models of species ranges. J Biogeogr 27: 441–455.

Cumming SG, Lefevre KL, Bayne E, Fontaine T, Schmiegelow FKA, et al. (2010) Toward conservation of canada's boreal forest avifauna: design and application of ecological models at continental extents. Avian Conserv Ecol 5: 8.

Cunningham DJ, Anderson WH, Anthony RM (1996) An image-processing program for automated counting. Wildlife Soc B 24: 345–346.

Custer T, Bickham J, Lyne T, Lewis T, Ruedas L, et al. (1994) Flow cytometry for monitoring contaminant exposure in black-crowned night-herons. Arch Environ Contam Toxicol 27: 176–179.

Custer T, Custer C, Hines R, Sparks D, Melancon M, et al. (2000) Mixed-function oxygenases, oxidative stress, and chromosomal damage measured in lesser scaup wintering on the Indiana Harbor Canal. Arch Environ Contam Toxicol 38: 522–529.

Custer TW, Albers PH (1980) Response of captive, breeding mallards to oiled water. The J Widlife Manag: 915–918.

Custer TW, Custer CM, Dickerson K, Allen K, Melancon MJ, et al. (2001) Polycyclic aromatic hydrocarbons, aliphatic hydrocarbons, trace elements, and monooxygenase activity in birds nesting on the North Platte River, Casper, Wyoming, USA. Environ Toxicol Chem 20: 624–631.

Cuthill IC, Partridge JC, Bennett AT, Church SC, Hart NS, et al. (2000) Ultraviolet vision in birds. Adv Study Behav 29: 159–214.

Cutler TL, Morrison ML (1998) Habitat use by small vertebrates at two water developments in southwestern Arizona. Southwest Nat 43: 155–162.

Cutler TL, Swann DE (1999) Using remote photography in wildlife ecology: a review. Wildlife Soc B 27: 571–581.

D'Andrea MA, Reddy GK (2013) Health consequences among subjects involved in gulf oil spill clean-up activities. Am J Med 126: 966–974.

Dagenais L (2008) Effects of oil sands extraction activities on breeding birds in the Athabasca oil sands region. Master of Science, University of Alberta, Edmonton, Alberta, Canada.

Damstra T, Barlow S, Bergman A, Kavlock R, Van Der Kraak G (2002) Global assessment of the state-of-the-science of endocrine disruptors. Global assessment of the state-of-the-science of endocrine disruptors. World Health Organization.

Dangles O, Irschick D, Chittka L, Casas J (2009) Variability in sensory ecology: expanding the bridge between physiology and evolutionary biology. Q Rev Biol 84: 51–74.

Danz NP, Regal RR, Niemi GJ, Brady VJ, Hollenhorst T, et al. (2005) Environmentally stratified sampling design for the development of Great Lakes environmental indicators. Environ Monit Assess 102: 41–65.

Davies J, Eaton B (2011) Community level physiological profiling for monitoring oil sands impacts. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-10. 44 pp.

Davies JBE, Humphries D (2012) Microcosm evaluation of community level physiological profiling in oil sands process affected water. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-28. 33 pp.

Davis J, Farmer V, Kreider R, Straub A, Reese K (1972) Migration of petroleum products in soil and ground water: principles and countermeasures. API Publication No. 4149.

Davis MA, Chew MK, Hobbs RJ, Lugo AE, Ewel JJ, et al. (2011) Don't judge species on their origins. Nature 474: 153–154.

Dawood M, Narayanan RM (2001) Receiver operating characteristics for the coherent UWB random noise radar. IEEE Trans Aerosp Electron Syst 37: 586–594.

Dawson DK, Efford MG (2009) Bird population density estimated from acoustic signals. J Appl Ecol 46: 1201–1209.

Day RH, Murphy SM, Wiens JA, Hayward GD, Harner EJ, et al. (1997) Effects of the Exxon Valdez oil spill on habitat use by birds along the Kenai Peninsula, Alaska. Condor 99: 728–742.

Day RH, Murphy SM, Wiens JA, Hayward GD, Harner EJ, et al. (1997) Effects of the Exxon Valdez oil spill on habitat use by birds in Prince William Sound, Alaska. Ecol Appl 7: 593–613.

De Jager S, Belterman T (1970) Treatment of oiled eider duck in Holland. Mar Pollut Bull 1: 156.

De Jong A, Magnhagen C, Thulin C-G (2013) Variable flight initiation distance in incubating Eurasian curlew. Behav Ecol and Sociobiology: 1–8.

De Lucas M, Janss GF, Ferrer M (2004) The effects of a wind farm on birds in a migration point: the Strait of Gibraltar. Biodivers Conserv 13: 395–407.

Deer Creek Energy Ltd. (2006) Joslyn North Mine project: Alberta Energy and Utilities Board Alberta Environment integrated application.

DeFusco RP (2007) A successful case study: The bird control program of Waste Management Outer Loop Recycling and Disposal Facility, Louisville, Kentucky, USA.Bird Strike Committee USA/Canada, 9th Annual Meeting, Kingston, Ontario.Paper 5. Available: http://digitalcommons.unl.edu/birdstrike2007/5

Degernes LA (1995) Toxicities in waterfowl. Seminars in Avian and Exotic Pet Medicine. 4:15–22.

Degernes LA (2008) Waterfowl toxicology: a review. Veterinary Clinics of North America: Exotic Animal Practice *VCONA/EAP* 11: 283–300.

Delaney DK, Grubb TG, Garcelon DK (1998) An infrared video camera system for monitoring diurnal and nocturnal raptors. J Raptor *Res* 32: 290–296.

Descamps S, Béchet A, Descombes X, Arnaud A, Zerubia J (2011) An automatic counter for aerial images of aggregations of large birds. Bird Study 58: 302–308.

Desholm M (2009) Avian sensitivity to mortality: Prioritising migratory bird species for assessment at proposed wind farms. J Environ Manag 90: 2672–2679.

Desholm M, Fox A, Beasley P, Kahlert J (2006) Remote techniques for counting and estimating the number of bird-wind turbine collisions at sea: a review. Ibis 148: 76–89.

Desholm M, Laursen K (2005) Preliminary investigations of bird-turbine collisions at Nysted offshore wind farm and final quality control of Thermal Animal Detection System (TADS). Report commissioned by Energi E2. National Environmental Research Institute, Ministry of the Environment. Denmark.

Devenny DW (2010) A Screening Study of Oil Sand Tailings Technologies and Practices. Prepared for Alberta Energy Research Institute. AERI Contract 2008 0326.

Devenny DW (2010) A Screening Study of Oil Sands Tailings Technologies and Practices - Part B Report. Prepared for Alberta Energy Research Institute. AERI Contract 2008 0326.

Devereux CL, Whittingham MJ, Fernández-Juricic E, Vickery JA, Krebs JR (2006) Predator detection and avoidance by starlings under differing scenarios of predation risk. Behav Ecol 17: 303–309.

Di Lazzaro P, Murra D, Schwortz B (2012) Pattern recognition after image processing of low-contrast images, the case of the Shroud of Turin. Pattern Recognition.46: 1964-1970

Diaz-Gómez D, Toxopeus A, Groen T, Muñoz A, Skidmore A, et al. (2013) Measuring the Insecurity Index of species in networks of protected areas using species distribution modeling and fuzzy logic: The case of raptors in Andalusia. Ecol Indic 26: 174–182.

Diefenbach DR, Brauning DW, Mattice JA, Thompson III F (2003) Variability in grassland bird counts related to observer differences and species detection rates. Auk 120: 1168–1179.

Diehl RH, Larkin RP (2002) Introduction to the WSR-88D (NEXRAD) for ornithological research. Bird conservation implementation and integration in the Americas. Proceedings of the Third International Partners in Flight Conference. pp. 20–24.

Diehl RH, Larkin RP, Black JE, Moore F (2003) Radar observations of bird migration over the Great Lakes. Auk 120: 278–290.

Dieter MP (1977) Acute and chronic studies with waterfowl exposed to petroleum hydrocarbons. Program Review Proceedings Of: Environmental Effects of Energy Related Activities On Marine/Esturine Ecosytems. OEMI-77-025.

Dinata Y, Nugroho A, Haidir IA, Linkie M (2008) Camera trapping rare and threatened avifauna in west-central Sumatra. Bird Conserv Int 18: 30.

Dipper B (1998) Monitoring and post-auditing in environmental impact assessment: a review. J Environ Plan Manag 41: 731–747.

Dixon DG, Farwell AJ (2002) The use of stable isotopes (13C/a_Cand 15N/14N) to trace energy sources and trophic interactions in aquatic systems influenced by mining of the Athabasca Oil Sands. Final report prepared for Syncrude Canada Ltd., Grant #: E3166, University of Waterloo. Waterloo, ON, Canada.

Dixon R, Maier M, Sandilya A, Schneider T (2012) Qualifying environmental trusts as financial security for oil sands reclamation liabilities. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-24. 32 pp.

Dokter AM, Baptist MJ, Ens BJ, Krijgsveld KL, van Loon EE (2013) Bird radar validation in the field by time-referencing line-transect surveys. PLoS ONE 8: e74129.

Dokter AM, Liechti F, Stark H, Delobbe L, Tabary P, et al. (2011) Bird migration flight altitudes studied by a network of operational weather radars. J R Soc Interface 8: 30–43.

Dokter AM, Shamoun-Baranes J, Kemp MU, Tijm S, Holleman I (2013) High altitude bird migration at temperate latitudes: A synoptic perspective on wind assistance. PLoS ONE 8: e52300.

Dolan RW, Carter T, Ryan T, Salsbury C, Dolan TE, et al. (2013) Trials of the urban ecologist. Front Ecol Environ 11: 163–164.

Dolbeer RA (2009) Birds and aircraft—fighting for airspace in ever more crowded skies. Human–Wildlife Interactions. Paper 10. Available: http://digitalcommons.unl.edu/hwi/10

Dominguez J, Vidal M (2009) Factors affecting plumage oiling levels in a Kentish Plover Charadrius alexandrinus population after a major oil spill. Acta Ornithologica 44: 119–126.

Domjan M (2003) The principles of learning and behaviour. 5th ed. Belmont, CA.: Thompson-Wadsworth.

Douben PE (2003) PAHs: an ecotoxicological perspective. Chichester: John Wiley & Sons Ltd. 388 p.

Dowdeswell E (2011) A Foundation for the Future: Building an Environmental Monitoring System for the Oil Sands. Environment Canada. Ottawa, On, Canada.

Drake V, Harman I, Wang H (2002) Insect monitoring radar: stationary-beam operating mode. Computers and Electronics in Agriculture 35: 111–137.

Draulans D (1987) The effectiveness of attempts to reduce predation by fish-eating birds: a review. Biol Cons 41: 219–232.

Draulans D, Van Vessem J (1985) The effect of disturbance on nocturnal abundance and behaviour of grey herons (*Ardea cinerea*) at a fish-farm in winter. J Appl Ecol: 19–27.

Drewitt AL, Langston RHW (2006) Assessing the impacts of wind farms on birds. Ibis 148: 29–42.

Duffy DC (1977) Incidence of oil contamination on breeding Common Terns. Bird-Banding 48: 370–371.

Duffy L, Bowyer RT, Testa JW, Faro JB (1993) Differences in blood haptoglobin and length-mass relationships in river otters (*Lutra canadensis*) from oiled and nonoiled areas of Prince William Sound, Alaska. J Wildl Dis 29: 353–359.

Duffy LK, Bowyer RT, Testa JW, Faro JB (2009) Chronic effects of the Exxon Valdez oil spill on blood and enzyme chemistry of river otters. Environ Toxicol Chem 13: 643–647.

Duffy LK, Hecker MK, Blundell GM, Bowyer RT (1999) An analysis of the fur of river otters in Prince William Sound, Alaska: oil related hydrocarbons 8 years after the Exxon Valdez oil spill. Polar Biol 21: 56–58.

Duguay AM, Davies RW (1997) The chronic ecotoxicology of oil sands tailings on freshwater invertebrates. Calgary, Alberta: Department of Biological Sciences, University of Calgary.

Dumont F, Pasquaretta C, Réale D, Bogliani G, Hardenberg A (2012) Flight initiation distance and starting distance: biological effect or mathematical artefact? Ethology 118: 1051–1062.

Dunn EH, Larivée J, Cyr A (2001) Site-specific observation in the breeding season improves the ability of checklist data to track population trends. J Field Ornithol: 547–555.

Dunnet G, Crisp D, Conan G, Bourne W (1982) Oil pollution and seabird populations [and discussion]. Philos Trans R Soc Lond B Biol Sci 297: 413–427.

Dunnet G, Samiullah Y, Stone D, Tasker M, Dunnet G, et al. (1987) Seabirds and north sea oil [and discussion]. Philos Trans R Soc Lond B Biol Sci 316: 513–524.

Dyer SJ, O'Neill JP, Wasel SM, Boutin S (2002) Quantifying barrier effects of roads and seismic lines on movements of female woodland caribou in northeastern Alberta. Canadian J Zool 80: 839–845.

Eason PK, Sherman PT, Rankin O, Coleman B (2006) Factors affecting flight initiation distance in American robins. J Widlife Manag 70: 1796–1800.

Eastin Jr WC, Murray HC (1981) Effects of crude oil ingestion on avian intestinal function. Can J Physiol Pharmacol 59: 1063–1068.

Eastin W, Hoffman D (1978) Biological effects of petroleum on aquatic birds. Proceedings of the conference on assessment of ecological impacts of oil spills. American Institute for Biological Sciences, Washington, DC. pp. 561–582.

Eastin WC Jr. (1979) Methods used at Patuxent Wildlife Research Center to study the effects of oil on birds. Proceedings of the 1979 U.S. Fish and Wildlife Service Pollution Response Workshop. St. Petersburg, Florida. pp. 60–66.

Eastin WC, Hoffman DJ, O'Leary CT (1983) Lead accumulation and depression of delta-aminolevulinic acid dehydratase (ALAD) in young birds fed automotive waste oil. Arch Environ Contam Toxicol 12: 31–35.

Eastin WC, Rattner BA (1982) Effects of dispersant and crude oil ingestion on mallard ducklings (*Anas platyrhynchos*). Bull Environ Contam Toxicol 29: 273–278.

Eastwood E (1967) Radar techniques for detection, tracking and navigation. Nature 213: 1072.

Eaton BTMJTF, Chai S-L (2013) Potential impacts of beaver on oil sands reclamation success – an analysis of available literature. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-37. 65 pp.

Eichholz MW, Dassow JA, Stafford JD, Weatherhead PJ (2012) Experimental evidence that nesting ducks use mammalian urine to assess predator abundance. Auk 129: 638–644.

Eisler R (1987) Polycyclic aromatic hydrocarbon hazards to fish, wildlife, and invertebrates. A Synoptic Review. Biological Report 85 (1.11); U.S. Fish and Wildlife Service: Laurel, MD, U.S.A.

Eisler R (2000) Handbook of Chemical Risk Assessment: Health Hazards to Humans, Plants, and Animals; Lewis Publishers: Boca Raton, U.S.A.

Elkins N (1983) Weather and Bird Behaviour. Calton, Staffordshire: T&AD Pyser Ltd.

Elkins N (2004) Weather and Bird Behaviour (3rd Edition). London, GBR: A & C Black.

Ellenton JA (1982) Teratogenic activity of aliphatic and aromatic fractions of Prudhoe Bay crude and fuel oil No. 2 in the chicken embryo. Toxicol Appl Pharmacol 63: 209–215.

Ellis JI, Wilhelm SI, Hedd A, Fraser GS, Robertson GJ, et al. (2009) Mortality of migratory birds from marine commercial fisheries and offshore oil and gas production in Canada. Avian Conserv Ecol 8: 4.

Elphick C (2008) Editor's choice: New research on wind farms. J Appl Ecol 45: 1840–1840.

Elphick CS (2008) How you count counts: the importance of methods research in applied ecology. J Appl Ecol 45: 1313–1320.

Elvidge CD, Keith DM (2010) Spectral signatures of night time lights. In: Art, Science and Applications of Reflectance Spectroscopy Symposium, Boulder.

Elzinga CL, Salzer DW, Willoughby JW, Gibbs JP (2001) Monitoring plant and animal populations. Blackwell Science, Oxford, UK.

Emery N (2000) The eyes have it: the neuroethology, function and evolution of social gaze. Neurosci Biobehav Rev 24: 581–604.

Energy and Natural Resources, Energy Resources Conservation Board, Alberta Oil Sands Technology and Research Authority (1977) The Alberta Oil Sands. ENR Report No. 26. Edmonton, Alberta: Alberta Energy and Natural Resources.

Englander JG, Bharadwaj S, Brandt AR (2013) Historical trends in greenhouse gas emissions of the Alberta oil sands (1970-2010). Environ Res Lett 8: 044036.

Environment Canada (2001) Assessment of Natural and Anthropogenic Impacts of Oil Sands Contaminants within the Northern River Basins. Progress Report for PERD Project No. 521006

Environment Canada (2009) Peace-Athabasca Delta, Alberta - Ramsar Site. Available: http://www.mb.ec.gc.ca/nature/whp/ramsar/df02s06.en.html. Accessed 15 February 2014.

Environment Canada, Government of Alberta (2012) Joint Canada/Alberta Implementation Plan of Oil Sands Monitoring. Edmonton, Alberta: Environment and Sustainable Resource Development.

EPA (1999) Understanding Oil Spills and Oil Spill Response. Available: http://www.epa.gov/osweroe1/content/learning/pdfbook.htm.

Eppley Z (1992) Assessing indirect effects of oil in the presence of natural variation: The problem of reproductive failure in South Polar Skuas during the Bahia Paraiso oil spill. Mar Pollut Bull 25: 307–312.

Eppley ZA, Rubega MA (1989) Indirect effects of an oil spill. Nature 340: 513.

Eppley ZA, Rubega MA (1990) Indirect effects of an oil spill: Reproductive failure in a population of south Polar skuas following the" Bahia Paraiso" oil spill in Antarctica. Mar Ecol Prog Ser 67: 1–6.

Erasmus T, Randall R, Randall B (1981) Oil pollution, insulation and body temperatures in the jackass penguin Spheniscus demersus. Comp Biochem Physiol A 69: 169–171.

ERCB (2012) ST98-2012 Alberta's Energy Reserves 2011 and Supply/Demand Outlook 2012—2021.

ERCB (2013) 2012 tailings management assessment report: oil sands mining industry. Energy Resources Conservation Board. Calgary, AB, Canada.

ERCB (2013) ST98-2013 Alberta's Energy Reserves 2012 and Supply/Demand Outlook 2013–2022.

Erickson WA, Marsh RE, Salmon TP (1992) High frequency sound devices lack efficacy in repelling birds. Proceedings of the Fifteenth Vertebrate Pest Conference 1992. Paper 26. Available: http://digitalcommons.unl.edu/vpc15/26

Erlich N, Lipp OV, Slaughter V (2013) Of hissing snakes and angry voices: human infants are differentially responsive to evolutionary fear-relevant sounds. Develop Sci. DOI: 10.1111/desc.12091.

Erwin RM, Smith GJ, Clapp RB (1986) Winter distribution and oiling of Common Terns in Trinidad: a further look. J Field Ornithol 57: 300–308.

Esler D (2000) Harlequin duck demography during winter in Prince William Sound, Alaska: effects of the Exxon Valdez oil spill. Doctor of Philosophy, Oregon State University, Oregon, U.S.A..

Esler D, Bowman TD, Trust KA, Ballachey BE, Dean TA, et al. (2002) Harlequin duck population recovery following the 'Exxon Valdez' oil spill: Progress, process and constraints. Mar Ecol Prog Ser 241: 271–286.

Esler D, Trust KA, Ballachey BE, Iverson SA, Lewis TL, et al. (2010) Cytochrome P4501A biomarker indication of oil exposure in harlequin ducks up to 20 years after the Exxon Valdez oil spill. Environ Toxicol Chem 29: 1138–1145.

Evans CS, Macedonia JM, Marler P (1993) Effects of apparent size and speed on the response of chickens, Gallus gallus, to computer-generated simulations of aerial predators. Anim Behav 46: 1–11.

Evans JE, Cuthill IC, Bennett AT (2006) The effect of flicker from fluorescent lights on mate choice in captive birds. Anim Behav72: 393–400.

Evans M, Symens P, Pilcher C (1993) Short-term damage to coastal bird populations in Saudi Arabia and Kuwait following the 1991 Gulf War marine pollution. Mar Pollut Bull 27: 157–161.

Evans MS, Talbot A (2012) Investigations of mercury concentrations in walleye and other fish in the Athabasca River ecosystem with increasing oil sands developments. J Environ Monit 14: 1989–2003.

Evans Ogden LJ (1996) Collision course: the hazards of lighted structures and windows to migrating birds. Prepared for World Wildlife Fund Canada and the Fatal Light Awareness Program.

EVS Consultants (1986) Biological Effects Study of Dredged Material Discharge to the Athabasca River near Fort McMurray, Alberta. Prepared for Suncor Inc. North Vancouver, British Columbia: EVS Environment Consultants Limited.

EVS Consultants (1990) Water quality inventory of tailings ponds and assessment of in situ experiment tanks. Prepared for Suncor Inc. North Vancouver, British Columbia: EVS Environment Consultants Limited.

EVS Consultants (1992) Chemical and Toxicological Characteristics of Interstitial Water and Fine Tails. Prepared for Suncor Inc. North Vancouver, British Columbia: EVS Environment Consultants Limited.

EVS Consultants (1992) Constructed Wetlands for the Treatment of Oil Sands Waste Water Technical Report #1. Prepared for Suncor Inc. North Vancouver, British Columbia: EVS Environment Consultants Limited.

EVS Consultants (1992) Microbiological aspects of Suncor's reclamation program. Prepared for Suncor Inc. North Vancouver, British Columbia: EVS Environment Consultants Limited.

EVS Consultants (1992) Natural Wetlands for the Treatment of Leachate from Oil Sands Tailings Ponds Technical Report #1. Prepared for Suncor Inc. North Vancouver, British Columbia: EVS Environment Consultants Limited.

EVS Consultants (1993) A field-scale study of Suncor's sustainable pond development research technical report #1. Prepared for Suncor Inc. North Vancouver, British Columbia: EVS Environment Consultants Limited.

EVS Consultants (1993) Natural wetlands for the treatment of dyke drainage water - technical report #2. Prepared for Suncor Inc. North Vancouver, British Columbia: EVS Environment Consultants Limited.

EVS Consultants (1994) A field-scale study of Suncor's sustainable pond development research technical report #2. Prepared for Suncor Inc. North Vancouver, British Columbia: EVS Environment Consultants Limited.

EVS Consultants (1994) Constructed Wetlands for the Treatment of Oil Sands Waste Water Technical Report #3. Prepared for Suncor Inc. North Vancouver, British Columbia: EVS Environment Consultants Limited.

EVS Consultants (1995) Constructed Wetlands for the Treatment of Oil Sands Waste Water Technical Report #4. Prepared for Suncor Inc. North Vancouver, British Columbia: EVS Environment Consultants Limited.

EVS Consultants (1995) Constructed wetlands for the treatment of oil sands wastewater - Technical Report #4 Appendices. Prepared for Suncor Inc. North Vancouver, British Columbia: EVS Environment Consultants Limited.

EVS Consultants (1995) Wetlands design criteria for treatment of dyke drainage water. Prepared for Suncor Inc. North Vancouver, British Columbia: EVS Environment Consultants Limited.

EVS Consultants (1996) Constructed Wetlands for the Treatment of Oil Sands Waste Water Technical Report #5. Prepared for Suncor Inc. North Vancouver, British Columbia: EVS Environment Consultants Limited.

EVS Consultants (1996) Suncor Lease 86: Baseline Aquatic Monitoring. Prepared for Suncor Inc. North Vancouver, British Columbia: EVS Environment Consultants Limited.

Faaborg J, Holmes RT, Anders AD, Bildstein KL, Dugger KM, et al. (2010) Recent advances in understanding migration systems of New World land birds. Ecoll Monogr 80: 3–48.

Fairbrother A, Smits J, Grasman KA (2004) Avian immunotoxicology. J Toxicol Environ Health B 7: 105–137.

Farmer A, Durbian F (2006) Estimating shorebird numbers at migration stopover sites. Condor 108: 792–807.

Farnsworth A (2005) Flight calls and their value for future ornithological studies and conservation research. Auk 122: 733–746.

Farnsworth A, Russell RW (2007) Monitoring flight calls of migrating birds from an oil platform in the northern Gulf of Mexico. J Field Ornithol 78: 279–289.

Farquharson J, Faszer C (2005) Noise Impact Assessment - Deer Creek Energy Limited Joslyn North Mine. Prepared for Millennium EMS Solutions. Calgary, Alberta: Faszer Farquharson & Associates Ltd.

Farrell A, Kennedy C, Kolok A (2004) Effects of wastewater from an oil-sand-refining operation on survival, hematology, gill histology, and swimming of fathead minnows. Canadian J Zool 82: 1519–1527.

Farwell A, Harms N, Smits J, Dixon D (2014) Stable nitrogen isotopes of nestling tree swallows indicate exposure to different types of oil sands reclamation. J Toxicol Environ Health A 77: 415–425.

Favreau-Peigné A, Calandreau L, Constantin P, Gaultier B, Bertin A, et al. (2014) Emotionality Modulates the Effect of Chronic Stress on Feeding Behaviour in Birds. PLoS ONE 9: e87249.

Ferguson EL, Gilmer DS (1980) Small-format cameras and fine-grain film used for waterfowl population studies. The J Widlife Manag 44: 691–694.

Fernández-Juricic E (2012) Sensory basis of vigilance behavior in birds: synthesis and future prospects. Behav Processes 89: 143–152.

Fernández-Juricic E, Blumstein DT, Abrica G, Manriquez L, Adams LB, et al. (2006) Relationships of anti-predator escape and post-escape responses with body mass and morphology: a comparative avian study. Evol Ecol Res 8: 731–752.

Fernández-Juricic E, Deisher M, Stark AC, Randolet J (2012) Predator detection is limited in microhabitats with high light intensity: an experiment with brown-headed cowbirds. Ethology 118: 341–350.

Fernández-Juricic E, Gaffney J, Blackwell BF, Baumhardt P (2011) Bird strikes and aircraft fuselage color: a correlational study. Hum-Wildl Interact 5: 224–234.

Fernández-Juricic E, Gall MD, Dolan T, O'Rourke C, Thomas S, et al. (2011) Visual systems and vigilance behaviour of two ground-foraging avian prey species: white-crowned sparrows and California towhees. Anim Behav81: 705–713.

Fernández-Juricic E, Jimenez MD, Lucas E (2001) Alert distance as an alternative measure of bird tolerance to human disturbance: implications for park design. Environ Conserv 28: 263–269.

Fernández-Juricic E, Kowalski V (2011) Where does a flock end from an information perspective? A comparative experiment with live and robotic birds. Behav Ecol 22: 1304–1311.

Ferrari MCO, Elvidge CK, Jackson CD, Chivers DP, Brown GE (2010) The responses of prey fish to temporal variation in predation risk: sensory habituation or risk assessment? Behav Ecol 21: 532–536.

Ferrer M, de Lucas M, Janss GF, Casado E, Munoz AR, et al. (2012) Weak relationship between risk assessment studies and recorded mortality in wind farms. J Appl Ecol 49: 38–46.

Field R, North MR, Wells J (1993) Nesting activity of yellow-billed loons on the Colville River Delta, Alaska, after the Exxon Valdez Oil Spill. Wilson Bull: 325–332.

Field SA, O'Connor PJ, Tyre AJ, Possingham HP (2007) Making monitoring meaningful. Austral Ecol 32: 485–491.

Fielding AH, Bell JF (1997) A review of methods for the assessment of prediction errors in conservation presence/absence models. Environ Conserv 24: 38–49.

Fifield D, Lewis K, Gjerdrum C, Robertson G, Wells R (2009) Offshore seabird monitoring program. Environment Studies Research Funds Report 183.

Finch BE, Wooten KJ, Faust DR, Smith PN (2012) Embryotoxicity of mixtures of weathered crude oil collected from the Gulf of Mexico and Corexit 9500 in mallard ducks (*Anas platyrhynchos*). Sci Total Environ. 426: 155–159.

Finch BE, Wooten KJ, Smith PN (2011) Embryotoxicity of weathered crude oil from the Gulf of Mexico in mallard ducks (*Anas platyrhynchos*). Environ Toxicol Chem 30: 1885–1891.

Fine Tailings Fundamentals Consortium (1995) Advances in Oil Sands tailings research. Edmonton, Alberta: Alberta Department of Energy, Oil Sands and Research Division.

Fischer RA, Gauthreaux SA, Valente JJ, Guilfoyle MP, Kaller MD (2012) Comparing transect survey and WSR-88D radar methods for monitoring daily changes in stopover migrant communities. J Field Ornithol 83: 61–72.

Fisher JT, Wilkinson L (2005) The response of mammals to forest fire and timber harvest in the North American boreal forest. Mamm Rev35: 51–81.

Fleishman E, Murphy DD (2009) A realistic assessment of the indicator potential of butterflies and other charismatic taxonomic groups. Conserv Biol 23: 1109–1116.

Fleming WJ, Sileo L, Franson JC (1982) Toxicity of Prudhoe Bay crude oil to sandhill cranes. The J Widlife Manag 46: 474–478.

Flickinger EL (1981) Wildlife mortality at petroleum pits in Texas. The J Widlife Manag 45: 560–564.

Flickinger EL, Bunck CM (1987) Number of oil-killed birds and fate of bird carcasses at crude oil pits in Texas. Southwest Nat 32: 377–381.

Flint PL, Schamber JL, Trust KA, Miles AK, Henderson JD, et al. (2012) Chronic hydrocarbon exposure of harlequin ducks in areas affected by the Selendang Ayu oil spill at Unalaska Island, Alaska. Environ Toxicol Chem. 31: 2828–2831.

Floater GJ (2002) A risk model for predicting the effects of lake-side development on wildfowl populations. J Environ Manag 66: 307–316.

Flock WL, Green JL (1974) The detection and identification of birds in flight, using coherent and noncoherent radars. Proceedings of the IEEE 62: 745–753.

U.S. Fish and Wildlife Service (2014) General Flyways Info. Washington D.C: United States Fish and Wildlife Service. Available: http://www.flyways.us/flyways/info. Accessed 5 February 2014.

Focardi S, De Marinis AM, Rizzotto M, Pucci A (2001) Comparative evaluation of thermal infrared imaging and spotlighting to survey wildlife. Wildlife Soc B 29: 133–139.

Foote L (2012) Threshold Considerations and Wetland Reclamation in Alberta's Mineable Oil Sands. Ecol Soc 17. DOI: http://dx.doi.org/10.5751/ES-04673-170135

Ford RG, Weins JA, Heinemann D, Hunt GL (1982) Modelling the sensitivity of colonially breeding marine birds to oil spills: guillemot and kittiwake populations on the Pribilof Islands, Bering Sea. J Appl Ecol: 1–31.

Forsman JT, Mönkkönen M, Korpimäki E, Thomson RL (2013) Mammalian nest predator feces as a cue in avian habitat selection decisions. Behav Ecol 24: 262–266.

Fowler A, Flint PL (1997) Persistence rates and detection probabilities of oiled king eider carcasses on St Paul Island, Alaska. Mar Pollut Bull 34: 522–526.

Fowler GS, Wingfield JC, Boersma PD (1995) Hormonal and reproductive effects of low levels of petroleum fouling in Magellanic penguins (*Spheniscus magellanicus*). Auk: 382–389.

Fox A, Desholm M, Kahlert J, Christensen TK, Krag Petersen I (2006) Information needs to support environmental impact assessment of the effects of European marine offshore wind farms on birds. Ibis 148: 129–144.

Fox A, Madsen J (1997) Behavioural and distributional effects of hunting disturbance on waterbirds in Europe: implications for refuge design. J Appl Ecol: 1–13.

Fox G (2001) Effects of endocrine disrupting chemicals on wildlife in Canada: past, present and future. Water Qual Res J Canada 36: 233–251.

Franci CD, Guillemette M, Pelletier E, Chastel O, Bonnefoi S, et al. (2013) Endocrine status of a migratory bird potentially exposed to the Deepwater Horizon oil spill: A case study of northern gannets breeding on Bonaventure Island, Eastern Canada. Sci Total Environ 473-474C: 110–116.

Francis CD, Paritsis J, Ortega CP, Cruz A (2011) Landscape patterns of avian habitat use and nest success are affected by chronic gas well compressor noise. Lands Ecol 26: 1269–1280.

Francis CM, Blancher PJ, Phoenix RD (2009) Bird monitoring programs in Ontario: What have we got and what do we need? Forest Chron85: 202–217.

Frank RA, Fischer K, Kavanagh R, Burnison BK, Arsenault G, et al. (2008) Effect of carboxylic acid content on the acute toxicity of oil sands naphthenic acids. Environ Sci Technol 43: 266–271.

Frank RA, Roy JW, Bickerton G, Rowland S, Headley JV, et al. (2014) Profiling oil sands mixtures from industrial developments and natural groundwaters for source identification. Environ Sci Technol.DOI: dx.doi.org/10.1021/es500131k

Fraser GS, Ellis J (2009) The Canada-Newfoundland Atlantic Accord Implementation Act: Transparency of the Environ Manag of the offshore oil and gas industry. Mar Policy 33: 312–316.

Fraser GS, Russell J, Von Zharen W (2006) Produced water from offshore oil and gas installations on the Grand Banks, Newfoundland and Labrador: are the potential effects to seabirds sufficiently known. Mar Ornithol 34: 147–156.

Frederiksen M, Moe B, Daunt F, Phillips RA, Barrett RT, et al. (2012) Multicolony tracking reveals the winter distribution of a pelagic seabird on an ocean basin scale. Divers Distrib 18: 530–542.

Fritcher DL, Mazet JA, Ziccardi MH, Gardner IA (2002) Evaluation of two direct immunoassays for rapid detection of petroleum products on marine birds. Mar Pollut Bull 44: 388–395.

Fry DM, Lowenstine LJ (1985) Pathology of common murres and Cassin's auklets exposed to oil. Arch Environ Contam Toxicol 14: 725–737.

Fry DM, Swenson J, Addiego L, Grau C, Kang A (1986) Reduced reproduction of wedge-tailed shearwaters exposed to weathered Santa Barbara crude oil. Arch Environ Contam Toxicol 15: 453–463.

Fuller-Perrine LD, Tobin ME (1993) A method for applying and removing bird-exclusion netting in commercial vineyards. Wildlife Soc B 21: 47–51.

Furness RW, Wade HM, Masden EA (2013) Assessing vulnerability of marine bird populations to offshore wind farms. J Environ Manag 119: 56–66.

Gagné F, Douville M, André C, Debenest T, Talbot A, et al. (2012) Differential changes in gene expression in rainbow trout hepatocytes exposed to extracts of oil sands process-affected water and the Athabasca River. Comp Biochem and Phys C Toxicol Pharmacol 155: 551–559.

Gagnon F, Ibarzabal J, Savard JPL, Vaillancourt P, Bélisle M, et al. (2011) Weather effects on autumn nocturnal migration of passerines on opposite shores of the St. Lawrence estuary. Auk 128: 99–112.

Gallup Jr GG, Cummings WH, Nash RF (1972) The experimenter as an independent variable in studies of animal hypnosis in chickens (*Gallus gallus*). Anim Behav20: 166–169.

Gandini P, Boersma PD, Frere E, Gandini M, Holik T, et al. (1994) Magellanic penguins (*Spheniscus magellanicus*) affected by chronic petroleum pollution along coast of Chubut, Argentina. Auk: 20–27.

Garcia-Borboroglu P, Boersma PD, Ruoppolo V, Reyes L, Rebstock GA, et al. (2006) Chronic oil pollution harms Magellanic penguins in the Southwest Atlantic. Mar Pollut Bull 52: 193–198.

Gardner RC, Davidson NC (2011) The Ramsar convention. Wetlands. Springer. pp. 189–203.

Garvey ME, Nol E, Howerter DW, Armstrong LM (2013) A spatial analysis of factors affecting nesting success of shorebirds in the canadian prairies. Condor 115: 58–66.

Gasc A, Sueur J, Pavoine S, Pellens R, Grandcolas P (2013) Biodiversity sampling using a global acoustic approach: contrasting sites with microendemics in new caledonia. PLoS ONE 8: e65311.

Gauthreaux (2003) Radar Ornithology and Biol Conservation. Auk 120: 266–277.

Gauthreaux Jr SA, Belser CG (1998) Displays of bird movements on the wsr-88d: patterns and quantification. Weather and Forecasting 13: 453–464.

Gauthreaux SA, Livingston JW (2006) Monitoring bird migration with a fixed-beam radar and a thermal-imaging camera. J Field Ornithol 77: 319–328.

Gay M, Belisle A, Patton J (1980) Quantification of petroleum-type hydrocarbons in avian tissue. J Chromatogr A187: 153–160.

Gendron M, Smyth SA, Stewart GR, Pollard JB (2001) Pease Athabasca delta waterbird inventory 2000 surveys: final report. Edmonton, Alberta: Ducks Unlimited Canada.

Gentes M-L (2006) Health assessment of tree swallows (*Tachycineta bicolor*) nesting on the Athabasca Oil Sands, Alberta. Master of Science, University of Saskatchewan, Saskatoon, SK, Canada.

Gentes M-L, McNabb A, Waldner C, Smits JEG (2007) Increased thyroid hormone levels in tree swallows (*Tachycineta bicolor*) on reclaimed wetlands of the Athabasca oil sands. Arch Environ Contam Toxicol 53: 287–292.

Gentes M-L, Waldner C, Papp Z, Smits JEG (2006) Effects of oil sands tailings compounds and harsh weather on mortality rates, growth and detoxification efforts in nestling tree swallows (*Tachycineta bicolor*). Environ Pollut 142: 24–33.

Gentes M-L, Waldner C, Papp Z, Smits JEG (2007) Effects of exposure to naphthenic acids in tree swallows (*Tachycineta bicolor*) on the Athabasca oil sands, Alberta, Canada. J Toxicol Environ Health A 70: 1182–1190.

Gentes M-L, Whitworth TL, Waldner C, Fenton H, Smits JE (2007) Tree swallows (*Tachycineta bicolor*) nesting on wetlands impacted by oil sands mining are highly parasitized by the bird blow fly Protocalliphora spp. J Wildl Dis 43: 167–178.

Gibson JJ, Birks SJ, Moncur M, Yi Y, Tattrie K, et al. (2011) Isotopic and geochemical tracers for fingerprinting process-affected waters in the oil sands industry: A Pilot Study. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-12. 109 pp.

Gibson JJ, Fennell J, Birks J, Yi Y, Moncur MC, et al. (2013) Evidence of discharging saline formation water to the Athabasca River in the oil sands mining region, northern Alberta. Can J Earth Sci. 50: 1244–1257.

Giese M, Goldsworthy SD, Gales R, Brothers N, Hamill J (2000) Effects of the Iron Baron oil spill on little penguins (*Eudyptula minor*). III. Breeding success of rehabilitated oiled birds. Wildlife Res 27: 583–591.

Giesy JP, Anderson JC, Wiseman SB (2010) Alberta oil sands development. PNAS 107: 951–952.

Gill RB (2007) Ornithology. 3rd ed. New York: W.H. Freeman and Co.

Gilmer DS, Brass JA, Strong LL, Card DH (1988) Goose counts from aerial photographs using an optical digitizer. Wildlife Soc B 16: 204–206.

Gilsdorf JM, Hygnstrom SE, VerCauteren KC (2002) Use of frightening devices in wildlife damage management. Integr Pest Manag Rev 7: 29–45.

Giraudeau M, Duval C, Guillon N, Bretagnolle V, Gutierrez C, et al. (2010) Effects of access to preen gland secretions on mallard plumage. Naturwissenschaften 97: 577–581.

Glahn JF, Ellis G, Fioranelli P, Dorr BS (2000) Evaluation of moderate and low-powered lasers for dispersing double-crested cormorants from their night roosts. Wildlife Damage Management Conferences-Proceedings. p. 11.

Glen AS, Cockburn S, Nichols M, Ekanayake J, Warburton B (2013) Optimising camera traps for monitoring small mammals. PLoS ONE 8: e67940.

Gochfeld M (1979) Prevalence of oiled plumage of terns and skimmers on Western Long Island, New York: baseline data prior to petroleum exploration. Environ Pollut 20: 123–130.

Godfrey WE (1966) The Birds of Canada. Ottawa: National Museum of Natural Sciences, National Museums of Canada. 595p.

Godwalt C, Kotecha P, Aumann C (2010) Oil sands tailings management project. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-7. 64 pp.

Goethe F (1968) The effects of oil pollution on populations of marine and coastal birds. Helgoländer wissenschaftliche Meeresuntersuchungen 17: 370–374.

Golder Associates (1996) Toxicity and bioaccumulation potential of fine tails and tailings water from oil sands extraction. Golder Associates Ltd, submitted to the Alberta Oil Sands Technology and Research Authority. Golder Project No. 942-2287. Calgary, AB, Canada.

Golder Associates (1997) Field scale trials to assess effects of consolidated tails release water on plants and wetland ecology. Golder Associates Ltd. for Suncor Energy Inc., Oil Sands. Calgary, AB, Canada.

Golder Associates (1999) The effects of consolidated tailings release water and deposits on terrestrial and wetlands ecosystems summary report #2. Prepared for Suncor Energy Ltd., Oil Sands, by Golder Associates. Calgary, AB, Canada.

Golder Associates (2009) The Suncor bird deterrent and monitoring program: History and evolution. Prepared for Suncor Inc. Oil Sands Group. Calgary, AB, Canada.

Golder Associates (2012) Literature Search Regarding the Effects of Spilled Oil on Birds. Prepared for Canadian National Railway. Golder Project Report Project No. 11-1421-0040. Burnaby, British Columbia, Canada.

Golder Associates Ltd (2000) Report on oil sands tailings pond bird deterrent systems-a review of research and current practices. Suncor Energy Inc., Syncrude Canada Ltd., and Albian Sands Energy Inc. Calgary. Alberta, Canada.

Golder Associates Ltd. (1995) 1993/1994 Reclamation landscape model development project. Submitted to Suncor Inc., Oil Sands Group and Syncrude Canada Limited. Golder Project Report No. 952-2339. Calgary, AB, Canada.

Golder Associates Ltd. (1996) Aquatic baseline report for the Athabasca, Steepbank and Muskeg rivers in the vicinity of the Steepbank and Aurora mines. Published by Suncor Energy Inc. . Calgary, AB, Canada.

Golder Associates Ltd. (1996) Athabasca river water releases impact assessment. Prepared for Suncor Inc. Oil Sands Group by Golder Associates Ltd. Golder Project Report No. 952-2307. 123 p. . Calgary, AB, Canada.

Golder Associates Ltd. (1996) Fish flavour impairment study. Prepared for Suncor Inc Oil Sands Group by Golder Associates Ltd. Golder Report No. 952-2307. Calgary, AB, Canada.

Golder Associates Ltd. (1996) Impact analysis of aquatic issues associated with Steepbank mine. Prepared for Suncor Inc. Oil Sands Group by Golder Associates Ltd. Golder Project Report No. 952-2307. Calgary, AB, Canada.

Golder Associates Ltd. (1996) Impact analysis of human health issues associated with the Steepbank mine. Prepared for Suncor Inc. Oil Sands Group by Golder Associates Ltd. Golder Project Report No. 952-2307. Calgary, AB, Canada.

Golder Associates Ltd. (1996) Impact analysis of terrestrial resources associated with the Steepbank mine. Prepared for Suncor Inc. Oil Sands Group by Golder Associates Ltd. Calgary Alberta, Canada.

Golder Associates Ltd. (1996) Laboratory studies on trophic effects and fish health. Prepared for Suncor Inc. Oil Sands Group by Golder Associates Ltd. Golder Project Report No. 952-2307. Calgary Alberta, Canada.

Golder Associates Ltd. (1996) Suncor mine advance plan (D&R) and cumulative effects assessment. Prepared for Suncor Inc. Oil Sands Group by Golder Associates Ltd. Calgary, AB, Canada

Golder Associates Ltd. (1996) Terrestrial baseline report for the steepbank mine. Prepared for Suncor Inc. Oil Sands Group by Golder Associates Ltd. Calgary, AB, Canada.

Golder Associates Ltd. (1996) Visual impact of Suncor Steepbank mine development. Prepared for Suncor Inc. Oil Sands Group by Golder Associates Ltd. Calgary, AB, Canada.

Golder Associates Ltd. (1997) Field scale trials to assess effects of consolidated tailings release water on plants and wetlands ecology: Technical Report #2. Golder Associates Ltd. for Suncor Energy Inc., Oil Sands.

Golder Associates Ltd. (1998) 1997 synthesis of environmental information on consolidated/composite tailings (CT). Report No. 972-2205.6045. Golder Associates Ltd.: Calgary, Alberta, Canada.

Golder Associates Ltd. (1998) Report on sands regional aquatic monitoring program (RAMP) 1997. Golder Associates Ltd. submitted to RAMP Steering Committee

Golder Associates Ltd. (1999) Report on sands regional aquatic monitoring program (RAMP) 1998. Golder Associates Ltd. submitted to RAMP Steering Committee.

Golder Associates Ltd. (2003) Oil sands regional aquatic monitoring program: Five Year Report. Golder Associates Ltd. submitted to RAMP Steering Committee

Golder Associates Ltd. (2003) Oil sands regional aquatics monitoring program (RAMP) 2002. Golder Associates Ltd. submitted to RAMP Steering Committee

Golder Associates, Can-AG_Enterprises_Ltd. (1996) Baseline soil survey for the proposed Suncor steepbank mine. Prepared for Suncor Inc. Oil Sands Group. Calgary, AB, Canada.

Golder Associates, EnviResource_Consulting_Ltd. (1996) Suncor Inc. Mine Expansion: Baseline Forestry Report. Prepared for Suncor Inc. Oil Sands Group. Calgary, AB, Canada.

Golder Associates, Westworth B (1996) Abundance & Distribution of Moose in the Suncor Study Area. Prepared for Suncor Inc. Oil Sands Group. Calgary, AB, Canada.

Golder Associates, Westworth B (1996) Baseline Study for Fur Trapping in the Suncor Study Area. Baseline Study for Fur Trapping in the Suncor Study Prepared for Suncor Inc. Oil Sands Group. Calgary, AB, Canada.

Golder Associates, Westworth B (1996) Habitat Suitability Models for the Suncor Study Area. Prepared for Suncor Inc. Oil Sands Group. Calgary, AB, Canada.

Golder Associates, Westworth B (1996) Herpetofauna in the Steepbank study area. Suncor Inc., Oil Sands Group. Calgary, AB, Canada.

Golder Associates, Westworth B (1996) Impact analysis Suncor Steepbank Mine environemtnal wildlife component. Prepared for Suncor Inc. Oil Sands Group. Calgary, AB, Canada.

Golder Associates, Westworth B (1996) Waterfowl, Raptors and Breeding Terrestrial Birds of the Suncor Lease in 1995. Prepared for Suncor Inc. Oil Sands Group. Calgary, AB, Canada.

Goldsworthy S, Gales R, Giese M, Brothers N (2000) Effects of the Iron Baron oil spill on little penguins (*Eudyptula minor*). I. Estimates of mortality. Wildlife Res 27: 559–571.

Goldsworthy SD, Giese M, Gales R, Brothers N, Hamill J (2000) Effects of the Iron Baron oil spill on little penguins (*Eudyptula minor*). II. Post-release survival of rehabilitated oiled birds. Wildlife Res 27: 573–582.

Golet GH, Seiser PE, McGuire AD, Roby DD, Fischer JB, et al. (2002) Long-term direct and indirect effects of the "Exxon Valdez" oil spill on pigeon guillemots in Prince William Sound, Alaska. Mar Ecol Prog Ser 241: 287–304.

Golightly RT, Newman SH, Craig EN, Carter HR, Mazet JAK (2002) Survival and behavior of Western Gulls following exposure to oil and rehabilitation. Wildlife Soc B: 539–546.

Gómez C, Bayly NJ, Rosenberg KV (2013) Seasonal variation in stopover site use: Catharus thrushes and vireos in northern Colombia. J Ornithol 154: 107–117.

González-Solis J, Shaffer SA (2009) Introduction and synthesis: spatial ecology of seabirds at sea. Mar Ecol Prog Ser 391: 117–120.

Goodenough AE, Smith AL, Stubbs H, Williams R, Hart AG (2012) Observer variability in measuring animal biometrics and fluctuating asymmetry when using digital analysis of photographs. Annales Zoologici Fennici. 49: 81–92.

Gorenzel P, Salmon T (2008) Bird Hazing Manual: Techniques and Strategies for Dispersing Birds from Spill Sites. Davis, California: Department of Wildlife, Fish, and Conservation Biology University of California.

Gorenzel W, Blackwell B, Simmons G, Salmon T, Dolbeer R (2002) Evaluation of lasers to disperse American crows, *Corvus brachyrhynchos*, from urban night roosts. Int J Pest Manag 48: 327–331.

Gorman ML, Milne H (1971) Seasonal changes in the adrenal steroid tissue of the Common eider Somateria mollissima and its relation to organic metabolism in normal and oil-polluted birds. Ibis 113: 218–228.

Gorman ML, Simms CE (1978) Lack of effect of ingested forties field crude oil on avian growth. Mar Pollut Bull 9: 273–276.

Gorsline J, Holmes WN (1981) Effects of petroleum on adrenocortical activity and on hepatic naphthalene-metabolizing activity in mallard ducks. Arch Environ Contam Toxicol 10: 765–777.

Gorsline J, Holmes WN (1982) Adrenocortical function and hepatic naphthalene metabolism in Mallard ducks (*Anas platyrhynchos*) consuming petroleum distillates. Environ Res 28: 139–146.

Gorsline J, Holmes WN (1982) Ingestion of petroleum by breeding mallard ducks: some effects on neonatal progeny. Arch Environ Contam Toxicol 11: 147–153.

Gorsline J, Holmes WN (1982) Suppression of adrenocortical activity in Mallard ducks exposed to petroleum-contaminated food. Arch Environ Contam Toxicol 11: 497–502.

Gorsline J, Holmes WN (1982) Variations with age in the adrenocortical responses of mallard ducks (*Anas platyrhynchos*) consuming petroleum-contaminated food. Bull Environ Contam Toxicol 29: 146–152.

Gorsline J, Holmes WN, Cronshaw J (1981) The effects of ingested petroleum on the naphthalene-metabolizing properties of liver tissue in seawater-adapted mallard ducks (*Anas platyrhynchos*). Environ Res 24: 377–390.

Gosler A, Kenward R, Horton N (1995) The effect of gull roost deterrence on roost occupancy, daily gull movements and wintering wildfowl. Bird Study 42: 144–157.

Gosselin P, Hrudey SE, Naeth M, Plourde A, Therrien R, Van Der Kraak G, Xu Z (2010) Environmental and health impacts of Canada's oil sands industry. Royal Society of Canada Expert Panel: Ottawa, ON, Canada.

Gosser AL, Conover MR, Messmer TA (1997) Managing problems caused by urban Canada geese.Berryman Institute Publication 13, Utah State University, Logan. 8 p.

Gotelli NJ (2000) Null model analysis of species co-occurrence patterns. Ecology 81: 2606–2621.

Goudie R, Jones IL (2004) Dose-response relationships of harlequin duck behaviour to noise from low-level military jet over-flights in central Labrador. Environ Conserv 31: 289–298.

Goudie RI (2006) Multivariate behavioural response of harlequin ducks to aircraft disturbance in Labrador. Environ Conserv 33: 28–35.

Government of Canada (1994) The Migratory Birds Convention Act. Chapter 22. Available: http://laws-lois.justice.gc.ca/PDF/M-7.01.pdf. Accessed 2013 October 1.

Grant J, Huot M, Lemphers N, Dyer S, Dow M (2013) Beneath the surface: a review of key facts in the oilsands debate. Drayton Valley, Alberta: Pembina Institute.

Grasman KA (2002) Assessing immunological function in toxicological studies of avian wildlife. Integr Comp Biol 42: 34–42.

Grau CR, Roudybush T, Dobbs J, Wathen J (1977) Altered yolk structure and reduced hatchability of eggs from birds fed single doses of petroleum oils. Science 195: 779–781.

Green TJ (1997) Target recognition theory for laser radar imaging. In: AeroSense 97 Conference on Photonic Quantum Computing, Orlando.

Green RH (1989) Power analysis and practical strategies for environmental monitoring. Environ Res 50: 195–205.

Greenley WB, Barta WD, Eddinger RT (1982) Literature review of toxicological properties of petroleum and coal-derived oil products. Fuel Process Technol 6: 9–25.

Greenwood J, Donally R, Feare C, Gordon NJ, Waterston G (1971) A massive wreck of oiled birds: northeast Britain, winter 1970. Scottish birds 6: 235–250.

Griffin AS, Blumstein DT, Evans CS (2000) Training captive-bred or translocated animals to avoid predators. Conserv Biol 14: 1317–1326.

Griffin AS, Evans CS, Blumstein DT (2001) Learning specificity in acquired predator recognition. Anim Behav 62: 577–589.

Groom G, Krag Petersen I, Anderson MD, Fox AD (2011) Using object-based analysis of image data to count birds: mapping of Lesser Flamingos at Kamfers Dam, Northern Cape, South Africa. Int J Remote Sens 32: 4611–4639.

Groom G, Stjernholm M, Nielsen RD, Fleetwood A, Petersen IK (2013) Remote sensing image data and automated analysis to describe marine bird distributions and abundances. Ecological Inform 14: 2–8.

Groom MJ, Meffe GK, Carroll CR, others (2006) Principles of conservation biology. Sinauer Associates Sunderland.

Gu W, Swihart RK (2004) Absent or undetected? Effects of non-detection of species occurrence on wildlife-habitat models. Biol Cons 116: 195–203.

Guarino JL (1972) Methiocarb, a chemical bird repellent: a review of its effectiveness on crops. Proceedings of the 5th Vertebrate Pest Conference (1972). Paper 20. Available: http://digitalcommons.unl.edu/vpc5/20

Guillaume G, Matthieu G, du Rau Pierre D, Patrick G (2014) Effects of mute swans on wetlands: a synthesis. Hydrobiologia 723: 195–204.

Gulley JR (1979) Waterfowl investigation reports 1977-1980, Avifauna report – 1979, Rehabilitation report - 1980. Prepared for Suncor Inc.

Gulley JR (1980) Factors influencing the efficacy of human effigies in deterring waterfowl from polluted ponds. Master of Science, University of Alberta, Edmonton, Canada.

Gulley JR (1981) Avifauna studies on crown lease no. 86 for Suncor Inc. 1976 - 1980. Prepared for Suncor Inc.. Boreas Environmental Consulting Services Ltd.

Gulley JR (1981) Waterfowl investigations for Suncor Inc. Fort McMurray - 1981. Prepared for Suncor Inc.. Boreas Environmental Consulting Services Ltd.

Gulley JR (1982) Avifauna studies on crown lease no. 86 1981-1982 update for Suncor Inc. Prepared for Suncor Inc.. Boreas Environmental Consulting Services Ltd.

Gulley JR (1982) Daily avifauna summaries for lease no. 86, Suncor Inc. 1976 - 1982. Prepared for Suncor Inc.. Boreas Environmental Consulting Services Ltd.

Gulley JR (1982) Waterfowl investigations for Suncor Inc. Fort McMurray, Alberta - 1982. Prepared for Suncor Inc.. Boreas Environmental Consulting Services Ltd.

Gulley JR (1983) Avifauna studies on crown lease no. 86 for Suncor Inc. 1976 - 1983. Prepared for Suncor Inc.. Boreas Environmental Consulting Services Ltd.

Gulley JR (1983) Investigations of avian activities on crown lease no. 86 for Suncor Inc., Fort McMurray, Alberta. 1983. Prepared for Suncor Inc.. Boreas Environmental Consulting Services Ltd.

Gulley JR (1985) Investigations of avian activities on crown lease no. 86 for Suncor Inc., Fort McMurray, Alberta. 1984. Prepared for Suncor Inc.. Boreas Environmental Consulting Services Ltd.

Gulley JR (1986) Investigations of avian activities on crown lease no. 86 for Suncor Inc., Fort McMurray, Alberta. 1985. Prepared for Suncor Inc.. Boreas Environmental Consulting Services Ltd.

Gulley JR (1987) Examination of waterfowl trends on crown lease no. 86 for Suncor Inc., Fort MeMurray, Alberta. 1976 - 1986. Prepared for Suncor Inc.. Boreas Environmental Consulting Services Ltd.

Gulley JR (1987) Investigations of avian activities on crown lease no. 86 for Suncor Inc. Fort McMurray, Alberta, 1986. Prepared for Suncor Inc.. Boreas Environmental Consulting Services Ltd.

Gulley JR (1988) The Suncor avifauna program: summary of the 1988 activites and results. Prepared for Suncor Inc.. Boreas Environmental Consulting Services Ltd.

Gulley JR (1990) The Suncor avifauna program: summary of the 1989 activites and results. Prepared for Suncor Inc.. Boreas Environmental Consulting Services Ltd.

Gulley JR (1990) The Suncor avifauna program: summary of the 1990 activites and results. Prepared for Suncor Inc.. Boreas Environmental Consulting Services Ltd.

Gulley JR (1992) Suncor Avifauna Program Ponds 1, 1A, 2, 3, and 4 January 30 - September 29, 1992. Prepared for Suncor Inc.. Boreas Environmental Consulting Services Ltd.

Gulley JR (1992) The Suncor avifauna program: summary of the 1991 activites and results. Prepared for Suncor Inc.. Boreas Environmental Consulting Services Ltd.

Gulley JR (1993) 1992-1993 Fine Tails Reclamation Research and Development Program. Fort McMurray, Alberta: Suncor Inc..

Gulley JR, Hamilton HR, Taylor B (1995) Oil Sands tailings leachability and toxicity evaluation. Fort McMurray, Alberta: Suncor Inc., Oil Sands Group.

Gulley JR, Klynn DJ (1992) Wetland treatment of oil sands operation waste waters. In: Environmental Issues and Waste Management in Energy and Minerals Production, Rotterdam.

Gulley JR, MacKinnon M (1993) Fine tails reclamation utilizing a wet landscape approach. In: Oil Sands – Our Petroleum Future Conference, Edmonton.

Gunn WWH (1951) Analysis of unusual bird migration in North America during the storm of April 4-7, 1947. Auk 68: 139–163.

Gupta N (2009) Effects of oil sands process-affected water and substrates on wood frog (*Rana sylvatica*) eggs and tadpoles. Master of Science, University of Saskatchewan, SK, Canada.

Gurney KE, Williams TD, Smits JE, Wayland M, Trudeau S, et al. (2005) Impact of oil-sands based wetlands on the growth of mallard (*Anas platyrhynchos*) ducklings. Environ Toxicol Chem 24: 457–463.

Gutiérrez-Ibáñez C, Iwaniuk AN, Wylie DR (2009) The independent evolution of the enlargement of the principal sensory nucleus of the trigeminal nerve in three different groups of birds. Brain Behav Evol 74: 280–294.

Haché S, Villard M-A, Bayne EM (2013) Experimental evidence for an ideal free distribution in a breeding population of a territorial songbird. Ecology 94: 861–869.

Hagen MO (2013) Analysis of goldfish innate immunity following exposure to oil sands process affected water. Master of Science, University of Alberta, Edmonton, Alberta, Canada.

Hagen MO, Katzenback B, Islam M, El-Din MG, Belosevic M (2013) The analysis of goldfish (*Carassius auratus L.*) innate immune responses after acute and sub chronic exposures to oil sands process-affected water. Toxicol Sci. 138: 59-68.

Hall RJ, Coon NC (1988) Interpreting residues of petroleum hydrocarbons in wildlife tissues. Laurel, Maryland: Patuxent Wildlife Research Center, U.S. Fish and Wildlife Service.

Hallett D, Onuska F, Comba M (1983) Aliphatic and polyaromatic components of weathered and unweathered southern Louisiana crude oil. Mar Environ Res 8: 73–85.

Halvorsen CF, Center USCGRT, Yorktown V (1979) Types and Characteristics of Oils and Hazardous Chemicals. Proceedings of the 1979 US Fish and Wildlife Service Pollution Response Workshop, 8-10 May 1979, St. Petersburg, Florida. p. 55.

Hamel S, Killengreen ST, Henden J-A, Eide NE, Roed-Eriksen L, et al. (2012) Towards good practice guidance in using camera-traps in ecology: influence of sampling design on validity of ecological inferences. Methods Ecol Evol 4: 105–113.

Hamer TE, Cooper BA, Ralph CJ (1995) Use of radar to study the movements of Marbled Murrelets at inland sites. Northwest Nat 76: 73–78.

Hampton RR (1994) Sensitivity to information specifying the line of gaze of humans in sparrows (*Passer domesticus*). Behaviour 130: 41–51.

Han X, MacKinnon MD, Martin JW (2009) Estimating the in situ biodegradation of naphthenic acids in oil sands process waters by HPLC/HRMS. Chemosphere 76: 63–70.

Hanley JA (1982) Characteristic (ROC) Curve. Radiol 743: 29–36.

Harding EG, Curtis PD, Vehrencamp SL (2007) Assessment of management techniques to reduce woodpecker damage to homes. The J Widlife Manag 71: 2061–2066.

Harfenist A, Gilman AP, Maus KL (1990) The effects of exposure of incubating adult and young Herring Gulls to a simulated No. 2 fuel oil slick. Arch Environ Contam Toxicol 19: 902–906.

Harmata AR, Podruzny KM, Zelenak JR, Morrison ML (1999) Using marine surveillance radar to study bird movements and impact assessment. Wildlife Soc B 27: 44–52.

Harmata AR, Podruzny KM, Zelenak JR, Morrison ML (2000) Passage rates and timing of bird migration in Montana. Am Midl Nat 143: 30–40.

Harms NJ, Fairhurst GD, Bortolotti GR, Smits JEG (2010) Variation in immune function, body condition, and feather corticosterone in nestling tree swallows (*Tachycineta bicolor*) on reclaimed wetlands in the Athabasca oil sands, Alberta, Canada. Environ Pollut 158: 841–848.

Haroldson BS, Wiggers EP, Beringer J, Hansen LP, McAninch JB (2003) Evaluation of aerial thermal imaging for detecting white-tailed deer in a deciduous forest environment. Wildlife Soc B 31: 1188–1197.

Harriman L, Muhlhausen J (2013) A new eye in the sky: Eco-drones. Article reproduced from United Nations Environment Programme (UNEP) Global Environmental Alert Service (GEAS). Environ Dev 7: 155-164.

Harris G, Thompson R, Childs JL, Sanderson JG (2010) Automatic storage and analysis of camera trap data. Bull Ecol Soc Am 91: 352–360.

Harris RE, Davis RA (1998) Evaluation of the efficacy of products and techniques for airport bird control. King City, Ontario: LGL Limited Environmental Research Associates.

Hart N, Partridge J, Cuthill I (1998) Visual pigments, oil droplets and cone photoreceptor distribution in the European starling (*Sturnus vulgaris*). The J Exp Biol 201: 1433–1446.

Hart NS (2001) Variations in cone photoreceptor abundance and the visual ecology of birds. J Comp Physiol A 187: 685–697.

Hart NS, Hunt DM (2007) Avian visual pigments: characteristics, spectral tuning, and evolution. Am Nat 169: S7–S26.

Hart V, Malkemper EP, Kušta T, Begall S, Nováková P, et al. (2013) Directional compass preference for landing in water birds. Front Zool 10: 38.

Hartung R (1965) Some effects of oiling on reproduction of ducks. The J Widlife Manag: 872–874.

Hartung R (1967) Energy metabolism in oil-covered ducks. The J Widlife Manag 31: 798–804.

Hartung R, Hunt GS (1966) Toxicity of some oils to waterfowl. The J Widlife Manag: 564–570.

Harvey S, Klandorf H, Phillips JG (1981) Reproductive performance and endocrine responses to ingested petroleum in domestic ducks (*Anas platyrhynchos*). Gen Comp Endocrinol 45: 372–380.

Harvey S, Sharp PJ, Phillips JG (1982) Influence of ingested petroleum on the reproductive performance and pituitary-gonadal axis of domestic ducks (*Anas platyrhynchos*). Comp Biochem Physiol C Pharmacol Comp Pharmacol 72: 83–89.

Harvey S, Sharp PJ, Phillips JG (1982) Influence of sublethal doses of ingested petroleum on reproductive performance and the pituitary adrenal axis in domestic ducks (*Anas platyrhynchos*). Comp Biochem Physiol C 72: 83–89.

Harwell MA, Gentile JH, Parker KR, Murphy SM, Day RH, et al. (2012) Quantitative Assessment of Current Risks to Harlequin Ducks in Prince William Sound, Alaska, from the Exxon Valdez Oil Spill. Hum Ecol Risk Assess 18: 261–328.

Hatfield Consultants Ltd. (2006) Aquatic resources environmental impact assessment: surface water quality, sediment quality, and benthic invertebrate communities. Prepared for Millennium EMS Solutions Ltd.. West Vancouver, British Columbia: Hatfield Consultants Ltd..

Haughland DL, Hero J-M, Schieck J, Castley JG, Boutin S, et al. (2010) Planning forwards: biodiversity research and monitoring systems for better management. Trends Ecol Evol 25: 199–200.

Haukos DA, Smith LM (1994) The importance of playa wetlands to biodiversity of the Southern High Plains. Landsc Urban Plan 28: 83–98.

Hausmann F, Arnold KE, Marshall NJ, Owens IPF (2003) Ultraviolet signals in birds are special. Roc R Soc Lond B Biol Sci 270: 61–67.

Havens KJ, Sharp EJ (1998) Using thermal imagery in the aerial survey of animals. Wildlife Soc B 26: 17–23.

He Y, Wiseman SB, Hecker M, Zhang X, Wang N, et al. (2011) Effect of ozonation on the estrogenicity and androgenicity of oil sands process-affected water. Environ Sci Technol 45: 6268–6274.

He Y, Wiseman SB, Wang N, Perez-Estrada LA, El-Din MG, et al. (2012) Transcriptional responses of the brain-gonad-liver axis of fathead minnows exposed to untreated and ozone-treated oil sands process-affected water. Environ Sci Technol 46: 9701–9708.

Headley JV, McMartin DW (2004) A review of the occurrence and fate of naphthenic acids in aquatic environments. J Environ Sci Health A 39: 1989–2010.

Heatley JJ (2002) Antipredator conditioning in Mississippi Sandhill cranes (*Grus canadensis pulla*). Master of Science, Louisiana State University and Agricultural and Mechanical College.

Hebert CE, Campbell D, Kindopp R, MacMillan S, Martin P, et al. (2013) Mercury trends in colonial waterbird eggs downstream of the oil sands region of Alberta, Canada. Environ Sci Technol 47: 11785–11792.

Hebert CE, Weseloh DVC, Macmillan S, Campbell D, Nordstrom W (2011) Metals and polycyclic aromatic hydrocarbons in colonial waterbird eggs from Lake Athabasca and the Peace-Athabasca Delta, Canada. Environ Toxicol Chem 30: 1178–1183.

Hedd A, Montevecchi W, McFarlane Tranquilla L, Burke C, Fifield D, et al. (2011) Reducing uncertainty on the Grand Bank: tracking and vessel surveys indicate mortality risks for common murres in the North-West Atlantic. Anim Conserv 14: 630–641.

Hein F (2000) Historical Overview of the Fort McMurray Area and Oil Sands Industry in Northeast Alberta (with expanded bibliographies on oil sands, surficial geology, hydrogeology, minerals and bedrock in Northeast Alberta). Earth Sciences Report 5: 32.

Henderson J, Nicol C, Lines J, White R, Wathes C (2001) Behaviour of domestic ducks exposed to mobile predator stimuli. 1. Flock responses. British Poult Sci 42: 433–438.

Hennan EG (1972) Status of waterfowl in the Peace-Athabasca Delta. Winnipeg, Manitoba: Ducks Unlimited (Canada).

Hennan EG (1973) Peace-Athabasca Delta: fall-staging census results, 1971-73. Winnipeg, Manitoba: Ducks Unlimited (Canada).

Hennan EG (1974) Peace-Athabasca Delta: breeding and fall- staging waterfowl census results, 1974. Winnipeg, Manitoba: Ducks Unlimited (Canada).

Hennan EG, Ambrock KR (1977) A Summary of Waterfowl Investigations in the Peace-Athabasca Delta, 1971-1976. Prepared for Parks Canada. Winnipeg, Manitoba: Ducks Unlimited (Canada) and Canadian Wildlife Service.

Hennan EG, Munson B (1979) Species Distribution and Habitat Relationships of Waterfowl in Northeastern Alberta; LS 22.1.2 Canadian Wildlife Service: Edmonton, Alberta, Canada.

Henny CJ, Hallock RJ, Hill EF (1994) Cyanide and migratory birds at gold mines in Nevada, USA. Ecotoxicology 3: 45–58.

Henry JD (1977) The use of urine marking in the scavenging behavior of the red fox (*Vulpes vulpes*). Behaviour: 82–106.

Herbert A (1970) Spatial disorientation in birds. Wilson Bull 82: 400–419.

Hernández L, Laundré JW (2005) Foraging in the 'landscape of fear' and its implications for habitat use and diet quality of elk *Cervus elaphus* and bison *Bison bison*. Widlife Biol 11: 215–220.

Hersikorn BD (2010) In-situ caged wood frog (*Rana sylvatica*) survival and development in wetlands formed from oil sands process-affected materials (OSPM).

Hersikorn BD, Ciborowski JJ, Smits JE (2010) The effects of oil sands wetlands on wood frogs (*Rana sylvatica*). Toxicol Environ Chem 92: 1513–1527.

Hersikorn BD, Smits JE (2011) Compromised metamorphosis and thyroid hormone changes in wood frogs (*Lithobates sylvaticus*) raised on reclaimed wetlands on the Athabasca oil sands. Environ Pollut 159: 596–601.

Hesse G, Rea RV, Booth AL (2010) Wildlife management practices at western Canadian airports. J Air Transp Manag16: 185–190.

Heubeck M, Camphuysen KCJ, Bao R, Humple D, Rey AS, et al. (2003) Assessing the impact of major oil spills on seabird populations. Mar Pollut Bull 46: 900–902.

Hewitt OH, Austin-Smith PJ (1966) A simple wing tag for field-marking birds. The J Widlife Manag: 625–627.

Heyers D, Manns M, Luksch H, Güntürkün O, Mouritsen H (2007) A visual pathway links brain structures active during magnetic compass orientation in migratory birds. PLoS ONE 2: e937.

Hilgerloh G, Bingman VP (1992) Radar observations of passerine migration over Frankfurt and Hannover, Germany. J Ornithol 133: 23–31.

Hilgerloh G, Caprano T, Griebeler EM (2010) Calibrating the operational beam width and maximum range of a ship radar used for bird observations. J Navig 63: 363–371.

Hill D, Fasham M, Tucker G, Shewry M, Shaw P (2005) Handbook of biodiversity methods: survey, evaluation and monitoring. Cambridge: Cambridge University Press. 589 p.

Hilton GM, Cresswell W, Ruxton GD (1999) Intraflock variation in the speed of escape-flight response on attack by an avian predator. Behav Ecol 10: 391–395.

Hilton GM, Ruxton GD, Cresswell W (1999) Choice of foraging area with respect to predation risk in redshanks: the effects of weather and predator activity. Oikos: 295–302.

Hinsley S, Hill R, Gaveau DLA, Bellamy PE (2002) Quantifying woodland structure and habitat quality for birds using airborne laser scanning. Funct Ecol 16: 851–857.

Hlady DA, Burger AE (1993) Drift-block experiments to analyse the mortality of oiled seabirds off Vancouver Island, British Columbia. Mar Pollut Bull 26: 495–501.

Ho CL, Karim H (1978) Impact of adsorbed petroleum hydrocarbons on marine organisms. Mar Pollut Bull 9: 156–162.

Hobson KA, Wilson AG, Van Wilgenburg SL, Bayne EM (2013) An estimate of nest loss in Canada due to industrial forestry operations. Avian Conserv Ecol 8: 5.

Hockin D, Ounsted M, Gorman M, Hill D, Keller V, et al. (1992) Examination of the effects of disturbance on birds with reference to its importance in ecological assessments. J Environ Manag 36: 253–286.

Hoen AL, Hagan M, Bratton M (2009) Burrowing owl and other migratory bird mitigation for a runway construction project at Edwards AFB. Hum-Wildl Interact 3: 251–256.

Hoffman DJ (1978) Embryotoxic effects of crude oil in mallard ducks and chicks. Toxicol Appl Pharmacol 46: 183–190.

Hoffman DJ (1979) Embryotoxic and teratogenic effects of crude oil on mallard embryos on day one of development. Bull Environ Contam Toxicol 22: 632–637.

Hoffman DJ (1979) Embryotoxic and teratogenic effects of petroleum hydrocarbons in mallards (*Anas platyrhynchos*). J Toxicol Environ Health 5: 835–844.

Hoffman DJ (1979) Embryotoxic effects of crude oil containing nickel and vanadium in Mallards. Bull Environ Contam Toxicol 23: 203–206.

Hoffman DJ, Albers PH (1984) Evaluation of potential embryotoxicity and teratogenicity of 42 herbicides, insecticides, and petroleum contaminants to mallard eggs. Arch Environ Contam Toxicol 13: 15–27.

Hoffman DJ, Eastin Jr WC, Gay ML (1982) Embryotoxic and biochemical effects of waste crankcase oil on birds' eggs. Toxicol Appl Pharmacol 63: 230–241.

Hoffman DJ, Gay ML (1981) Embryotoxic effects of benzo a pyrene, chrysene, and 7,12-dimethylbenz a anthracene in petroleum hydrocarbon mixtures in mallard ducks. J Toxicol Environ Health Part 7: 775–787.

Hoffmann L, Hafner H, Salathé T (1996) The contribution of colonial waterbird research to wetland conservation in the Mediterranean region. Colon Waterbird 19: 12–30.

Holevinski RA, Curtis PD, Malecki RA (2007) Hazing of Canada geese is unlikely to reduce nuisance populations in urban and suburban communities. Hum-Wildl Interact. Paper 97. Available: http://digitalcommons.unl.edu/hwi/97

Holleman I (2005) Quality control and verification of weather radar wind profiles. J Atmos Oceanic Technol 22: 1541–1550.

Holleman I, van Gasteren H, Bouten W (2008) Quality assessment of weather radar wind profiles during bird migration. J Atmos Oceanic Technol 25: 2188–2198.

Holmes W, Cavanaugh K (1990) Some evidence for an effect of ingested petroleum on the fertility of the mallard drake (*Anas platyrhynchos*). Arch Environ Contam Toxicol 19: 898–901.

Holmes W, Cavanaugh K, Cronshaw J (1978) The effects of ingested petroleum on oviposition and some aspects of reproduction in experimental colonies of mallard ducks (*Anas platyrhynchos*). J Reprod Fertil 54: 335–347.

Holmes W, Cronshaw J (1977) Biological effects of petroleum on marine birds. Effects of petroleum on arctic and subarctic marine environments and organisms 2: 359–398.

Holmes WN, Cronshaw J, Gorsline J (1978) Some effects of ingested petroleum on seawater-adapted ducks (*Anas platyrhynchos*). Environ Res 17: 177–190.

Holmes WN, Gorsline J, Cronshaw J (1979) Effects of mild cold stress on the survival of seawater-adapted mallard ducks (*Anas platyrhynchos*) maintained on food contaminated with petroleum. Environ Res 20: 425–444.

Holowenko FM, MacKinnon MD, Fedorak PM (2002) Characterization of naphthenic acids in oil sands wastewaters by gas chromatography-mass spectrometry. Water Res 36: 2843–2855.

Holroyd P, Simieritsch T (2009) The waters that bind us: transboundary implications of oil sands development. Drayton Valley, Alberta: Pembina Institute for Appropriate Development.

Holt M, Fox K, Grießbach E, Johnsen S, Kinnunen J, et al. (2000) Monitoring, modelling and environmental exposure assessment of industrial chemicals in the aquatic environment. Chemosphere 41: 1799–1808.

Homan HJ, Slowik A, Blackwell B, Linz G (2010) Field Testing Class IIIb Handheld Lasers to Disperse Roosting Blackbirds. In: National Sunflower Association Sunflower Research Forum, Fargo.

Honda T (2012) Line color affects the collision risk and deterrence of crows. J Ethology 30: 11–14.

Honza M, Pola\vciková L (2008) Experimental reduction of ultraviolet wavelengths reflected from parasitic eggs affects rejection behaviour in the blackcap *Sylvia atricapilla*. J Exp Biol 211: 2519–2523.

Hood GA, Bayley SE (2008) Beaver (*Castor canadensis*) mitigate the effects of climate on the area of open water in boreal wetlands in western Canada. Biol Cons 141: 556–567.

Hoover AK, Rohwer FC, Richkus KD (2004) Evaluation of nest temperatures to assess female nest attendance and use of video cameras to monitor incubating waterfowl. Wildlife Soc B 32: 581–587.

Horváth G, Bernáth B, Molnár G (1998) Dragonflies find crude oil visually more attractive than water: multiple-choice experiments on dragonfly polarotaxis. Naturwissenschaften 85: 292–297.

Horváth G, Kriska G, Malik P, Robertson B (2009) Polarized light pollution: a new kind of ecological photopollution. Front Ecol Environ 7: 317–325.

Horváth G, Varjú D (2004) Polarized light in animal vision: polarization patterns in nature. Heidelberg: Springer. 447 p.

Houk AP, Delwiche MJ, Gorenzel WP, Salmon TP (2005) Electronic broadcast call unit for bird control in orchards. Appl Eng Agric 21: 721-727.

Hounsell R, Reilly TJ (1995) New Brunswick bird deterrent study. Marine Spill Response Corporation. In: 1997 International oil spill conference, Fort Lauderdale.

Howell S, Clarke A, Freitag S, McNaughton C, Kapustin V, et al. (2013) An airborne assessment of atmospheric particulate emissions from the processing of Athabasca oil sands. Atmos Chem Phys Discuss 13: 21301–21343.

Howlett M, Craft J (2013) Application of federal legislation to Alberta's mineable oil sands. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-33. 94 pp.

Hsu CS, Dechert G, Robbins W, Fukuda E (2000) Naphthenic acids in crude oils characterized by mass spectrometry. Energy Fuels 14: 217–223.

Hudson M-AR, Bird DM (2009) Recommendations for design and management of golf courses and green spaces based on surveys of breeding bird communities in Montreal. Landsc Urban Plan 92: 335–346.

Huffman S (1999) The effects of oil on mallards (Anas platyrhynchos). Tex J Sci 51: 181–190.

Hughes MR, Kasserra C, Thomas BR (1990) Effect of externally applied bunker fuel on body mass and temperature, plasma concentration, and water flux of Glaucous-winged Gulls, *Larus glaucescens*. Can J Zool 68: 716–721.

Humple DL, Nevins HM, Phillips EM, Gibble C, Henkel L, et al. (2011) Demographics of Aechmophorus grebes killed in three mortality events in California. Mar Ornithol 39: 235–242.

Hüppop O, Hilgerloh G (2012) Flight call rates of migrating thrushes: effects of wind conditions, humidity and time of day at an illuminated offshore platform. J Avian Biol 43: 85–90.

Hutto RL, Stutzman RJ (2009) Humans versus autonomous recording units: a comparison of point-count results. J Field Ornithol 80: 387–398.

Hwang S, Cutright TJ (2004) Preliminary exploration of the relationships between soil characteristics and PAH desorption and biodegradation. Environ Int 29: 887–894.

Hwang S, Joe Y, Cha H, Joo G, Lee H, et al. (2013) Direct Look from a Predator Shortens the Risk-Assessment Time by Prey. PLoS ONE 8: e64977.

HydroQual Laboratories Ltd. (1996) Laboratory Studies on Trophic Level Effects and Fish Health Effects of Suncor Tar Island Dyke Wastewater. Prepared for Suncor Inc. Calgary, Alberta: HydroQual Laboratories Ltd.

Imber M (1975) Behaviour of petrels in relation to the moon and artificial lights. Notornis 22: 302-306.

Irons DB, Kendall SJ, Erickson WP, McDonald LL, Lance BK (2000) Nine years after the Exxon Valdez oil spill: effects on marine bird populations in Prince William Sound, Alaska. Condor 102: 723–737.

Islam T, Rico-Ramirez MA, Han D, Srivastava PK (2012) Artificial intelligence techniques for clutter identification with polarimetric radar signatures. Atmos Res 109: 95–113.

Iverson SA, Esler D (2010) Harlequin Duck population injury and recovery dynamics following the 1989 Exxon Valdez oil spill. Ecol Appl 20: 1993–2006.

Jachowski D, Slotow R, Millspaugh J (2013) Good virtual fences make good neighbors: opportunities for conservation. Anim Conserv. doi:10.1111/acv.12082

Jacobsen OW, Ugelvik M (1992) Anti-predator behavior of breeding Eurasian Wigeon. J Field Ornithol 63: 324–330.

Jacobson HA, Kroll JC, Browning RW, Koerth BH, Conway MH (1997) Infrared-triggered cameras for censusing white-tailed deer. Wildlife Soc B 25: 547-556.

Jaime M, Lopez JP, Lickliter R (2009) Bobwhite quail (*Colinus virginianus*) hatchlings track the direction of human gaze. Anim Cogn 12: 559–565.

James D, Vold T (2010) Establishing a world class public information and reporting system for ecosystems in the oil sands region: report and appendices. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-5. 189 pp.

James JB, Hellgren EC, Masters RE (1999) Effects of deterrents on avian abundance and nesting density in electrical substations in Oklahoma. J Widlife Manag 63: 1009–1017.

James W (1994) Water Quality Review and Treatment Recommendations for Water Released From Suncor's Tailings: FinalReport. Prepared for CANMET Energy.

Jantzen C, Slagle R (1997) Investigations of high-speed digital imaging of low-light-level events using pulsed near-infrared laser light sources. 22nd Int'l Congress on High-Speed Photography and Photonics. pp. 827–833.

Jantzie T, Croft B, Coutts R, Biggs D, Loman A (1979) Proposed design for a program of toxicological research for the Alberta oil sands environmental research program. AOSERP Program Management Report PM-3; RMD report L 41.

Jehl Jr JR (1993) Observations on the fall migration of eared grebes, based on evidence from a mass downing in Utah. Condor 95: 470–473.

Jehl Jr JR (1996) Mass mortality events of eared grebes in North America. J Field Ornithol 67: 471–476.

Jenkins KJ, Manly B (2008) A double-observer method for reducing bias in faecal pellet surveys of forest ungulates. J Appl Ecol 45: 1339–1348.

Jenssen BM (1989) Effects of ingested crude and dispersed crude oil on thermoregulation in ducks (*Anas platyrhynchos*). Environ Res 48: 49–56.

Jenssen BM (1994) Review article: effects of oil pollution, chemically treated oil, and cleaning on the thermal balance of birds. Environ Pollut 86: 207–215.

Jenssen BM, Ekker M (1988) A method for evaluating the cleaning of oiled seabirds. Wildlife Soc B 16: 213–215.

Jenssen BM, Ekker M (1989) Rehabilitation of oiled birds: a physiological evaluation of four cleaning agents. Mar Pollut Bull 20: 509–512.

Jenssen BM, Ekker M (1991) Dose dependent effects of plumage-oiling on thermoregulation of Common Eiders *Somateria mollissima* residing in water. Polar Res10: 579–584.

Jenssen BM, Ekker M (1991) Effects of plumage contamination with crude oil dispersant mixtures on thermoregulation in common eiders and mallards. Arch Environ Contam Toxicol 20: 398–403.

Jenssen BM, Ekker M, Bech C (1985) Thermoregulation in a naturally oil-contaminated Black-billed Murre *Uria aalge*. Bull Environ Contam Toxicol 35: 9–14.

Jenssen BM, Ekker M, Zahlsen K (1990) Effects of ingested crude oil on thyroid hormones and on the mixed function oxidase system in ducks. Comp Biochem Physiol C Pharmacol Comp Pharmacol 95: 213–216.

Jetz W, Steffen J, Linsenmair KE (2003) Effects of light and prey availability on nocturnal, lunar and seasonal activity of tropical nightjars. Oikos 103: 627–639.

Joensen AH (1972) Oil pollution and seabirds in Denmark 1935-1968. Danish Rev Game Biol 6: 1–24.

Johansson CA, Hardin PJ, White CM (1997) An inexpensive fully automated system reduces avian landings on a 45 acre "defended" pond by 97%. Washington, DC: US Fish and Wildlife Service.

Johnson DH (2008) In defense of indices: the case of bird surveys. The J Widlife Manag 72: 857–868.

Johnson E, Miyanishi K (2008) Creating new landscapes and ecosystems. Anna NY Acad Sci 1134: 120–145.

Johnson SR (1998) Distribution and movements of brood-rearing lesser snow geese in relation to petroleum development in Arctic Alaska. Arctic: 336–344.

Johnson WP, Baar L, Matlack RS, Barron RB (2010) Hatching chronology of ducks using playas in the southern high plains of Texas. Am Midl Nat 163: 247–253.

Johnston DW, Haines T (1957) Analysis of mass bird mortality in October, 1954. Auk 4: 447–458.

Johnston NN, Bradley JE, Pomeroy AC, Otter KA (2013) Flight Paths of Migrating Golden Eagles and the Risk Associated with Wind Energy Development in the Rocky Mountains. Avian Conserv Ecol 8: 12.

Jones D, Scarlett AG, West CE, Rowland SJ (2011) Toxicity of individual naphthenic acids to *Vibrio fischeri*. Environ Sci Technol 45: 9776–9782.

Jones J, Francis CM (2003) The effects of light characteristics on avian mortality at lighthouses. J Avian Biol 34: 328–333.

Jones NF, Pejchar L (2013) Comparing the Ecological Impacts of Wind and Oil & Gas Development: A Landscape Scale Assessment. PLoS ONE 8(11): e81391. doi:10.1371/journal.pone.0081391

Jones PH, Monnat JY, Cadbury C, Stowe T (1978) Birds oiled during the Amoco Cadiz incident—An interim report. Mar Pollut Bull 9: 307–310.

Jones R (1980) Reactions of male domestic chicks to two-dimensional eye-like shapes. Anim Behav 28: 212–218.

Jones R, Forrest D (2010) Oil sands mining reclamation challenge dialogue: report and appendices. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-4A. 18 pp.

Jordaan SM, Keith DW, Stelfox B (2009) Quantifying land use of oil sands production: a life cycle perspective. Environ Res Lett 4: 024004.

Jørgensen LF, Refsgaard JC, Højberg AL (2007) The inadequacy of monitoring without modelling support. J Environ Monit 9: 931–942.

Journey L, Drury JP, Haymer M, Rose K, Blumstein DT (2013) Vivid birds respond more to acoustic signals of predators. Behav Ecol Sociobiol 67: 1285-1293.

Juel A, Ejrnæs R, Fredshavn J, Groom G (2012) Integrating field survey and orthophoto information to monitor coastal habitats - A pilot study to develop methods and resolve key issues. Ecol Inform 14:48-52.

Kahlert J, Fox AD, Ettrup H (1996) Nocturnal feeding in moulting Greylag Geese Anser anseran anti-predator response. Ardea 84: 15–22.

Kalal Z, Mikolajczyk K, Matas J (2010) Face-TLD: Tracking-Learning-Detection applied to faces. Image Processing (ICIP), 2010 17th IEEE International Conference on. pp. 3789–3792.

Kalal Z, Mikolajczyk K, Matas J (2010) Forward-backward error: Automatic detection of tracking failures. Pattern Recognition (ICPR), 2010 20th International Conference on. pp. 2756–2759.

Kalies EL, Chambers CL, Covington WW (2010) Wildlife responses to thinning and burning treatments in southwestern conifer forests: A meta-analysis. Forest Ecol Manag 259: 333–342.

Kammerer M, Mastain O, Le Dréan-Quenech'du S, Pouliquen H, Larhantec M (2004) Liver and kidney concentrations of vanadium in oiled seabirds after the Erika wreck. Sci Total Environ 333: 295–301.

Kannel PR, Gan TY (2012) Naphthenic acids degradation and toxicity mitigation in tailings wastewater systems and aquatic environments: A review. J Environ Sci Health A 47: 1–21.

Kannel PR, Gan TY (2013) Application of WASP for Modelling and Management of Naphthenic Acids along Athabasca River, Alberta, Canada. Water Air Soil Pollut 224: 1764.

Karanth KU, Nichols JD (1998) Estimation of tiger densities in India using photographic captures and recaptures. Ecology 79: 2852–2862.

Karanth KU, Nichols JD, Seidenstricker J, Dinerstein E, Smith JLD, et al. (2003) Science deficiency in conservation practice: the monitoring of tiger populations in India. Anim Conserv 6: 141–146.

Kardynal K, Morissette J, Van Wilgenburg S, Bayne E, Hobson K (2011) Avian responses to experimental harvest in southern boreal mixedwood shoreline forests: implications for riparian buffer management. Can J Forest Res 41: 2375–2388.

Karlsson H, Bäckman J, Nilsson C, Alerstam T (2010) Exaggerated orientation scatter of nocturnal passerine migrants close to breeding grounds: comparisons between seasons and latitudes. Behav Ecol and Sociobiology 64: 2021–2031.

Karlsson H, Nilsson C, Bäckman J, Alerstam T (2011) Nocturnal passerine migration without tailwind assistance. Ibis 153: 485–493.

Karlsson H, Nilsson C, Bäckman J, Alerstam T (2012) Nocturnal passerine migrants fly faster in spring than in autumn: a test of the time minimization hypothesis. Anim Behav 83: 87–93.

Karr JR (1987) Biological monitoring and environmental assessment: a conceptual framework. Environ Manag 11: 249–256.

Karr JR, Dudley DR (1981) Ecological perspective on water quality goals. Environ Manag 5: 55–68.

Kavanagh RJ (2012) The Effects of Oil Sands Process-Affected Waters and their Associated Constituents on Fathead Minnow (*Pimephales promelas*) Reproductive Physiology. Doctor of Philosophy, University of Guelph. Guelph, Canada.

Kavanagh RJ, Frank RA, Burnison BK, Young RF, Fedorak PM, et al. (2012) Fathead minnow (*Pimephales promelas*) reproduction is impaired when exposed to a naphthenic acid extract. Aquat Toxicol 116: 34–42.

Kelly EN, Schindler DW, Hodson PV, Short JW, Radmanovich R, et al. (2010) Oil sands development contributes elements toxic at low concentrations to the Athabasca River and its tributaries. PNAS 107: 16178–16183.

Kelly EN, Short JW, Schindler DW, Hodson PV, Ma M, et al. (2009) Oil sands development contributes polycyclic aromatic compounds to the Athabasca River and its tributaries. PNAS 106: 22346–22351.

Kelly T, West T, Davenport J (2009) Challenges and solutions of remote sensing at offshore wind energy developments. Mar Pollut Bull 58: 1599–1604.

Kelly TA (2009) Beware of the birds. Potentials, IEEE 28: 31–35.

Kennedy HL (2008) Comparison of MHT and PDA track initiation performance. Radar, 2008 International Conference on. pp. 508–512.

Kennicutt MCI (1990) Oil spillage in Antarctica. Environ Sci Technol 24: 620–624.

Kerley G, Crellin C, Erasmus T (1987) Gravimetric determination of water-repellancy in rehabilitated oiled seabirds. Mar Pollut Bull 18: 609–611.

Kerley G, Erasmus T (1986) Oil pollution of Cape gannets: To clean or not to clean? Mar Pollut Bull 17: 498–500.

Kerley GIH, Erasmus T (1987) Cleaning and rehabilitation of oiled Jackass Penguins. S Afr J Wildl Res 17: 64–70.

Kerley GIH, Erasmus T (1987) The management of oiled penguins. Proc. 1987 Oil Spill Conf. (Prevention, Behaviour, Control, Cleanup) Amer. Petroleum Inst., Washington 4452: 465–468.

Kerlinger P (2000) Avian mortality at communication towers: a review of recent literature, research, and methodology. Prepared for: United States Fish and Wildlife Service Office of Migratory Bird Management.

Kerlinger P, Gauthreaux S (1985) Flight behavior of raptors during spring migration in south Texas studied with radar and visual observations. J Field Ornithol 56: 394–402.

Kertész V, Hlubik I (2002) Plasma ALP activity and blood PCV value changes in chick fetuses due to exposure of the egg to different xenobiotics. Environ Pollut 117: 323–327.

Kéry M, Royle JA, Schmid H (2005) Modeling avian abundance from replicated counts using binomial mixture models. Ecol Appl 15: 1450–1461.

Kessler WB, Booth AL (1998) Professor Leopold, what is education for? Wildlife Soc B 26: 707–712.

Khan R, Nag K (1993) Estimation of hemosiderosis in seabirds and fish exposed to petroleum. Bull Environ Contam Toxicol 50: 125–131.

Khan R, Ryan P (1991) Long term effects of crude oil on common murres (*Uria aalge*) following rehabilitation. Bull Environ Contam Toxicol 46: 216–222.

Kight CR, Swaddle JP (2011) How and why environmental noise impacts animals: an integrative, mechanistic review. Ecol Lett 14: 1052–1061.

Kim C-H, Jeung K-M, Choi Y-S, Lee S-U, Kim C-H, et al. (2011) Design of the unmanned ground vehicle for bird expellant in an airport. Control, Automation and Systems (ICCAS), 2011 11th International Conference on. pp.1094–1097.

Kimmins J, Rempel R, Welham CVJ, Seely B, Van Rees KCJ (2007) Biophysical sustainability, process-based monitoring and forest ecosystem management decision support systems. Forest Chron 83: 502–514.

Kindzierski W, El-Din MG, Jin J (2012) Review of health effects of naphthenic acids: data gaps and implications for understanding human health risk. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-20. 43 pp.

Kindzierski W, Jin J, El-Din MG (2011) Plain language explanation of human health risk assessment. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-14. 37 pp.

King DI, DeGraaf RM (2006) Predators at bird nests in a northern hardwood forest in New Hampshire. J Field Ornithol 77: 239–243.

King JR, Bendell-Young LI (2000) Toxicological significance of grit replacement times for juvenile mallards. J Wildl Manage 64: 858–862.

King KA, Lefever CA (1979) Effects of oil transferred from incubating gulls to their eggs. Mar Pollut Bull 10: 319–321.

King KA, Macko S, Parker PL, Payne E (1979) Resuspension of oil: probable cause of brown pelican fatality. Bull Environ Contam Toxicol 23: 800–805.

King KA, Stafford CJ, Cain BW, Mueller AJ, Hall HD (1987) Industrial, agricultural, and petroleum contaminants in cormorants wintering near the Houston Ship Channel, Texas, USA. Colon Waterbird 10: 93–99.

Kirschel AN, Cody ML, Harlow ZT, Promponas VJ, Vallejo EE, et al. (2011) Territorial dynamics of Mexican Ant-thrushes Formicarius moniliger revealed by individual recognition of their songs. Ibis 153: 255–268.

Kissane C (2013) Reduction of wildlife on airfields through turfgrass selection. Master of Science. University of Illinois. Urbana-Champaign, USA.

Kistler C, Hegglin D, von Wattenwyl K, Bontadina F (2013) Is electric fencing an efficient and animal-friendly tool to prevent stone martens from entering buildings? European J Wildl Res: 1–5.

Klaassen M, Biebach H (2003) Flight altitude of trans-Sahara migrants in autumn: a comparison of radar observations with predictions from meteorological conditions and water and energy balance models. J Avian Biol 31: 47–55.

Klem D, Groot KLD, Krebs EA, Fort KT, Elbin SB, Prince A (2012) A second critique of "Collision mortality has no discernable effect on population trends of North American Birds." PLoS ONE 6.

Klope MW, Beason RC, Nohara TJ, Begier MJ (2009) Role of near-miss bird strikes in assessing hazards. Hum-Wildl Interact 3: 208–215.

Knag AC, Sebire M, Mayer I, Meier S, Renner P, Katsiadaki I (2013) In vivo endocrine effects of naphthenic acids in fish. Chemosphere 93: 2356-2364.

Knight RL, Kadlec RH, Ohlendorf HM (1999) The use of treatment wetlands for petroleum industry effluents. Environ Sci Technol 33: 973–980.

Koistinen J (2000) Bird migration patterns on weather radars. Phys Chem Earth B 25: 1185–1193.

Kopischke ED (1972) The effect of 2, 4-D and diesel fuel on egg hatchability. The J Widlife Manag: 1353–1356.

Korosi J, Irvine G, Skierszkan E, Doyle J, Kimpe L, et al. (2013) Localized enrichment of polycyclic aromatic hydrocarbons in soil, spruce needles, and lake sediments linked to in-situ bitumen extraction near Cold Lake, Alberta. Environ Pollut 182: 307–315.

Koski WR, Kevan SD, Richardson WJ (1993) Bird dispersal and deterrent techniques for oil spills in the Beaufort Sea.

Koyuncu T, Lule F (n.d.) Design, Manufacture and Test of a Solar Powered Audible Bird Scarer.

Kozar J (2011) Historical avian deterrence at Suncor Energy Inc. Lease No. 86 from 1977 to 1992.

Kramer DL, Bonenfant M (1997) Direction of predator approach and the decision to flee to a refuge. Anim Behav54: 289–295.

Kreutzweiser D, Beall F, Webster K, Thompson D, Creed I (2013) Impacts and prognosis of natural resource development on aquatic biodiversity in Canada's boreal zone. Environ Rev 21: 227–259.

Kriska G, Barta A, Suhai B, Bernáth B, Horváth G (2008) Do brown pelicans mistake asphalt roads for water in deserts. Acta Zool Acad Sci H 54: 157–165.

Kross SM, Nelson XJ (2011) A portable low-cost remote videography system for monitoring wildlife. Methods Ecol Evol 2: 191–196.

Kullberg C, Henshaw I, Jakobsson S, Johansson P, Fransson T (2007) Fuelling decisions in migratory birds: geomagnetic cues override the seasonal effect. Proc R Soc B Biol Sci 274: 2145–2151.

Kunz TH, Arnett EB, Cooper BM, Erickson WP, Larkin RP, et al. (2007) Assessing impacts of wind-energy development on nocturnally active birds and bats: a guidance document. The J Widlife Manag 71: 2449–2486.

Kurek J, Kirk JL, Muir DC, Wang X, Evans MS, et al. (2013) Legacy of a half century of Athabasca oil sands development recorded by lake ecosystems. PNAS 110: 1761–1766.

Köth T, Vauk-Hentzelt E (1988) Influence of plumage and stomach oiling on body and organ growth in young kittiwakes. Mar Pollut Bull 19: 71–73.

Lack D, Varley G (1945) Detection of birds by radar. Nature 156: 446.

Laffon B, Aguilera F, Ríos-Vázquez J, Valdiglesias V, Pásaro E (2013) Follow-up study of genotoxic effects in individuals exposed to oil from the tanker Prestige, seven years after the accident. Mutat Res.

Lai JWS, Pinto L, Kiehlmann E, Bendell-Young LL, Moore MM (1995) The mineralization of naphthenic acids in Oil Sands wastewater by indigenous microbial communities.

Lai JWS, Pinto LJ, Kiehlmann E, Bendell-Young LI, Moore MM (1995) Determining the Ecological Viability of Constructed Wetlands for the Treatment of Oil Sands Wastewater.

Lakshmanan V, Zhang J, Howard K (2010) A technique to censor biological echoes in radar reflectivity data. J Appl Met Climatol 49: 453–462.

Laliberte AS, Ripple WJ (2003) Automated wildlife counts from remotely sensed imagery. Wildlife Soc B: 362–371.

Lamb EG, Bayne E, Holloway G, Schieck J, Boutin S, et al. (2009) Indices for monitoring biodiversity change: Are some more effective than others? Ecol Indic 9: 432–444.

Lambert G, Peakall D, Philogene B, Engelhardt F (1982) Effect of oil and oil dispersant mixtures on the basal metabolic rate of ducks. Bull Environ Contam Toxicol 29: 520–524.

Lance BK, Irons DB, Kendall SJ, McDonald LL (2001) An evaluation of marine bird population trends following the Exxon Valdez oil spill, Prince William Sound, Alaska. Mar Pollut Bull 42: 298–309.

Lanctot R, Goatcher B, Scribner K, Talbot S, Pierson B, et al. (1999) Harlequin duck recovery from the Exxon Valdez oil spill: a population genetics perspective. Auk: 781–791.

Landowne R, Wainright R (1971) A chemical tagging system for use in the prevention of oil spills. American Cyanamid Company Central Research Division: 179–182.

Langton R, Davies I, Scott B (2011) Seabird conservation and tidal stream and wave power generation: Information needs for predicting and managing potential impacts. Mar Policy 35: 623–630.

Larkin RP (1991) Flight speeds observed with radar, a correction: slow "birds" are insects. Behav Ecol and Sociobiology 29: 221–224.

Larkin RP (2012) Radar techniques for Widlife Biol. In: Nova J. Silvy, editor. Techniques for wildlife investigations and management. Baltimore: John Hopkins University Press, Vol. 1. pp. 448–464.

Larkin RP, Frase BA (1988) Circular paths of birds flying near a broadcasting tower in cloud. J Comp Psychol 102: 90-93.

Larkin RP, Sutherland PJ (1977) Migrating birds respond to Project Seafarer's electromagnetic field. Science 195: 777–779.

Larkin RP, Szafoni RE (2008) Evidence for widely dispersed birds migrating together at night. Integr Comp Biol 48: 40–49.

Larkin RP, Thompson D (1980) Flight speeds of birds observed with radar: evidence for two phases of migratory flight. Behav Ecol and Sociobiology 7: 301–317.

Larsen EM, Richardson SA (1990) Some effects of a major oil spill on wintering shorebirds at Grays Harbor, Washington. Northwestern Naturalist: 88–92.

Larsson C, Gustafsson M, Kristensson G (2010) Calibration methods for wideband forward RCS measurements. Antennas and Propagation (EuCAP), 2010 Proceedings of the Fourth European Conference on. pp. 1–5.

Laundré JW, Hernández L, Ripple WJ (2010) The landscape of fear: ecological implications of being afraid. Open Ecol J 3: 1–7.

Lawler G, Holmes J, Adamkiewicz D, Shields M, Monnat J, et al. (1979) Characterization of petroleum hydrocarbons in tissues of birds killed by the Amoco Cadiz oil spill. Amoco Cadiz-Conséquences d'une pollution accidentelle par les hydrocarbures, Actes du Colloque International Centre Océanologique de Bretagne Brest. Vol. 19. p. 22.

Lawler G, Holmes J, Fiorito B, Laseter J, Szaro R (1978) Quantification of petroleum hydrocarbons in selected tissues of male mallard ducklings chronically exposed to South Louisiana crude oil. Conference on Assessment of Ecological Impacts of Oil Spills" American Institute of Biological Sciences, Keystone, Colorado.

Lawler GC, Loong WA, Laseter JL (1978) Accumulation of saturated hydrocarbons in tissues of petroleum-exposed mallard ducks (*Anas platyrhynchos*). Environ Sci Technol 12: 47–51.

Lawrence GA, Ward PRB, Yassien HA (1996) Potential for formation of a permanent halocine in the base mine lake.

Leask M (1977) A physicochemical mechanism for magnetic field detection by migratory birds and homing pigeons. Nature 267: 144–145.

Lebeuf AP, Giroux J-F (2013) Sibling pairing in temperate-nesting Canada Geese. Wilson J Ornithol 125: 398–401.

Leclair LA, MacDonald GZ, Phalen LJ, Köllner B, Hogan NS, et al. (2013) the immunological effects of oilsands surface waters and naphthenic acids on rainbow trout (*Oncorhynchus mykiss*). Aquat Toxicol.142: 185–194.

Lee Jr RC (1990) Bird kills in contained oil: a biopolitical/enforcement strategy. Proceedings of the Annual Conference of Southeastern Association of Fish and Wildlife Agencies. Vol. 44. pp. 444–447.

Lee P, Boutin S (2006) Persistence and developmental transition of wide seismic lines in the western Boreal Plains of Canada. J Environ Manag 78: 240–250.

Lee PG (2011) Migration of whooping cranes (*Grus americana*) through Alberta's bitumen sands region.

Lee W-H, Gader PD, Wilson JN (2007) Optimizing the area under a receiver operating characteristic curve with application to landmine detection. IEEE Trans Geosci Remote Sens 45: 389–397.

Lee YZ, Leighton F, Peakall D, Norstrom R, O'Brien P, et al. (1985) Effects of ingestion of Hibernia and Prudhoe Bay crude oils on hepatic and renal mixed function oxidase in nestling herring gulls (*Larus argentatus*). Environ Res 36: 248–255.

Legg CJ, Nagy L (2006) Why most conservation monitoring is, but need not be, a waste of time. J Environ Manag 78: 194–199.

Leggett MA (2012) Managing populations of the Australasian harrier (*Circus approximans*) to reduce passerine bird damage in vineyards.

Lehman RN, Kennedy PL, Savidge JA (2007) The state of the art in raptor electrocution research: a global review. Biol Cons 136: 159–174.

Lehmann V (2012) Optimal Gabor-frame-expansion-based intermittent-clutter-filtering method for radar wind profiler. J Atmos Oceanic Technol 29: 141–158.

Lehoux D, Bordage D (2000) Deterrent Techniques and Bird Dispersal Approach for Oil Spills. Environment Canada, Canadian Wildlife Service.

Leibowitz SG (2003) Isolated wetlands and their functions: an ecological perspective. Wetlands 23: 517–531.

Leighton F (1985) Morphological lesions in red blood cells from herring gulls and Atlantic puffins ingesting Prudhoe Bay Crude Oil. Vet Pathol 22: 393–402.

Leighton FA (1986) Clinical, gross, and histological findings in herring gulls and atlantic puffins that ingested Prudhoe Bay crude oil. Vet Pathol 23: 254.

Leighton FA (1993) The toxicity of petroleum oil to birds. Environ Rev 1: 92–103.

Leighton FA, Lee YZ, Rahimtula AD, O'Brien PJ, Peakall DB (1985) Biochemical and functional disturbances in red blood cells of herring gulls ingesting Prudhoe Bay crude oil. Toxicol Appl Pharmacol 81: 25–31.

Leighton FA, Peakall DB, Butler RG (1983) Heinz-body hemolytic anemia from the ingestion of crude oil: a primary toxic effect in marine birds. Science 220: 871–873.

Lele SR, Moreno M, Bayne E (2012) Dealing with detection error in site occupancy surveys: what can we do with a single survey? J Plant Ecol 5: 22–31.

Leonhardt CL (2003) Zoobenthic succession in constructed wetlands of the Fort McMurray oil sands region: Developing a measure of zoobenthic recovery.

Lepage D, Francis CM (2002) Do feeder counts reliably indicate bird population changes? 21 years of winter bird counts in Ontario, Canada. Condor 104: 255–270.

Leskiw LA, Ag P (1998) Land and Capability Classification for Forest Ecossytems in the Oil Sands Region.

Leung SS, MacKinnon MD, Smith RE (2003) The ecological effects of naphthenic acids and salts on phytoplankton from the Athabasca oil sands region. Aquat Toxicol 62: 11–26.

Levy E (1980) Oil pollution and seabirds: Atlantic Canada 1976-1977 and some implications for northern environments. Mar Pollut Bull 11: 51–56.

Levy E (1983) Commentary: what impact will the oil industry have on seabirds in the Canadian Arctic? Arctic 30: 1–4.

Lewis HF (1942) Destruction of waterfowl by oil. Wilson Bull 54: 217.

Lewis P, Perry G, Morris T (2000) Ultraviolet radiation and laying pullets. British Poult Sci41: 131–135.

Lewis SJ, Malecki RA (1984) Effects of egg oiling on larid productivity and population dynamics. Auk: 584–592.

Lewis SJ, Malecki RA (1985) Reproductive success of great black-backed and herring gulls in response to egg oiling. New York Cooperative Wildlife Res Unit, Dept. of Natural Resources, Cornell University.

Lewison RL, Crowder LB (2003) Estimating fishery bycatch and effects on a vulnerable seabird population. Ecol Appl 13: 743–753.

Li S, McShea WJ, Wang D, Shao L, Shi X (2010) The use of infrared-triggered cameras for surveying phasianids in Sichuan Province, China. Ibis 152: 299–309.

Li X, Feng Y (1995) Dewatering fine tails by evaporation: A mathematical modelling approach.

Lichtenstein G, Sealy SG (1998) Nestling competition, rather than supernormal stimulus, explains the success of parasitic brown-headed cowbird chicks in yellow warbler nests. Roc R Soc Lond B Biol Sci 265: 249–254.

Liechti F, Bruderer B, Paproth H (1995) Quantification of nocturnal bird migration by moonwatching: comparison with radar and infrared observations. J Field Ornithol 66: 457–468.

Liechti F, Guélat J, Komenda-Zehnder S (2013) Modelling the spatial concentrations of bird migration to assess conflicts with wind turbines. Biol Cons 162: 24–32.

Lilley MB, Firestone J (2008) Wind power, wildlife, and the Migratory Bird Treaty Act: a way forward. Envtl L 38: 1167.

Lima SL (1995) Collective detection of predatory attack by social foragers: fraught with ambiguity? Anim Behav 50: 1097–1108.

Lima SL (1998) Stress and decision making under the risk of predation: recent developments from behavioral, reproductive, and ecological perspectives. Adv Study Behav 27: 215–290.

Lima SL, Dill LM (1990) Behavioral decisions made under the risk of predation: a review and prospectus. Canadian J Zool 68: 619–640.

Lindenmayer DB, Gibbons P, Bourke M, Burgman M, Dickman CR, et al. (2012) Improving biodiversity monitoring. Austral Ecol 37: 285–294.

Lindenmayer DB, Likens GE (2009) Adaptive monitoring: a new paradigm for long-term research and monitoring. Trends Ecol Evol 24: 482–486.

Lindenmayer DB, Likens GE (2010) Improving ecological monitoring. Trends Ecol Evol 25: 200–201.

Lindenmayer DB, Likens GE (2010) The science and application of ecological monitoring. Biol Cons 143: 1317–1328.

Lindstedt-Siva J, Albers PH, Fucik KW, Maynard NG (1984) Ecological considerations for the use of dispersants in oil spill response. Oil Spill Chemical Dispersants: Research, Experience, and Recommendations American Socieity for Testing and Materials, Philadelphia, PA STP 840: 363–377.

Link WA, Sauer JR (1998) Estimating population change from count data: application to the North American breeding bird survey. Ecol Appl 8: 258–268.

Lintelmann J, Katayama A, Kurihara N, Shore L, Wenzel A (2003) Endocrine disruptors in the environment (IUPAC Technical Report). Pure Appl Chem 75: 631–681.

Linz GM, Homan HJ, Werner SJ, Hagy HM, Bleier WJ (2011) Assessment of bird-management strategies to protect sunflowers. Biosci 61: 960–970.

Lister A, Nero V, Farwell A, Dixon D, Van Der Kraak G (2008) Reproductive and stress hormone levels in goldfish (*Carassius auratus*) exposed to oil sands process-affected water. Aquat Toxicol 87: 170–177.

Lokemoen JT, Woodward RO (1993) An assessment of predator barriers and predator control to enhance duck nest success on peninsulas. Wildlife Soc B: 275–282.

Longcore T, Rich C (2004) Ecological light pollution. Front Ecol Environ 2: 191–198.

Longcore T, Rich C, Mineau P, MacDonald B, Bert DG, et al. (2012) An estimate of avian mortality at communication towers in the United States and Canada. PLoS ONE 7: e34025.

Longcore T, Rich C, Mineau P, MacDonald B, Bert DG, et al. (2013) Avian mortality at communication towers in the United States and Canada: which species, how many, and where? Biol Cons 158: 410–419.

Longcore T, Smith PA (2013) On Avian Mortality Associated with Human Activities.

Loots S (2014) Automated monitoring of flying and landed birds at industrial ponds.

Lord DA, Kerley GIH, Randall RM, Reddering JSV, Schumann EH, et al. (1986) The Kapodistrias grounding and oil spill, Cape Recife, South Africa. Institute for Coastal Research, University of Port Elizabeth.

Lorimer H (2013) Scaring Crows. Geogr Rev 103: 177–189.

Lott EO, Jones RK (2010) Review of four major environmental effects monitoring programs in the oil sands area. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-6. 114 pp.

Lotz A, Allen CR (2007) Observer bias in anuran call surveys. The J Widlife Manag 71: 675–679.

Lowery GH (1949) A quantitative study of the nocturnal migration of birds.

Lowery GH, Newman RJ (1966) A continentwide view of bird migration on four nights in October. Auk 83: 547–586.

Lucas Z, MacGregor C (2006) Characterization and source of oil contamination on the beaches and seabird corpses, Sable Island, Nova Scotia, 1996-2005. Mar Pollut Bull 52: 778–789.

Lusimbo WS, Leighton FA (1996) Effects of Prudhoe Bay crude oil on hatching success and associated changes in pipping muscles in embryos of domestic chickens (*Gallus gallus*). J Wildl Dis 32: 209–215.

Lustick S (1973) The effect of intense light on bird behavior and physiology. Bird Control Seminars Proceedings. p. 119.

Luttenberg D, Turgeon D, Higgins J (2001) Report from the NOAA workshops to standardize protocols for monitoring toxic Pfiesteria species and associated environmental conditions. Environ Health Perspect 109: 707-710.

Lynn JC, Rosenstock SS, Chambers CL (2008) Avian use of desert wildlife water developments as determined by remote videography. Western North Am Nat 68: 107–112.

Lyytinen A, Alatalo RV, Lindström L, Mappes J (2001) Can ultraviolet cues function as aposematic signals? Behav Ecol 12: 65–70.

Ma X (2012) Distance sampling to estimate the abundance of birds with sector and radial radar detection methods. Procedia Environ Sci 13: 2284–2303.

Mabee TJ, Cooper BA, Plissner JH, Young DP (2006) Nocturnal bird migration over an Appalachian ridge at a proposed wind power project. Wildl. Soc. Bull. 34: 682–690.

Macaulay BM, Rees D (2014) Bioremediation of oil spills: a review of challenges for research advancement. Ann Environ Sci 8: 9-37.

MacDonald GZ, Hogan NS, Köllner B, Thorpe KL, Phalen LJ, et al. (2013) Immunotoxic effects of oil sands-derived naphthenic acids to rainbow trout. Aquat Toxicol 126: 95–103.

Machtans CS, Wedeles CH, Bayne EM (2013) A first estimate for Canada of the number of birds killed by colliding with building windows. Avian Conserv Ecol 8: 6.

MacKay WC, Verbeek AG (1993) The Characterization of Acutely Toxic Compounds In Fine Tails Pore Water and Tailings Pond Water. Alberta Oil Sands Technology and Research Authority Agreement #8878M. Pp. 1-46.

MacKenzie DI, Nichols JD, Lachman GB, Droege S, Andrew Royle J, et al. (2002) Estimating site occupancy rates when detection probabilities are less than one. Ecology 83: 2248–2255.

MacKenzie DI, Nichols JD, Sutton N, Kawanishi K, Bailey LL (2005) Improving inferences in population studies of rare species that are detected imperfectly. Ecology 86: 1101–1113.

MacKinnon M, Boerger H (1986) Description of two treatment methods for detoxifying oil sands tailings pond water. Water Qual Res J Can 21: 496–512.

MacKinnon M, Matthews J, Shaw W, Cuddy R (2001) Water quality issues associated with composite tailings (CT) technology for managing oil sands tailings. Int J Surface Mini Reclamat Environ 15: 235–256.

Macko SA, King SM (1980) Weathered oil: effect on hatchability of heron and gull eggs. Bull Environ Contam Toxicol 25: 316–320.

Madill RE, Orzechowski MT, Chen G, Brownlee BG, Bunce NJ (2001) Preliminary risk assessment of the wet landscape option for reclamation of oil sands mine tailings: bioassays with mature fine tailings pore water. Environ Toxicol16: 197–208.

Madliger CL (2012) Toward improved conservation management: a consideration of sensory ecology. Biodivers Conserv 21: 3277–3286.

Mahon CL, Bayne EM, Sólymos P, Matsuoka SM, Carlson M, et al. (2014) Does expected future landscape condition support proposed population objectives for boreal birds? Forest Ecol Manag 312: 28–39.

Main C, editor (2011) 2010 Regional Aquatics Monitoring Program (RAMP) Scientific Review. Integrated Water Management Program – Alberta Innovates – Technology Futures. Pp. 1-160.

Makal S, Kizilay A, Durak L (2008) On the target classification through wavelet-compressed scattered ultrawide-band electric field data and ROC analysis. Prog Electromagn Res Pier 82: 419–431.

Mandel JT, Bohrer G, Winkler DW, Barber DR, Houston CS, et al. (2011) Migration path annotation: cross-continental study of migration-flight response to environmental conditions. Ecol Appl 21: 2258–2268.

Manel S, Williams HC, Ormerod SJ (2002) Evaluating presence-absence models in ecology: the need to account for prevalence. J Appl Ecol 38: 921–931.

Mangat R, Pinto L, Moore M (1998) Investigation of the factors that suppress bicyclic naphthenic acid degradation in Natural Wetland sediment. Suncor OSG c/o J. Gulley, Golder Associates, Calgary, AB, Pp. 1-44.

Manville A, Albert M (2005) Bird strikes and electrocutions at power lines, communication towers, and wind turbines: state of the art and state of the science-next steps toward mitigation. USDA Forest Service General Technical Report PSW-GTR-191. Pp. 1051-1064.

Marcus Rowcliffe J, Carbone C, Jansen PA, Kays R, Kranstauber B (2011) Quantifying the sensitivity of camera traps: an adapted distance sampling approach. Methods Ecol Evol 2: 464–476.

Margalida A, Ecolan S, Boudet J, Bertran J, Martinez J-M, et al. (2006) A solar-powered transmitting video camera for monitoring cliff-nesting raptors. J Field Ornithol 77: 7–12.

Margules CR, Pressey RL (2000) Systematic conservation planning. Nature 405: 243–253.

Marr JB, Mohammed P, MacKinnon M, Rodgers M (1996) Removal of naphthenic acids from Syncrude consolidated tailings using acidification and sand filtration followed by activated carbon treatment. Alberta Science and Technology Inc.

Marsh DM, Trenham PC (2008) Current trends in plant and animal population monitoring. Conserv Biol 22: 647–655.

Marsh RE, Erickson WA, Salmon TP (1991) Bird Hazing and Frightening Methods and Techniques (with emphasis on containment ponds). Other Publications in Wildlife Management.Paper 51. http://digitalcommons.unl.edu/icwdmother/51

Marsh RE, Erickson WA, Salmon TP (1992) Scarecrows and predator models for frightening birds from specific areas. Proc. 15th Vertebrate Pest Conf. 112-114.

Martin GR (2011) Through birds' eyes: insights into avian sensory ecology. J Ornithol 153(Suppl 1):S23-S48.

Martin GR (2011) Understanding bird collisions with man-made objects: a sensory ecology approach. Ibis 153: 239–254.

Martin GR, Piersma T (2009) Vision and touch in relation to foraging and predator detection: insightful contrasts between a plover and a sandpiper. Proc R Soc B Biol Sci 276: 437–445.

Martin LR, Hagar S (1990) Bird control on containment pond sites. Proceedings of the Fourteenth Vertebrate Pest Conference 1990. Paper 60. http://digitalcommons.unl.edu/vpc14/60

Martin LR, Martin LM, Taber MR (1998) Containment basins and bird exclusion-a historical perspective. Proceedings of the Eighteenth Vertebrate Pest Conference (1998). Paper 60. http://digitalcommons.unl.edu/vpc18/60

Martin LR, Martin PC (1984) Research indicates propane cannons can move birds. Pest Control 52(10):52.

Martorello DA, Eason TH, Pelton MR (2001) A sighting technique using cameras to estimate population size of black bears. Wildlife Soc B: 560–567.

Marzluff JM, Raphael MG, Sallabanks R (2000) Understanding the effects of forest management on avian species. Wildlife Soc B: 1132–1143.

Mas AH, Dietsch TV (2004) Linking shade coffee certification to biodiversity conservation: butterflies and birds in Chiapas, Mexico. Ecol Appl 14: 642–654.

Masco C (2013) Neighbor-stranger discrimination on the basis of a threat vocalization in the Great Black-Backed Gull. Wilson J Ornithol 125: 342–347.

Mason JR, Clark L (1992) Nonlethal repellents: the development of cost-effective, practical solutions to agricultural and industrial problems. Proceedings of the Fifteenth Vertebrate Pest Conference 1992.Paper 51. http://digitalcommons.unl.edu/vpc15/51

Massey JG (2006) Summary of an oiled bird response. J Exotic Pet Med 15: 33–39.

Masson JF, Leblond V, Margeson J (2006) Bitumen morphologies by phase-detection atomic force microscopy. J Microscopy 221: 17–29.

Mateos M, Arroyo GM, Rodriguez A, Cuenca D, De La Cruz A (2010) Calibration of visually estimated distances to migrating seabirds with radar measurements. J Field Ornithol 81: 302–309.

Mateos-Rodríguez M, Bruderer B (2012) Flight speeds of migrating seabirds in the Strait of Gibraltar and their relation to wind. J Ornithol153: 881–889.

Matkovich C, Quillen CJ, Stucky S, Kavanagh R, Strajt P (2010) Radar-based automated bird control system at CNRL Horizon's tailing pond – Technology overview and efficacy. In: 2010 Oil Sands & Heavy Oil Technologies Conference, Calgary.

Matsuoka SM, Bayne EM, Sólymos P, Fontaine PC, Cumming SG, et al. (2012) Using binomial distance-sampling models to estimate the effective detection radius of point-count surveys across boreal Canada. Auk 129: 268–282.

Matthews LH, Matthews BHC (1939) Owls and infra-red radiation. Nature 143(3632): 983.

Matyjasiak P (2008) Methods of bird control at airports. In: Uchmański J, editor. Theoretical and applied aspects of modern ecology. Warsaw: Cardinal Stefan Wyszyński University Press. Pp. 171-203.

Mazet J, Gardner I, Jessup D, Rittenburg J (1997) Field assay for the detection of petroleum products on wildlife. Bull Environ Contam Toxicol 59: 513–519.

Mazet JAK, Newman SH, Gilardi KVK, Tseng FS, Holcomb JB, et al. (2002) Advances in oiled bird emergency medicine and management. J Avian Med Surg 16: 146–149.

MBCA (1994) Migratory Birds Convention Act, 1994. Available: http://lawslois.justice.gc.ca/PDF/M-7.01.pdf. Accessed October 2013.

MBTA (1918) Migratory Birds Treaty Act. Available: http://www.fws.gov/laws/lawsdigest/MIGTREA.HTML. Accessed February 2014.

McComb B, Zuckerberg B, Vesely D, Jordan C (2010) Monitoring animal populations and their habitats: a practitioner's guide. CRC Press. 296 p.

McEwan E (1978) The effect of crude oils on salt gland sodium secretion of orally imposed salt loads in glaucous-winged gulls, *Larus glaucescens*. Can J Zool 56: 1212–1213.

McEwan E, Koelink A (1973) The heat production of oiled mallards and scaup. Can J Zool 51: 27–31.

McEwan E, Whitehead P (1978) Influence of weathered crude oil on liver enzyme metabolism of testosterone in gulls. Can J Zool 56: 1922–1924.

McEwan E, Whitehead P (1980) Uptake and clearance of petroleum hydrocarbons by the Glaucous-winged Gull (Larus glaucescens) and the Mallard Duck (*Anas platyrhynchos*). Can J Zool 58: 723–726.

Energy Resources Conservation Board (2009) Directive 074: Tailings Performance Criteria and Requirements for Oil Sands Mining Schemes. Calgary, AB, Canada.

McGill PA, Richmond ME (1979) Hatching success of Great Black-backed Gull eggs treated with oil. Bird-Banding 50(2): 108–113.

McKee RH, North CM, Podhasky P, Charlap JH, Kuhl A (2013) Toxicological assessment of refined naphthenic acids in a repeated dose/developmental toxicity screening Test. Int J Toxicol: 33(1 Suppl):168S-180S.

McKnight A, Sullivan K, Irons D, Stephensen S, Howlin S (2006) Marine bird and sea otter population abundance of Prince William Sound, Alaska: trends following the T/V Exxon Valdez oil spill, 1989-2005. Exxon Valdez Oil Spill Restoration Project Final Report (Restoration Projects 040159/050751), U. S. Fish and Wildlife Service, Anchorage, Alaska. 149 pp.

McLaren MA, McLaren PL (1985) Bird migration watches on Crown Lease 17, Alberta, Fall 1984. Environmental Research Monograph 1985-2. Prepared by LGL Limited for Syncrude Canada Ltd. 33 pp.

McNeill SA, Arens CJ, Hogan NS, Köllner B, van den Heuvel MR (2012) Immunological impacts of oil sands-affected waters on rainbow trout evaluated using an in situ exposure. Ecotoxicol Environ Saf 84: 254–261.

McQuillen HL, Brewer LW (2000) Methodological considerations for monitoring wild bird nests using video technology. J Field Ornithol 71: 167–172.

Mead C (1997) Poor prospects for oiled birds. Nature 390: 449–450.

Mead C, Baillie S (1981) Seabirds and oil: the worst winter. Nature 292: 10–11.

Melnikov VM, Lee RR, Langlieb NJ (2012) Resonance effects within S-band in echoes from birds. IEEE Geosci Remote Sens 9: 413–416.

Melvin SD, Lanctôt CM, Craig PM, Moon TW, Peru KM, et al. (2013) Effects of naphthenic acid exposure on development and liver metabolic processes in anuran tadpoles. Environ Pollut 177: 22–27.

Melvin SD, Trudeau VL (2012) Growth, development and incidence of deformities in amphibian larvae exposed as embryos to naphthenic acid concentrations detected in the Canadian oil sands region. Environ Pollut 167: 178–183.

Melvin SD, Trudeau VL (2012) Toxicity of naphthenic acids to wood frog tadpoles (*Lithobates sylvaticus*). J Toxicol Environ Health A 75: 170–173.

Mennill DJ (2011) Individual distinctiveness in avian vocalizations and the spatial monitoring of behaviour. Ibis 153: 235–238.

Mennill DJ, Battiston M, Wilson DR, Foote JR, Doucet SM (2012) Field test of an affordable, portable, wireless microphone array for spatial monitoring of animal ecology and behaviour. Methods Ecol Evol 3: 704–712.

Merrell R (2012) Some successful methods to mitigate conflicts caused by common ravens in an industrial environment. Human-Wildlife Interactions 6(2):339-343.

Metcalfe J, Schmidt KL, Bezner Kerr W, Guglielmo CG, MacDougall-Shackleton SA (2013) White-throated sparrows adjust behaviour in response to manipulations of barometric pressure and temperature. Anim Behav 86: 1285-1290.

Mezache A, Soltani F (2008) Threshold optimization of decentralized CFAR detection in weibull clutter using genetic algorithms. SIViP 2(1): 1–7.

Miadonye A, Evans L (2010) The solubility of asphaltenes in different hydrocarbon liquids. Petrol Sci Technol 28 (14): 1407–1414.

Miall AD (2013) Environmental Management of the Alberta Oil Sands: Introduction to the Special Set of Articles. GeoSci Can 40(3): 165–166.

Miall AD (2013) Geoscience of Climate and Energy 13. The environmental hydrogeology of the oil sands, lower Athabasca area, Alberta. GeoSci Can 40(3): 215–233.

Microbial Technologies (1995) Interactions between nitrogen- cycling bacteria and toxicants during detoxification of tailings water at the Syncrude and Suncor leases. Prepared for Syncrude Canada and Suncor Inc. 42 pp.

Microbial Technologies (1999) Hydrogen sulphide production in gypsum-amended MFT and NST. Prepared for Syncrude Research, Ltd. 39 pp.

Microbial Technologies (1999) Survey of Anaerobic Bacteria in Tailings Sampled at the Syncrude Lease. Prepared for Syncrude Research, Ltd. 25 pp.

Mignucci-Giannoni A (1999) Assessment and rehabilitation of wildlife affected by an oil spill in Puerto Rico. Environ Pollut 104: 323–333.

Mikula RJ, Kasperski KL, Burns RD, MacKinnon MD (1996) Nature and fate of oil sands fine tailings. In Schramm LL, editor. Suspensions: Fundamentals and Applications in the Petroleum Industry. ACS Advances in Chemistry Series 251: 639-675.

Miles AK, Flint PL, Trust KA, Ricca MA, Spring SE, Arrieta DE, Hollmen T, Wilson BW (2007) Polycyclic aromatic hydrocarbon exposure in Steller's eiders (Polysticta stelleri) and Harlequin ducks (Histronicus histronicus) in the eastern Aleutian Islands, Alaska, USA. Environ Toxicol Chem 26 (12): 2694–2703.

Millennium EMS Solutions Ltd. (2005) Joslyn North Mine Project Conservation & Reclamation and Closure Plan. Prepared for Deer Creek Energy Limited. File #04-050. 52 pp.

Millennium EMS Solutions Ltd. (2005) Joslyn North Mine Project Environmental Impact Assessment Hydrogeology. Prepared for Deer Creek Energy Limited. File #04-050. 88 pp.

Millennium EMS Solutions_Ltd. (2005) Soil and Terrain Report Joslyn North Mine Project. Prepared for Deer Creek Energy Limited. File #04-050. 203 pp.

Miller D, Hallett D, Peakall D (1982) Which components of crude oil are toxic to young seabirds? Environ Toxicol Chem 1: 39–44.

Miller DS, Peakall DB, Kinter WB (1978) Ingestion of crude oil: sublethal effects in herring gull chicks. Science 199: 315–317.

Miller G, Connell D (1980) Occurrence of petroleum hydrocarbons in some Australian seabirds. Wildlife Res 7: 281–293.

Millington JD, Walters MB, Matonis MS, Laurent EJ, Hall KR, et al. (2011) Combined long-term effects of variable tree regeneration and timber management on forest songbirds and timber production. Forest Ecol Manag 262: 718–729.

Milne R, Bennett L (2012) Understanding landscape patterns of temporal variability in avian populations to improve environmental impact assessments. Ecol Inform 14:75-78.

Dodds-Smith M, Filas B (1996) Preventing bird deaths. Mining Environmental Management 4(4): 10-11.

Minister of Justice (2002) Species at Risk Act. Chapter 29; http://laws-lois.justice.gc.ca/PDF/S-15.3.pdf.

Minot EO (1988) Sudden reversal of phototaxis during the development of nestling birds. Anim Behav 36: 1619–1625.

Moffitt J, Orr RT (1938) Recent disastrous effects of oil pollution on birds in the San Francisco Bay region. Pr. in California State Press Office. 6pp.

Møller A, Ibáñez-Álamo J (2012) Escape behaviour of birds provides evidence of predation being involved in urbanization. Anim Behav 84: 341-348.

Monago C, Frank O, Erhabor O (2013) Assessment of polycyclic aromatic hydrocarbon and total petroleum hydrocarbon levels in blood of refinery workers. Int J Appl Med Scis 2:18-26

Montevecchi W, Hedd A, McFarlane Tranquilla L, Fifield D, Burke C, et al. (2012) Tracking seabirds to identify ecologically important and high risk marine areas in the western North Atlantic. Biol Cons 156: 62–71.

Montevecchi WA (1999) Seabird attraction to offshore platforms and seabird monitoring from offshore support vessels and other ships: literature review and monitoring designs. Environmental Studies Research Funds. Prepared for the Canadian Association of Petroleum Producers. 51 pp.

Montevecci WA (2005) Influences of Artificial Light on Marine Birds. In: Rich C, Longcore T, editors. Ecological Consequences of Artificial Night Lighting. Washington: Island Press. pp. 94–113.

Moore BA, Baumhardt P, Doppler M, Randolet J, Blackwell BF, et al. (2012) Oblique color vision in an open-habitat bird: spectral sensitivity, photoreceptor distribution and behavioral implications. J Exp Biol 215: 3442–3452.

Moore FR, Phillips JB (1988) Sunset, skylight polarization and the migratory orientation of yellow-rumped warblers, *Dendroica coronata*. Anim Behav 36: 1770–1778.

Moore MM, Rio LD, Hadwin A, Pinto L (2002) Enhancing oil sands reclamation technologies by optimizing the microbial degradation of naphthenic acids. Prepared for Alberta Energy Research Institude, Syncrude Canada Ltd., and Suncor OSG. 166 pp.

Moore S, Dwyer R (1974) Effects of oil on marine organisms: a critical assessment of published data. Water Res 8: 819–827.

Moreno R, Jover L, Diez C, Sanpera C (2011) Seabird feathers as monitors of the levels and persistence of heavy metal pollution after the Prestige oil spill. Environ Pollut 159: 2454–2460.

Morissette JL, Kardynal KJ, Bayne EM, Hobson KA (2013) Comparing Bird Community Composition Among Boreal Wetlands: Is Wetland Classification a Missing Piece of the Habitat Puzzle? Wetlands 33: 653–665.

Morris SR, Holmes DW, Richmond ME (1996) A ten-year study of the stopover patterns of migratory passerines during fall migration on Appledore Island, Maine. Condor: 395–409.

Morton M (2011) Factors to consider in estimating oil sands plant decommissioning costs. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-16. 62 pp.

Mugerwa B, Sheil D, Ssekiranda P, Heist M, Ezuma P (2012) A camera trap assessment of terrestrial vertebrates in Bwindi Impenetrable National Park, Uganda. Afr J Ecol 51: 21-31.

Munilla I, Velando A (2010) Oiling of live gulls as a tool to monitor acute oil spill effects on seabirds. Ibis 152: 405–409.

Munro RE, Kimball CF (1982) Population ecology of the mallard. VII. Distribution and derivation of the harvest. Fish and Wildlife Service. U.S. Department of the Interior. 133 pp.

Munro U, Munro JA, Phillips JB, Wiltschko W (1997) Effect of wavelength of light and pulse magnetisation on different magnetoreception systems in a migratory bird. Aust J Zool 45: 189–198.

Munro U, Wiltschko R (1995) The role of skylight polarization in the orientation of a day-migrating bird species. J Comp Physiol A Neuroethol Sens Neural Behav Physiol 177: 357–362.

Münzberg M, Schilling A, Schlemmer H, Vogel H, Cramer H, et al. (2011) The infrared-based early warning system for bird strike prevention at Frankfurt airport. Proc of SPIE Defense, Security, and Sensing. Infrared Technology and Applications. 7 pp.

Murphy SM, Day RH, Wiens JA, Parker KR (1997) Effects of the Exxon Valdez oil spill on birds: comparisons of pre-and post-spill surveys in Prince William Sound, Alaska. Condor: 299–313.

Murphy SM, Mabee TJ (2000) Status of black oystercatchers in Prince William Sound, Alaska, nine years after the Exxon Valdez oil spill. Waterbirds: 204–213.

Nadasdy K (2014) Killing Two Birds with One Stone: How an Incidental Take Permit Program under the MBTA Can Help Companies and Migratory Birds. B.C. Envtl. Aff. L. Rev. 41: 5.

Naeth MA, Howat D, McClure H (1999) Germination of plant species on Syncrude composite tailings sands. Department of Renewable Resources, University of Alberta. Prepared for Syncrude Canada Ltd. 28 pp.

Nagai S, Saitoh TM, Noh NJ, Yoon TK, Kobayashi H, et al. (2013) Utility of information in photographs taken upwards from the floor of closed-canopy deciduous broadleaved and closed-canopy evergreen coniferous forests for continuous observation of canopy phenology. Ecol Inform 18:10-19.

Naguib M, van Oers K, Braakhuis A, Griffioen M, de Goede P, Waas JR (2013) Noise annoys: effects of noise on breeding great tits depend on personality but not on noise characteristics. Anim Behav 85(5): 949-956.

Nakhaie S (2013) Avian Species-Area Relationships and Environmental Covariates in Natural and Constructed Wetlands of Northeastern Alberta. Master of Science, University of Windsor, Windsor, Canada.

National Energy Board (2006) Canada's Oil Sands Opportunities and Challenges to 2015: An update. Calgary, Canada: Government of Canada.

Navas JM, Babin M, Casado S, Fernández C, Tarazona JV (2006) The Prestige oil spill: a laboratory study about the toxicity of the water-soluble fraction of the fuel oil. Marine Environ Res 62: S352–S355.

Navedo JG, Masero JA, Overdijk O, Orizaola G, Sánchez-Guzmán JM (2010) Assessing the role of multiple environmental factors on Eurasian Spoonbill departure decisions from stopover sites. Ardea 98: 3–12.

Nebuloni R, Capsoni C, Vigorita V (2008) Quantifying bird migration by a high-resolution weather radar. IEEE Transactions on Geoscience and Remote Sensing 46: 1867–1875.

Negro JJ, Bustamante J, Melguizo C, Ruiz JL, Grande JM (2000) Nocturnal activity of lesser kestrels under artifical lighting conditions in Seville, Spain. J Raptor Res 34(4) 327-329.

Nero V, Farwell A, Lee L, Van Meer T, MacKinnon M, et al. (2006) The effects of salinity on naphthenic acid toxicity to yellow perch: Gill and liver histopathology. Ecotoxicol Environ Saf 65: 252–264.

Nero V, Farwell A, Lister A, Van Der Kraak G, Lee LEJ, et al. (2006) Gill and liver histopathological changes in yellow perch (*Perca flavescens*) and goldfish (*Carassius auratus*) exposed to oil sands process-affected water. Ecotoxicol Environ Saf 63: 365–377.

Newman S, Anderson D, Ziccardi M, Trupkiewicz J, Tseng F, et al. (2000) An experimental soft-release of oil-spill rehabilitated American coots (*Fulica americana*): II. Effects on health and blood parameters. Environ Pollut 107: 295–304.

Newman S, Golightly R, Craig E, Carter H, Kreuder C (2004) The effects of petroleum exposure and rehabilitation on post-release survival, behavior, and blood health indices: a common murre (*Uria aalge*) case study following the Stuyvesant petroleum spill. Final Report. Oiled Wildlife Care Network, Wildlife Health Centre, School of Veterinary Medicine, University of California, Davis. 46 pp.

Newman S, Mazet J, Ziccardi M, Lieske C, Fauquier D, et al. (1999) Haematological changes and anaemia associated with captivity and petroleum exposure in seabirds. Comp Haematol Int 9: 60–67.

Newton I (2006) Can conditions experienced during migration limit the population levels of birds? J Ornithol147: 146–166.

Newton I (2008) The migration ecology of birds. London, Burlington, San Diego: Academic Press. 976 pp.

Nichols JD, Hines JE, Sauer JR, Fallon FW, Fallon JE, et al. (2000) A double-observer approach for estimating detection probability and abundance from point counts. Auk 117: 393–408.

Nichols JD, Johnson FA, Williams BK (1995) Managing North American waterfowl in the face of uncertainty. Annu Rev Ecol Evol Syst 26: 177–199.

Nichols JD, Thomas L, Conn PB (2009) Inferences about landbird abundance from count data: recent advances and future directions. In: Thomson DL, Cooch EG, Conroy MJ, editors. Modeling demographic processes in marked populations. Springer. pp. 201–235.

Nichols JD, Williams BK (2006) Monitoring for conservation. Trends Ecol Evol 21: 668–673.

Nielsen SE, Haughland DL, Bayne E, Schieck J (2009) Capacity of large-scale, long-term biodiversity monitoring programmes to detect trends in species prevalence. Biodivers Conserv 18: 2961–2978.

Nieman DJ, Dirschl HJ (1971) Waterfowl populations of the Peace-Athabasca Delta, 1969 and 1970: 30 pp.

Nix P, Hamilton S, Bauer E, Gunter C (1993) Constructed Wetlands for the Treatment of Oil Sands Waste Water Technical Report #2. Prepared for Suncor Inc. 75pp.

Nix P, Martin R (1992) Detoxification and reclamation of Suncor's oil sand tailings ponds. Environ Toxicol Water Qual 7: 171–188.

Nix PG, Gulley JR (1995) Kinetic and Empirical Design Criteria for Constructed Wetlands. Proceedings of the 19th Annual British Columbia Mine Reclamation Symposium in Dawson Creek, BC. The Technical and Research Committee on Reclamation. 99 pp.

Nix PG, Martin RW (1991) Detoxification and Reclamation of Suncor's Oil Sand Tailings Ponds. In: Proceedings of the 17th Annual Aquatic ToxicityWorkshop, Nov 5-7, 1990. Vancouver, BC. Chapman P, Bishay F, Power E, Hall K, Harding L, McLeary D, Nassichuk M. Technical Report of Fisheries and Aquatic Sciences 1774: 428-442.

Noble B, Ketilson S, Aitken A, Poelzer G (2013) Strategic environmental assessment opportunities and risks for Arctic offshore energy planning and development. Mar Policy 39: 296–302.

Nodelcorp Consulting Inc. (2005) Greenhouse Gas Management Plan, Deer Creek Energy Limited – Joslyn North Mine Project. 18 pp.

Noga J, Wolbring G (2014) The Oil and Gas Discourse from the Perspective of the Canadian and Albertan Governments, Non-Governmental Organizations and the Oil and Gas Industry. Energies 7: 314–333.

Nohara TJ, Beason RC, Weber P (2012) The role of radar-activated waterfowl deterrents on tailings ponds. International Oil Sands Tailings Conference. 9 pp.

Nohara TJ, Eng B, Eng M, Weber P, Ukrainec A, et al. (2007) An overview of avian radar developments-past, present and future. 2007 Bird Strike Committee USA/Canada, 9th Annual Meeting, Kingston, Ontario. Paper 13. http://digitalcommons.unl.edu/birdstrike2007/13

Nohara TJ, Weber P, Premji A, Krasnor C, Gauthreaux S, et al. (2005) Affordable avian radar surveillance systems for natural resource management and BASH applications. Radar Conference, 2005 IEEE International. pp. 10–15.

Northrup JM, Wittemyer G (2013) Characterising the impacts of emerging energy development on wildlife, with an eye towards mitigation. Ecol Lett 16: 112–125.

Noss RF (1990) Indicators for monitoring biodiversity: a hierarchical approach. Conserv Biol 4: 355–364.

Nwokolo E, Ohale L (1986) Growth and anatomical characteristics of pullet chicks fed diets contaminated with crude petroleum. Bull Environ Contam Toxicol 37: 441–447.

O'Brien CS, Waddell RB, Rosenstock SS, Rabe MJ (2006) Wildlife use of water catchments in southwestern Arizona. Wildlife Soc B 34: 582–591.

O'Brien TG, Kinnaird MF (2008) A picture is worth a thousand words: the application of camera trapping to the study of birds. Bird Conserv Int 18: S144.

O'Connell AF, Nichols JD, Karanth KU (2011) Camera traps in animal ecology: methods and analyses. Springer Tokyo.

O'Connell TJ, Jackson LE, Brooks RP (2000) Bird guilds as indicators of ecological condition in the central Appalachians. Ecol Appl 10: 1706–1721.

O'Hara P, Morgan K (2006) Do low rates of oiled carcass recovery in beached bird surveys indicate low rates of ship-source oil spills. Mar Ornithol 34: 133–140.

O'Hara PD, Morandin LA (2010) Effects of sheens associated with offshore oil and gas development on the feather microstructure of pelagic seabirds. Mar Pollut Bull 60: 672–678.

O'Neal BJ, Stafford JD, Larkin RP (2010) Waterfowl on weather radar: applying ground-truth to classify and quantify bird movements. J Field Ornithol 81: 71–82.

Obrist MK, Boesch R, Flückiger PF (2004) Variability in echolocation call design of 26 Swiss bat species: consequences, limits and options for automated field identification with a synergetic pattern recognition approach. Mammalia 68: 307–322.

Government of Alberta (2013) Alberta Energy: Facts and Statistics. Available: http://www.energy.alberta.ca/oilsands/791.asp. Accessed 2 April 2013.

International Union for Conservation of Nature (2001) The IUCN Red List of Threatened Species. Available: http://www.iucnredlist.org/technical-documents/categories-and-criteria. Accessed 5 February 2014.

Ogden AG (2013) Dying for a Solution: Incidental Taking Under the Migratory Bird Treaty Act. Wm & Mary Envtl L & Pol'y Rev 38: 2.

Ogochukwu ES, Okechukwu AD, Nnaegbo OG (2012) Construction And Testing Of Ultrasonic Bird Repeller. J Nat Sci Res 2: 8–17.

Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-38. 108 pp.

Oil Sands Wetlands Working Group (2000) Guideline for Wetland Establishment on Reclaimed Oil Sands Leases. Neil Chymko (Editor) Report # ESD/LM/00-1, Alberta Environment, Environmental Service. ISBN 0-7785-1042-5, Publication No. T/517.

Oka N, Okuyama M (2000) Nutritional status of dead oiled rhinoceros auklets (*Cerorhinca monocerata*) in the southern Japan sea. Mar Pollut Bull 40: 340–347.

Oka N, Takahashi A, Ishikawa K, Watanuki Y (1999) The past and present impact of oil pollution in seabird mortality world-wide. J Yamashina Inst Ornithol 31: 108–133.

Orbell JD, Tan EK, Coutts M, Bigger SW, Ngeh LN (1999) Cleansing Oiled Feathers-Magnetically. Mar Pollut Bull 38: 219–221.

Oropesa A-L, Pérez-López M, Hernández D, Garcia J-P, Fidalgo L-E, et al. (2007) Acetylcholinesterase activity in seabirds affected by the Prestige oil spill on the Galician coast (NW Spain). Sci Total Environ 372: 532–538.

Osinga S (2013) Storage of oil sand waste in sub-surface salt caverns: a feasibility analysis based on experimental rock mechanics and geological setting. Master of Science. Utrecht University, Utrecht, The Netherlands. 209 pp.

OSRIN (2010) Glossary of terms and acronyms used in oil sands mining, processing and environ manag January 2013 update. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. SR-1. 119 pp.

OSRIN (2010) OSRIN Annual Report: 2009/10. Oil Sands Research and Information Network, School of Energy and the Environment, University of Alberta, Edmonton, Alberta. OSRIN Report No. SR-3. 26 pp.

OSRIN (2011) Equivalent land capability workshop summary notes. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-13. 83 pp.

OSRIN (2012) OSRIN Annual Report: 2011/12. Oil Sands Research and Information Network, School of Energy and the Environment, University of Alberta, Edmonton, Alberta. OSRIN Report No. SR-8. 25 pp.

OSRIN (2013) OSRIN Annual Report: 2012/13. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Report Number SR-9, Edmonton, Alberta, pp. 56.

OSRIN (2014) OSRIN Annual Report: 2013/14. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Report Number SR-10, Edmonton, Alberta, pp. 66.

OSRIN and Canadian Environmental Assessment Agency, (2012) summary of the oil sands groundwater – surface water interactions workshop. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-22. 125 pp.

Overton JMC, Theo Stephens R, Leathwick JR, Lehmann A (2002) Information pyramids for informed biodiversity conservation. Biodivers Conserv 11: 2093–2116.

Overton WS, Stehman SV (1996) Desirable design characteristics for long-term monitoring of ecological variables. Environ Ecol Stat 3: 349–361.

Pacific CWS, Region Y, Hooper TD, Szaro I, Vermeer K (1987) Oil pollution of birds: an annotated bibliography. Delta, BC: Canadian Wildlife Service.

Page GW, Carter HR, Ford RG (1990) Numbers of seabirds killed or debilitated in the 1986 Apex Houston oil spill in central California. Stud Avian Biol 14: 164–174.

Palmer J, Bing K, Sharma A, Greneker E, Selee T (2011) Detecting gait alterations due to concussion impairment with radar using information-theoretic techniques. SPIE Defense, Security, and Sensing. p. 80290U–80290U.

Parajulee A, Wania F (2014) Evaluating officially reported polycyclic aromatic hydrocarbon emissions in the Athabasca oil sands region with a multimedia fate model. PNAS.

Parnell JF, Shields MA, Frierson Jr D (1984) Hatching success of Brown Pelican eggs after contamination with oil. Colon Waterbird: 22–24.

Parr SJ, Haycock RJ, Smith ME (1997) The impact of the Sea Empress oil spill on birds of the Pembrokeshire coast and islands. International Oil Spill Conference. Washington, DC: American Petroleum Institute. pp. 217–225.

Paskey J, Steward G (2012) The Alberta oil sands, journalists, and their sources. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-17. 33 pp.

Paskey JGS, Williams A (2013) The Alberta oil sands then and now: an investigation of the economic, environmental and social discourses across four decades.

Patrick SC, Weimerskirch H (2014) Personality, Foraging and Fitness Consequences in a Long

Lived Seabird. PLoS ONE 9: e87269.

Pattee OH, Franson JC (1982) Short-term effects of oil ingestion on American kestrels (*Falco sparverius*). J Wildl Dis 18: 235–241.

Patton JF, Dieter MP (1980) Effects of petroleum hydrocarbons on hepatic function in the duck. Comp Biochem Physiol C Pharmacol Comp Pharmacol 65: 33–36.

Paul Gorenzel TSCH (2012) Evaluation of a Floating Bird Diverter.

Pauly D (1995) Anecdotes and the shifting baseline syndrome of fisheries. Trends Ecol Evol 10: 430.

Peakall D, Hallett D, Bend J, Foureman G, Miller D (1982) Toxicity of Prudhoe Bay crude oil and its aromatic fractions to nestling herring gulls. Environ Res 27: 206–215.

Peakall D, Jeffrey D, Boersma D (1987) Mixed-function oxidase activity in seabirds and its relationship to oil pollution. Comp Biochem Physiol C Pharmacol Comp Pharmacol 88: 151–154.

Peakall D, Jeffrey D, Miller D (1985) Weight loss of herring gulls exposed to oil and oil emulsion. Ambio: 108–110.

Peakall D, Miller D, Kinter W (1983) Toxicity of crude oils and their fractions to nestling herring gulls—1. Physiological and biochemical effects. Mar Environ Res 8: 63–71.

Peakall D, Norstrom R, Jeffrey D, Leighton F (1989) Induction of hepatic mixed function oxidases in the herring gull (*Larus argentatus*) by prudhoe bay crude oil and its fractions. Comp Biochem Physiol C Pharmacol Comp Pharmacol 94: 461–463.

Peakall D, Tremblay J, Kinter W, Miller D (1981) Endocrine dysfunction in seabirds caused by ingested oil. Environ Res 24: 6–14.

Peakall D, Wells P, Mackay D (1987) A hazard assessment of chemically dispersed oil spills and seabirds. Mar Environ Res 22: 91–106.

Peakall DB, Hallett D, Miller D, Butler R, Kinter W (1980) Effects of ingested crude oil on black guillemots: a combined field and laboratory study. Ambio 9.

Peckford ML, Taylor PD (2008) Within night correlations between radar and ground counts of migrating songbirds. J Field Ornithol 79: 207–214.

Pedroni PM, Jaramillo H, Torres C, Navarrete ZH, Bernal-Ramirez J, et al. (2013) A partnership approach to addressing applied ecological research needs of an oil and gas business. J Appl Ecol.

Peluc SI, Sillett TS, Rotenberry JT, Ghalambor CK (2008) Adaptive phenotypic plasticity in an island songbird exposed to a novel predation risk. Behav Ecol 19: 830–835.

Pembina_Institute (2013) Understanding the impacts of oilsands production.

Pennekamp F, Schtickzelle N (2013) Implementing image analysis in laboratory-based experimental systems for ecology and evolution: a hands-on guide. Methods Ecol Evol.

Pennycuick C (1961) Sun Navigation in Birds?

Pepperberg IM, McLaughlin MA (1996) Effect of avian-human joint attention in allospecific vocal learning by grey parrots (*Psittacus erithacus*). J Comp Psychol110: 286.

Pereira AS, Islam M, Gamal El-Din M, Martin JW (2013) Ozonation degrades all detectable organic compound classes in oil sands process-affected water; an application of high-performance liquid chromatography/obitrap mass spectrometry. Rapid Communications in Mass Spectrometry 27: 2317–2326.

Pérez C, Munilla I, López-Alonso M, Velando A (2010) Sublethal effects on seabirds after the Prestige oil-spill are mirrored in sexual signals. Biol Lett 6: 33–35.

Pérez C, Velando A, Munilla I, López-Alonso M, Oro D (2008) Monitoring polycyclic aromatic hydrocarbon pollution in the marine environment after the Prestige oil spill by means of seabird blood analysis. Environ Sci Technol 42: 707–713.

Pérez-Tris J, Telleria JL (2002) Migratory and sedentary blackcaps in sympatric non-breeding grounds: implications for the evolution of avian migration. J Anim Ecol 71: 211–224.

Perkins JS (1983) Oiled Magellanic Penguins in Golfo San Jose, Argentina. Mar Pollut Bull 14: 383–387.

Perrier C, De Lope F, Møller AP, Ninni P (2002) Structural coloration and sexual selection in the barn swallow *Hirundo rustica*. Behav Ecol 13: 728–736.

Perry M, Ferrigno F, Settle F (1978) Rehabilitation of birds oiled on two mid-Atlantic estuaries. Proceedings Southeastern Association Game and Fish Commissioners 32: 318–325.

Peters LE, MacKinnon M, Van Meer T, van den Heuvel MR, Dixon DG (2007) Effects of oil sands process-affected waters and naphthenic acids on yellow perch (*Perca flavescens*) and Japanese medaka (Orizias latipes) embryonic development. Chemosphere 67: 2177–2183.

Peterson RT (1961) A field guide to western birds. Houghton Mifflin, Co., Boston.

Peterson SA (2013) Using effigies to deter American crows and common ravens in snowy plover breeding habitat.

Petit LJ, Petit DR, Martin TE (1995) Landscape-level management of migratory birds: looking past the trees to see the forest. Wildlife Soc B: 420–429.

Phillips JB, Moore FR (1992) Calibration of the sun compass by sunset polarized light patterns in a migratory bird. Behav Ecol and Sociobiology 31: 189–193.

Piatt JF, Anderson P (1996) Response of common murres to the Exxon Valdez oil spill and long-term changes in the Gulf of Alaska marine ecosystem. American Fisheries Society Symposium. Vol. 18. pp. 720–737.

Piatt JF, Carter HR, Nettleship DN (1991) Effects of oil pollution on marine bird populations. In: White, J., editor. The Effects of Oil on Wildlife. Hanover, PA: The Sheridan Press. pp. 125–141.

Piatt JF, Lensink CJ (1989) Exxon Valdez bird toll. Nature 342: 865–866.

Piatt JF, Lensink CJ, Butler W, Kendziorek M, Nysewander DR (1990) Immediate impact of the Exxon Valdez'oil spill on marine birds. Auk 107: 387–397.

Piatt JF, Van Pelt TI (1997) Mass-mortality of guillemots (*Uria aalge*) in the Gulf of Alaska in 1993. Mar Pollut Bull 34: 656–662.

Pierce AJ, Pobprasert K (2007) A portable system for continuous monitoring of bird nests using digital video recorders. J Field Ornithol 78: 322–328.

Piplani A, Gupta A, Mahani S (2007) A Low Cost, Online, Computer Controlled Robot Architecture Using a CCTV Network as Sensors. iJOE 3: 35.

Pisces_Environmental_Consulting_Services_Ltd. (2006) Fish and fish habitat baseline conditions and environmental impact assessment for the proposed Joslyn North Mine Project.

Plank MJ, Codling EA (2009) Sampling rate and misidentification of Lévy and non-Lévy movement paths. Ecology 90: 3546–3553.

Plant AL, Crowe A, Ban H, Kermode A, Tang J, Bendell-Young L, Pollet I, Tang J, Kolok A, Farrell AP, Bennett K, Challanger W, Williams T (1998) Ecological characteristics of wetlands receiving oil sands effluent.

Plant AL, Crowe A, Ban H, Kermode A, Tang J, et al. (1998) Ecological characteristics of wetlands receiving oil sands effluent.

Plonczkier P, Simms IC (2012) Radar monitoring of migrating pink-footed geese: behavioural responses to offshore wind farm development. J Appl Ecol 49: 1187–1194.

Pohl H (1999) Spectral composition of light as a zeitgeber for birds living in the high Arctic summer. Physiol Behav 67: 327–337.

Pollard JB, Gendron M, Smyth SA, Richard AJ, Stewart GR (2000) Peace- Athabasca Delta Waterbird Inventory: 1999 Surveys Final Report.

Pollet I, Bendell-Young LI (2000) Amphibians as indicators of wetland quality in wetlands formed from oil sands effluent. Environ Toxicol Chem 19: 2589–2597.

Poon W, Wong C, Abdullah K, Lim E, Teo C (2011) Development of Migratory Birds Population Monitoring System Using Digital Single Reflex Camera. Computer Graphics, Imaging and Visualization (CGIV), 2011 Eighth International Conference on. pp. 136–140.

Poot H, Ens BJ, de Vries H, Donners MAH, Wernand MR, et al. (2008) Green light for nocturnally migrating birds. Ecol Soc 13: 47.

Pounder B (1974) Wildfowl and pollution in the Tay estuary. Mar Pollut Bull 5: 35–38.

Pouwels R, Opdam P, Jochem R (2011) Reconsidering the Effectiveness of Scientific Tools for Negotiating Local Solutions to Conflicts between Recreation and Conservation with Stakeholders. Ecol Soc 16.

Prach R, Surrendi D (1978) Migratory bird management and industrial cooling in northern climates. Can Water Resour J 3: 111–120.

Prichard AK, Roby DD, Bowyer RT, Duffy LK (1997) Pigeon guillemots as a sentinel species: A dose-response experiment with weathered oil in the field. Chemosphere 35: 1531–1548.

Proppe DS, Sturdy CB, St. Clair CC (2011) Flexibility in animal signals facilitates adaptation to rapidly changing environments. PLoS ONE 6: e25413.

Protection at the Speed of Sound (2011) \$1.1 Million HyperSpike Order for Bird Deterrence Announced by Ultra Electronics USSI. Available: http://www.ultra-ussi.com/latest-news/95-11-million-hyperspike-order-for-bird-deterrence-announced-by-ultra-electronics-ussi. Accessed June 2012.

Puttaswamy N, Turcotte D, Liber K (2010) Variation in toxicity response of Ceriodaphnia dubia to Athabasca oil sands coke leachates. Chemosphere 80: 489–497.

Pyle P (2007) Standardizing at-sea monitoring programs for marine birds, mammals, other organisms, debris, and vessels, including recommendations for west-coast national marine

sanctuaries. Prepared as part of the integration of a multi-sanctuary ecosystem observation effort: Gulf of the Farallones, Cordell Bank, and Monterey Bay National Marine Sanctuaries Point Reyes Statsion, CA: The Institute for Bird Populations.

Pöysä H (1996) Population estimates and the timing of waterfowl censuses. Ornis Fennica 73: 60–68.

R. v. Syncrude Canada Ltd. (2010). ABPC 229. Available: http://www.canlii.org/en/ab/abpc/doc/2010/2010abpc229/2010abpc229.pdf.

Radvanyi A (1980) Control of small mammal damage in the Alberta oil sands reclamation and afforestation program. Forest Sci 26: 687–702.

Rajabi H, Monsef H, Moghadami M, Zare M, Armandei A (2014) Design, development and demonstration of an improved bird washing machine. Ecotoxicology: 1–6.

Rajchard J (2008) Exogenous chemical substances in bird perception: a review. Veterinarni Medicina 53: 412–419.

Ramirez Jr P (2002) Oil field produced water discharges into wetlands in Wyoming.

Ramirez Jr P (2005) Oil field-produced water discharges into wetlands: Benefits and risks to wildlife. Environ Geosciences 12: 65–72.

Ramirez Jr P (2010) Bird mortality in oil field wastewater disposal facilities. Environ Manag 46: 820–826.

Ramsar (2001) Ramsar Directory of Wetlands of International Importance. Canada 7: Peace Athabasca Delta Information Sheet. Available:

http://ramsar.wetlands.org/Database/Searchforsites/tabid/765/language/en-US/Default.aspx. Accessed 24 October 2009.

Randall LA, Diehl RH, Wilson BC, Barrow WC, Jeske CW (2011) Potential use of weather radar to study movements of wintering waterfowl. The J Widlife Manag 75: 1324–1329.

Randall R, Randall B, Bevan J (1980) Oil pollution and penguins—Is cleaning justified? Mar Pollut Bull 11: 234–237.

Randler C (2004) Frequency of bird hybrids: does detectability make all the difference? J Ornithol 145: 123–128.

Randolet J, Lucas JR, Fernández-Juricic E (2014) Non-Redundant Social Information Use in Avian Flocks with Multisensory Stimuli. Ethology.

Rappl R, Wiltschko R, Weindler P, Berthold P, Wiltschko W (2000) Orientation behavior of Garden Warblers (*Sylvia borin*) under monochromatic light of various wavelengths. Auk 117: 256–260.

Rappole JH, Warner DW (1976) Relationships between behavior, physiology and weather in avian transients at a migration stopover site. Oecologia 26: 193–212.

Rattner BA (1981) Tolerance of adult mallards to subacute ingestion of crude petroleum oil. Toxicol Lett 8: 337–342.

Rattner BA (2009) History of wildlife toxicology. Ecotoxicology 18: 773–783.

Rattner BA, Eastin WC (1981) Plasma corticosterone and thyroxine concentrations during chronic ingestion of crude oil in mallard ducks (*Anas platyrhynchos*). Comp Biochem Physiol C Pharmacol Comp Pharmacol 68: 103–107.

Rattner BA, Eroschenko VP, Fox GA, Fry DM, Gorsline J (1984) Avian endocrine responses to environmental pollutants. J Exp Zool 232: 683–689.

Ravenscroft N, Beardall C (2003) The importance of freshwater flows over estuarine mudflats for wintering waders and wildfowl. Biol Cons 113: 89–97.

Read J (2001) A strategy for minimizing waterfowl deaths on toxic waterbodies. J Appl Ecol 36: 345–350.

Read J, Whiteoak D (2003) The Shell bitumen handbook. Thomas Telford Services Limited.

Rearden JD (1951) Identification of waterfowl nest predators. The J Widlife Manag: 386–395.

Reid LM (2007) The epidemiology of monitoring. AWRA 37: 815–820.

Reilly TJ (1995) Bird Deterrent Research and Development: Marine Oil Spills.

Rempel RS, Andison DW, Hannon SJ (2004) Guiding principles for developing an indicator and monitoring framework. Forest Chron80: 82–90.

Rempel RS, Francis CM, Robinson JN, Campbell M (2013) Comparison of audio recording system performance for detecting and monitoring songbirds. J Field Ornithol 84: 86–97.

Renault S (2001) Early Revegetation Strategies on CT and CT-amended Substrates.

Renault S, Zwiazek JJ (1996) Phytotoxicity of reclaimed fine tails and tailings sands.

Renault S, Zwiazek JJ (1997) Effects of CT Materials on Plants - Research Project 1997.

Renault S, Zwiazek JJ (1997) Salt tolerance of plants treated with consolidated tailings water.

Renault S, Zwiazek JJ, Croser C, Franklin J, Redfield E (1998) Effects of CT Materials on Plants - Progress Report 1998.

Rensel LJ, Wilder JD (2012) The Effects of Owl Decoys and Non-threatening Objects on Bird Feeding Behavior. Quercus: Linfield Journal of Undergraduate Research 1: 4.

Reside AE, VanDerWal JJ, Kutt AS, Perkins GC (2010) Weather, not climate, defines distributions of vagile bird species. PLoS ONE 5: e13569.

Response Corporation Ltd. E (n.d.) Bird hazing devices.

Rhodes OE, DeVault TL, Smith LM (2006) Seasonal variation in carcass composition of American Wigeon wintering in the Southern High Plains. J Field Ornithol 77: 220–228.

Ribic CA, Thompson III FR, Pietz PJ (2012) Video Surveillance of Nesting Birds. University of California Pr.

Richardson F (1956) Sea birds affected by oil from the freighter Seagate. The Murrelet: 20–22.

Richardson MG, Heubeck M, Lea D, Reynolds P (1982) Oil pollution, seabirds, and operational consequences, around the Northern Isles of Scotland. Environ Conserv 9: 315–321.

RICHARDSON T, Gardali T, Jenkins SH (2009) Review and meta-analysis of camera effects on avian nest success. The J Widlife Manag 73: 287–293.

Richardson WJ (1978) Timing and amount of bird migration in relation to weather: a review. Oikos: 224–272.

Richardson WJ (1991) Wind and orientation of migrating birds: a review. Orientation in birds. Springer. pp. 226–249.

Richardson WJ, Gunn WWH (1971) Radar observations of bird movements in east-central Alberta. Studies of Bird Hazards to Aircraft : 35–68.

Richman SE (n.d.) Sea duck predation on mussel farms: a growing conflict.

Riffaut L, McCoy KD, Tirard C, Friesen VL, Boulinier T (2005) Population genetics of the common guillemot *Uria aalge* in the North Atlantic: geographic impact of oil spills. Mar Ecol Prog Ser 291: 263–273.

Ringelman KM (2014) Predator foraging behavior and patterns of avian nest success: What can we learn from an agent-based model? Ecol Modell 272: 141–149.

Ringelman KM, Eadie JM, Ackerman JT (2012) Density-dependent nest predation in waterfowl: the relative importance of nest density versus nest dispersion. Oecologia 169: 695–702.

Rioux S, Savard J-PL, Gerick AA (2013) Avian mortalities due to transmission line collisions: a review of current estimates and field methods with an emphasis on applications to the Canadian electric network. Avian Conserv Ecol8.

Robbins CS, Brunn B, Zim HS (1966) A guide to field- identification: birds of North America. Western Publishing Co., Inc, Racine, Wisconsin.

Robinson WD, Bowlin MS, Bisson I, Shamoun-Baranes J, Thorup K, et al. (2009) Integrating concepts and technologies to advance the study of bird migration. Front Ecol Environ 8: 354–361.

Robson D, Barriocanal C (2011) Ecological conditions in wintering and passage areas as determinants of timing of spring migration in trans-Saharan migratory birds. J Anim Ecol 80: 320–331.

Rock P, Vaughan IP (2013) Long-term estimates of adult survival rates of urban Herring Gulls *Larus argentatus* and Lesser Black-backed Gulls *Larus fuscus*. Ringing & Migration 28: 21–29.

Rocke T, Yuill T, Hinsdill R (1984) Oil and related toxicant effects on mallard immune defenses. Enviro Res 33: 343–352.

Rodr'\iguez A, Rodr'\iguez B, Curbelo A, Pérez A, Marrero S, et al. (2012) Factors affecting mortality of shearwaters stranded by light pollution. Anim Conserv 15: 519–526.

Rodriguez A, Negro JJ, Bustamante J, Antolin J (2013) Establishing a Lesser Kestrel colony in an urban environment for research purposes. J Raptor Res 47: 214–218.

Rodriguez A, Negro JJ, Mulero M, Rodriguez C, Hernández-Pliego J, et al. (2012) The Eye in the Sky: Combined Use of Unmanned Aerial Systems and GPS Data Loggers for Ecological Research and Conservation of Small Birds. PLoS ONE 7: e50336.

Rogers AS, DeStefano S, Ingraldi M (2005) Quantifying Northern Goshawk diets using remote cameras and observations from blinds. J Raptor Res 39: 303.

Rogers V (2003) Mammalian Toxicity of Napthenic Acids Derived From the Athabasca Oil Sands.

Rogers VV, Liber K, MacKinnon MD (2002) Isolation and characterization of naphthenic acids from Athabasca oil sands tailings pond water. Chemosphere 48: 519–527.

Rogers VV, Wickstrom M, Liber K, MacKinnon MD (2002) Acute and subchronic mammalian toxicity of naphthenic acids from oil sands tailings. Toxicol Sci 66: 347–355.

Romero L, Wikelski M (2002) Severe effects of low-level oil contamination on wildlife predicted by the corticosterone-stress response: preliminary data and a research agenda. Spill Sci Technol Bull 7: 309–313.

Ronald KL, Fernández-Juricic E, Lucas JR (2012) Taking the sensory approach: how individual differences in sensory perception can influence mate choice. Anim Behav.

Ronchi-Virgolini AL, Lorenzon RE, Blake JG, Beltzer AH (2013) Temporal variation of bird assemblages in a wetland: influence of spatial heterogeneity. Avian Biol Res 6: 198–206.

Ronconi R (2011) Oil Sands Bird Contact Monitoring Proposed Methodology.

Ronconi RA (2006) Predicting bird oiling events at oil sands tailings ponds and assessing the importance of alternate waterbodies for waterfowl: a preliminary assessment. Can Field Nat 120: 1–9.

Ronconi RA (2011) Oil Sands Bird Contact Monitoring Plan for 2011.

Ronconi RA, St. Clair CC (2006) Efficacy of a radar-activated on-demand system for deterring waterfowl from oil sands tailings ponds. J Appl Ecol 43: 111–119.

Ronconi RA, St. Clair CC, O'Hara PD, Burger AE (2004) Waterbird deterrence at oil spills and other hazardous sites: potential applications of a radar-activated on-demand deterrence system. Mar Ornithol 32: 25–33.

Rooney RC, Bayley SE, Schindler DW (2012) Oil sands mining and reclamation cause massive loss of peatland and stored carbon. PNAS 109: 4933–4937.

Rosa Salva O, Regolin L, Vallortigara G (2007) Chicks discriminate human gaze with their right hemisphere. Behav Brain Res 177: 15–21.

Ross B, Lien J, Furness R (2001) Use of underwater playback to reduce the impact of eiders on mussel farms. ICES J Mar Sci 58: 517–524.

Rowan MK (1978) Scarecrows to save birds.

Rowcliffe J, Carbone C (2008) Surveys using camera traps: are we looking to a brighter future? Anim Conserv 11: 185–186.

Rowland SJ, Scarlett AG, Jones D, West CE, Frank RA (2011) Diamonds in the rough: identification of individual naphthenic acids in oil sands process water. Environ Sci Technol 45: 3154–3159.

Roy M-C, Mollard FP, Foote AL (2014) Do peat amendments to oil sands wet sediments affect Carex aquatilis biomass for reclamation success? J Environ Manag 139: 154–163.

Rozhok A (2008) Orientation and navigation in vertebrates. Springer.

Rubinstein I, Strausz O, Spyckerelle C, Crawford R, Westlake D (1977) The origin of the oil sand bitumens of Alberta: a chemical and a microbiological simulation study. Geochimica et Cosmochimica Acta 41: 1341–1353.

Rudd MA (2011) Scientists' opinions on the global status and management of biological diversity. Conservation Biology 25: 1165–1175.

Ruiz-Gutiérrez V, Zipkin EF (2011) Detection biases yield misleading patterns of species persistence and colonization in fragmented landscapes. Ecosphere 2.

Ruiz-Rodríguez M, Avilés J, Cuervo J, Parejo D, Ruano F, et al. (2013) Does avian conspicuous colouration increase or reduce predation risk? Oecologia: 1–11.

Rullan CG (2010)

Responses of Gulls to a Remote Control Operated Hawk Model: Implications for Bird Strikes.

Russell Mason J, Clark L (1995) Avian Repellents: Options, Modes of Action, and Economic Considerations.

Ruth JM, Barrow WC, Sojda RS, Dawson DK, Diehl RH, et al. (2005) Advancing migratory bird conservation and management by using radar: an interagency collaboration. US Geological Survey.

Ruth JM, Buler JJ, Diehl RH, Sojda RS (2008) Management and research applications of long-range surveillance radar data for birds, bats, and flying insects. US Department of the Interior, US Geological Survey.

Ruth JM, Diehl RH, Felix Jr RK (2012) Migrating Birds' use of Stopover Habitat in The Southwestern United States. Condor 114: 698–710.

Ruth, JM (2007) Applying Radar Technology to Migratory Bird Conservation and Management: Strengthening and Expanding a Collaborative. Reston, Virginia: U.S. Geological Survey.

Ryan PG (2003) Estimating the demographic benefits of rehabilitating oiled African penguins. Rehabilitation of Oiled African Penguins: a Conservation Success story Birdlife South Africa and the Avian Demography Unit, Cape Town.

Saino N, Rubolini D, Von Hardenberg J, Ambrosini R, Provenzale A, et al. (2010) Spring migration decisions in relation to weather are predicted by wing morphology among trans-Mediterranean migratory birds. Funct Ecol 24: 658–669.

Sallabanks R, Arnett EB, Marzluff JM (2000) An evaluation of research on the effects of timber harvest on bird populations. Wildlife Soc B: 1144–1155.

Salt WR, Salt JR (1976) The Birds of Alberta. Hurtig Publishers, Edmonton.

Samuels WB, Ladino A (1984) Calculations of seabird population recovery from potential oilspills in the mid-Atlantic region of the United States. Ecol Modell 21: 63–84.

Sandel B, Smith AB (2009) Scale as a lurking factor: incorporating scale-dependence in experimental ecology. Oikos 118: 1284–1291.

Sander B, Nix P, Gunter C (1995) Biological Treatment Options for Consolidated Tailings Release Waters.

Sands JP, Pope MD (2010) A survey of galliform monitoring programs and methods in the United States and Canada. Widlife Biol 16: 342–356.

Santos CD, Lourenço PM, Miranda AC, Granadeiro JP, Palmeirim JM (2008) Birds after dark: an efficient and inexpensive system for making long-range observations at night. J Field Ornithol 79: 329–335.

Santos CD, Miranda AC, Granadeiro JP, Lourenço PM, Saraiva S, et al. (2010) Effects of artificial illumination on the nocturnal foraging of waders. Acta oecologica 36: 166–172.

Santos H, Rodrigues L, Jones G, Rebelo H (2013) Using species distribution modelling to predict bat fatality risk at wind farms. Biol Cons 157: 178–186.

Saracco JF, Desante DF, Kaschube DR (2008) Assessing landbird monitoring programs and demographic causes of population trends. The J Widlife Manag 72: 1665–1673.

Saunders W (1930) The destruction of birds at Long Point Light-House, Ontario, on four nights in 1929. Auk: 507–511.

Sauter A, Korner-Nievergelt F, Jenni L (2010) Evidence of climate change effects on within-winter movements of European Mallards *Anas platyrhynchos*. Ibis 152: 600–609.

Scaife M (1976) The response to eye-like shapes by birds II. The importance of staring, pairedness and shape. Anim Behav 24: 200–206.

Scaife M (1976) The response to eye-like shapes by birds. I. The effect of context: a predator and a strange bird. Anim Behav 24: 195–199.

Scanes CG, McNabb F (2003) Avian models for research in toxicology and endocrine disruption. Avian Poultry Biol Rev 14: 21–52.

Scaring Birds (n.d.) Scarecrow: wind powered constantly revolving visual deterrent. Available: http://www.scaringbirds.com/assets/productinfo/scarecrow.pdf. Accessed 10 February 2014.

Scarlett AG, West CE, Jones D, Galloway TS, Rowland SJ (2012) Predicted toxicity of naphthenic acids present in oil sands process-affected waters to a range of environmental and human endpoints. Sci Total Environ 425: 119–127.

Schaub M (2012) Spatial distribution of wind turbines is crucial for the survival of red kite populations. Biol Cons 155: 111–118.

Schaub M, Kery M, Korner P, Korner-Nievergelt F (2011) A critique of "Collision mortality has no discernable effect on population trends of North American Birds." PLoS ONE 6.

Schick CD, Ambrock DR (1974) Waterfowl investigations in the Athabasca tar sands area. Canadian Wildlife Service: Canada.

Schieck J, Song SJ (2006) Changes in bird communities throughout succession following fire and harvest in boreal forests of western North America: literature review and meta-analyses. Can J Forest Res 36: 1299–1318.

Schley P, Pinto L, Moore M (1998) Biodegradation of Naphthenic Acids in Sediments Receiving Oil Sands Wastewater.

Schlierf R, Hight R, Payne S, Shaffer J, Missimer B, et al. (n.d.) KENNEDY SPACE CENTER (KSC) LAUNCH PAD AVIAN ABATEMENT EFFORTS INCLUDING RELATED KSC ROAD KILL REDUCTION EFFORT.

Schmaljohann H, Liechti F, Bächler E, Steuri T, Bruderer B (2008) Quantification of bird migration by radar-a detection probability problem. Ibis 150: 342–355.

Schmaljohann H, Liechti F, Bruderer B (2009) Trans-Sahara migrants select flight altitudes to minimize energy costs rather than water loss. Behav Ecol and Sociobiology 63: 1609–1619.

Schmidt KA, Dall SRX, Van Gils JA (2009) The ecology of information: an overview on the ecological significance of making informed decisions. Oikos 119: 304–316.

Schneider DC (2001) The rise of the concept of scale in ecology. Biosci 51: 545–553.

Schneider R, Dyer S, Parks C (2006) Death by a thousand cuts: impacts of in situ oil sands development on Alberta's boreal forest. Pembina Institute.

Schneider RR, Hauer G, Farr D, Adamowicz W, Boutin S (2011) Achieving conservation when opportunity costs are high: optimizing reserve design in Alberta's oil sands region. PLoS ONE 6: e23254.

Schneider RR, Stelfox JB, Boutin S, Wasel S (2003) Managing the cumulative impacts of landuses in the western Canadian sedimentary basin: a modeling approach. Conserv Ecol 7: 8.

Schneider T (2011) Accounting for environmental liabilities under international financial reporting standards. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-9. 16 pp.

Scott AC (2007) Biodegradation, Ozonation, and Characterization of Naphthenic Acids.

Scott AC, MacKinnon MD, Fedorak PM (2005) Naphthenic acids in Athabasca oil sands tailings waters are less biodegradable than commercial naphthenic acids. Environ Sci Technol 39: 8388–8394.

Scott JM, Ramsey FL (1981) Effects of abundant species on the ability of observers to make accurate counts of birds. Auk 98: 610–613.

Seamans TW (2004) Response of roosting turkey vultures to a vulture effigy. USDA National Wildlife Res Center-Staff Publications: 385.

Seamans TW, Barras SC, Patton ZJ (2003) Are birds scared by rotating mirrors? 2003 Bird Strike Committee USA/Canada, 5th Joint Annual Meeting, Toronto, ONT. p. 12.

Seamans TW, Bernhardt GE (2004) Response of Canada geese to a dead goose effigy. USDA National Wildlife Research Center - Staff Publications. Paper 384.

Seamans TW, Bernhardt GE (2004) Response of Canada geese to a dead goose effigy. USDA National Wildlife Research Center - Staff Publications. Paper 384.

Seamans TW, Blackwell BF (2011) Electric shock strips as bird deterrents: does experience count? Int J Pest Manag 57: 357–362.

Seamans TW, Blackwell BF, Gansowski JT (2002) Evaluation of the Allsopp Helikite as a bird scaring device. Proceedings of the Vertebrate Pest Conference. Vol. 20. pp. 129–134.

Seamans TW, Clemons SE, Gosser AL (2009) Observations of neck-collared Canada geese near John F. Kennedy International Airport, New York.

Seamans TW, Helon DA (2006) Evaluation of the Chroma-Flair\textregistered Crow Buster as a Starling Repellent at Nest Sites. Proceedings-Vertebrate Pest Conference. p. 228.

Seamans TW, Lovell CD, Dolbeer RA, Cepek JD (2001) Evaluation of mirrors to deter nesting starlings. Wildlife Soc B: 1061–1066.

Seip KL, Sandersen E, Mehlum F, Ryssdal J (1991) Damages to seabirds from oil spills: comparing simulation results and vulnerability indexes. Ecol Modell 53: 39–59.

Seiser PE, Duffy LK, David McGuire A, Roby DD, Golet GH, et al. (2000) Comparison of pigeon guillemot, *Cepphus columba*, blood parameters from oiled and unoiled areas of Alaska eight years after the Exxon Valdez oil spill. Mar Pollut Bull 40: 152–164.

Sellick A, Boyd M, Singh K (2006) Applications for the Deer Creek Energy Limited ("DCEL") Joslyn North Mine Project (the "Project").

Selman W, Hess Jr TJ, Salyers B, Salyers C (2012) Short-Term Response of Brown Pelicans (Pelecanus occidentalis) to Oil Spill Rehabilitation and Translocation. Southeastern Naturalist 11: 1–16.

Semenchuk, GP (1992) Atlas of the breeding birds of Alberta. Nature Alberta. 391 p.

Semm P, Beason R (1990) Sensory basis of bird orientation. Cell Molec Life Scie 46: 372–378.

Serafin J (1983) Nutrition and disease relationships that may serve as models for feeding oiled birds. US Fish and Wildlife Service.

Seymour RJ, Geyer RA (1992) Fates and effects of oil spills. Annual Review of Energy and the Environment 17: 261–283.

Seys J, Offringa H, Meire P, Van Waeyenberge J, Kuijken E (2002) Long-term changes in oil pollution off the Belgian coast: evidence from beached bird monitoring.

Shakeri M, Zhang H (2012) Real-time bird detection based on background subtraction. Proceedings of the 10th World Congress on Intelligent Control and Automation.

Shamoun-Baranes J, Van Loon E, Alon D, Alpert P, Yom-Tov Y, et al. (2006) Is there a connection between weather at departure sites, onset of migration and timing of soaring-bird autumn migration in Israel? Global Ecology and Biogeography 15: 541–552.

Sharp BE (1996) Post-release survival of oiled, cleaned seabirds in North America. Ibis 138: 222–228.

Sharp L (1987) Chasing birds from oil spills—two experiments. Coastal Zone'87. pp. 1–15.

Sharp PL, Birdsall DA, Richardson WJ (1975) Inventory studies of birds on and near crown lease number 17, Athabasca Tar Sands, 1974. Syncrude Canada Limited.

Shaw RD, Hamilton HR, MacKinnon MD, Gulley J (1995) A Modelling Framework for Reclamation Planning of Oil Sands Mines in Northern Alberta.

Shell (2012) Overview of tailings and Shell technologies.

Sherman DE, Barras AE (2004) Efficacy of a laser device for hazing Canada geese from urban areas of northeast Ohio. Ohio J Sci 104: 38–42.

Sherrington M (2005) Biodiversity assessment in the oil sands region, northeastern Alberta, Canada. Impact Assessment and Project Appraisal 23: 73–81.

Shirose LJ, Bishop CA, Green DM, MacDonald CJ, Brooks RJ, et al. (1997) Validation tests of an amphibian call count survey technique in Ontario, Canada. Herpetologica: 312–320.

Short J, Kelley M, SPEELMAN III R, McCarty R (2000) Birdstrike Prevention: Applying Aeroscience and Bio-Science. International Bird Strike Committee IBSC25/WP-RS4 463: 485.

Shortt R (1985) Bird use of alternate waterbodies in the Syncrude Canada Ltd. area.

Shrout PE, Fleiss JL (1979) Intraclass correlations: uses in assessing rater reliability. Psychological bulletin 86: 420.

Siemers BM, Schaub A (2011) Hunting at the highway: traffic noise reduces foraging efficiency in acoustic predators. Proc R Soc B Biol Sci 278: 1646–1652.

Sih A (1992) Prey uncertainty and the balancing of antipredator and feeding needs. Am Nat: 1052–1069.

Sih A (2013) Understanding variation in behavioural responses to human-induced rapid environmental change: a conceptual overview. Anim Behav.

Sih A, Bell AM, Johnson JC, Ziemba RE (2004) Behavioral syndromes: an integrative overview. The Quarterly Review of Biology 79: 241–277.

Sih A, Ferrari MC, Harris DJ (2011) Evolution and behavioural responses to human-induced rapid environmental change. Evolutionary Applications 4: 367–387.

Silvy NJ, Lopez RR, Peterson MJ (2005) Wildlife marking techniques. Techniques for wildlife investigations and management Bethesda: The Wildlife Society: 339–363.

Simons TR, Moore FR, Gauthreaux Jr S (2004) Mist netting trans-Gulf migrants at coastal stopover sites: the influence of spatial and temporal variability on capture data. Stud Avian Biol 29: 135–143.

Singh R, Dev K, Kour DN, Sahi D (2014) Bird communities structure along with species diversity, relative abundance and habitat use of Tehsil Udhampur, Jammu and Kashmir, India. Int J Pure Appl Zool 2: 26–40.

Sirot E (2006) Social information, antipredatory vigilance and flight in bird flocks. Anim Behav 72: 373–382.

Siwik PL, Van Meer T, MacKinnon MD, Paszkowski CA (2000) Growth of fathead minnows in oilsand-processed wastewater in laboratory and field. Environ Toxicol Chem 19: 1837–1845.

Slabbekoorn H (2013) Songs of the city: noise-dependent spectral plasticity in the acoustic phenotype of urban birds. Anim Behav.

Smart S, Henrys P, Purse B, Murphy J, Bailey M, et al. (2012) Clarity or confusion?-Problems in attributing large-scale ecological changes to anthropogenic drivers. Ecol Indic 20: 51–56.

Smith GE, Vespe M, Woodbridge K, Baker CJ (2008) Radar classification evaluation. Radar Conference, 2008. RADAR'08. IEEE. pp. 1–6.

Smith LM, Haukos DA, Prather RM (2004) Avian response to vegetative pattern in playa wetlands during winter. Wildlife Soc B 32: 474–480.

Smith REH, Almond MJR, McIntyre S, Taylor WD, Dixon DG (1997) Analysis and assessment of plankton data collected from experientnal ponds on the Syncrude Canda Ltd. lease (1993, 1994, 1995).

Smits J, Bortolotti GR, Tella JL (1999) Simplifying the phytohaemagglutinin skin-testing technique in studies of avian immunocompetence. Funct Ecol 13: 567–572.

Smits J, Miller M, Zimmer C, Wayland M (1998) Impacts of Wetland Contaminants on Tree Swallows and Mallard Ducks - Year 2 Annual Report. Report.

Smits JE, Fernie KJ (2013) Avian wildlife as sentinels of ecosystem health. Comp Immunol Microbiol Infect Dis 36: 333–342.

Smits JE, Wayland ME, Miller MJ, Liber K, Trudeau S (2000) Reproductive, immune, and physiological end points in tree swallows on reclaimed oil sands mine sites. Environ Toxicol Chem 19: 2951–2960.

Smits JE, Williams TD (1999) Validation of immunotoxicology techniques in passerine chicks exposed to oil sands tailings water. Ecotoxicol Environ Saf 44: 105–112.

Smits JEG, Hersikorn BD, Young RF, Fedorak PM (2012) Physiological effects and tissue residues from exposure of leopard frogs to commercial naphthenic acids. Sci Total Environ 437: 36–41.

Smits JEG, Liber K, Zimmer C, Wayland M (1997) Impacts of Wetland Contaminants on Tree Swallows and Mallard Ducks - Year 1 Annual Report. Report.

Snäll T, Kindvall O, Nilsson J, Pärt T (2011) Evaluating citizen-based presence data for bird monitoring. Biol Cons 144: 804–810.

Soanes L, Atkinson P, Gauvain R, Green J (2012) Individual consistency in the foraging behaviour of Northern Gannets: Implications for interactions with offshore renewable energy developments. Marine Policy.

Sohrabi V, Ross M, Martin J, Barker J (2013) Potential for in situ chemical oxidation of acid extractable organics in oil sands process affected groundwater. Chemosphere.

Sojda RS, Ruth JM, Barrow WC, Dawson DK, Diehl RH, et al. (2005) Using Radar to Advance Migratory Bird Management: an Interagency Collaboration. Fort Collins Science Center: US Geological Survey.

Sol D, Lapiedra O, González-Lagos C (2013) Behavioural adjustments for a life in the city. Anim Behav.

Soldatini C, Albores-Barajas YV, Torricelli P, Mainardi D (2008) Testing the efficacy of deterring systems in two gull species. Applied Anim Behav Science 110: 330–340.

Sólymos P, Lele S, Bayne E (2012) Conditional likelihood approach for analyzing single visit abundance survey data in the presence of zero inflation and detection error. Environmetrics 23: 197–205.

Sólymos P, Matsuoka SM, Bayne EM, Lele SR, Fontaine P, et al. (2013) Calibrating indices of avian density from non-standardized survey data: making the most of a messy situation. Methods Ecol Evol 4: 1047–1058.

Somers CM, Morris RD (2002) Birds and wine grapes: foraging activity causes small-scale damage patterns in single vineyards. J Appl Ecol 39: 511–523.

Song D, Xu Y (2010) A low false negative filter for detecting rare bird species from short video segments using a probable observation data set-based EKF method. Image Processing, IEEE Transactions on 19: 2321–2331.

Song D, Xu Y (n.d.) Monocular Vision-based Detection of a Flying Bird.

Soper, JD (1951) Waterfowl and related investigations in the Peace-Athabasca Delta region of Alberta. Canadian Wildlife Service: Wildlife Management, Ottawa, Ontario, Canada.

Soudat-Soucaze J, Ferri M (1997) A means of scaring birds: the laser gun, description and applications to cormorants and other birds. Desman SAR L, France in cooperation with the Office for Wildlife Protection and Regulation of Hunting and Fishing, Modena Province Regione Emilia Romagna, Italy.

Sovacool BK (2009) Contextualizing avian mortality: A preliminary appraisal of bird and bat fatalities from wind, fossil-fuel, and nuclear electricity. Energy Policy 37: 2241–2248.

Spaling H, Zwier J, Ross W, Creasey R (2000) Managing regional cumulative effects of oil sands development in Alberta, Canada. J Environ Assess Policy Manag 2: 501–528.

Spanier E (1980) The use of distress calls to repel night herons (*Nycticorax nycticorax*) from fish ponds. J Appl Ecol: 287–294.

Speakman J, Ward S (1998) Infrared thermography: principles and applications. ZOOLOGY-JENA- 101: 224–232.

Speich SM, Manuwal DA, Wahl TR (1991) In my experience: the bird/habitat oil index: a habitat vulnerability index based on avian utilization. Wildlife Soc B: 216–221.

Spooner M (1971) Some ecological effects of marine oil pollution: 313–316.

Sprague JB, Holdway DA, Stendahl D (1978) Acute and chronic toxicity of vanadium to fish. Alberta Oil Sands Environmental Research Program.

Squires AJ, Dubé MG, Rozon-Ramilo LD (2013) Assessing the sublethal effects of in-river concentrations of parameters contributing to cumulative effects in the athabasca river basin using a fathead minnow bioassay. Env Toxicol Chem 32: 662–672.

- St. Clair C, Loots S (2012) Oil Sands Bird Contact Monitoring Plan for 2012.
- St. Clair C, Ronconi R, Ball J, Habib T (2011) Draft prospectus for instrument-based monitoring of birds in the oil sands.
- St. Clair CC (2009) Avian Deterrence in Alberta's Oil sands: Timing of Deployment, Habituation, and Minimum Program Requirements.
- St. Clair CC, Habib T, Loots S, Ball J, McCallum C (2012) 2011 Annual Report of the Regional Bird Monitoring Program for the Oil Sands Region.
- St. Clair CC, Habib T, Shore B (2012) Spatial and temporal correlates of mass bird mortality in oil sands tailings ponds. A report prepared for Alberta Environment. University of Alberta, Edmonton, Canada.
- St. Clair CC, Loots S, Gangadharan A, Gilhooly P, Jackson J (2014) 2013 Report of the Regional Bird Monitoring Program for the Oil Sands. Unpublished Work.
- St. Clair CC, Loots S, McCallum C, Gangadharan A (2014) Many thousands of migratory birds land annually, but few appear to die, in oil sands tailings ponds. (in progress).
- St. Clair CC, Loots S, McCallum C, Ronconi RA (2013) Oil Sands Bird Contact Monitoring Plan for 2013.
- St. Clair CC, Loots S, McCallum C, Thayer D, Fontaine T, Gilhooly P (2013) 2012 Report of the Regional Bird Monitoring Program for the Oil Sands; Department of Biological Sciences, University of Alberta: Edmonton, Alberta, Canada.
- St. Clair CC, Loots S, Ronconi RA (2014) 2014 Protocol Oil Sands Bird Contact Monitoring Program (OSBCMP).
- St. Clair CC, Ronconi RA (2009) Review of Alberta Environment 2009 compliance inspection reports and avian deterrence in the oil sands region. Prepared for Alberta Environment. 36 pp.
- St. Clair JJ, García-Peña GE, Woods RW, Székely T (2010) Presence of mammalian predators decreases tolerance to human disturbance in a breeding shorebird. Behav Ecol 21: 1285–1292.

Staddon JE (1975) A note on the evolutionary significance of supernormal stimuli. Am Nat: 541–545.

Stadt JJ, Schieck J, Stelfox HA (2006) Alberta Biodiversity Monitoring Program-monitoring effectiveness of sustainable forest management planning. Environ Monit Assess 121: 33–46.

Stamps J (1988) Conspecific attraction and aggregation in territorial species. Am Nat: 329–347.

Stankowich T, Blumstein DT (2005) Fear in animals: a meta-analysis and review of risk assessment. Proc R Soc B Biol Sci 272: 2627–2634.

Steen KA, Villa-Henriksen A, Therkildsen OR, Green O (2012) Automatic detection of animals in mowing operations using thermal cameras. Sensors (Basel, Switzerland) 12: 7587.

Steensma K (2009) Efficacy of bird deterrent devices in agricultural areas of the Fraser Valley of British Columbia: a pilot study.

Stein AB, Fuller TK, Marker LL (2008) Opportunistic use of camera traps to assess habitat-specific mammal and bird diversity in northcentral Namibia. Biodivers Conserv 17: 3579–3587.

Stephenson R (1997) Effects of oil and other surface-active organic pollutants on aquatic birds. Environ Conserv 24: 121–129.

Stephenson R, Andrews CA (1997) The effect of water surface tension on feather wettability in aquatic birds. Canadian J Zool 75: 288–294.

Stevens GR, Clark L (1998) Bird repellents: development of avian-specific tear gases for resolution of human-wildlife conflicts. Int Biodeterior Biodegradation 42: 153–160.

Stevens GR, Clark L, Weber RA (1998) The use of aerosol repellents as an avian deterrent strategy.

Stevens GR, Rogue J, Weber R, Clark L (2000) Evaluation of a radar-activated, demand-performance bird hazing system. Int Biodeterior Biodegradation 45: 129–137.

Stevens M, Cuthill IC (2007) Hidden messages: Are ultraviolet signals a special channel in avian communication? Biosci 57: 501–507.

Stevens M, PARraga CA, Cuthill IC, Partridge JC, Troscianko TS (2007) Using digital photography to study animal coloration. Biol J Linn Soc 90: 211–237.

Stewart GB, Pullin AS, Coles CF (2007) Poor evidence-base for assessment of windfarm impacts on birds. Environ Conserv 34: 1–11.

Stewart-Oaten A, Murdoch WW, Parker KR (1986) Environmental impact assessment: "Pseudoreplication" in time? Ecology 67: 929–940.

Stickel L, Dieter M (1979) Ecological and physiological/toxicological effects of petroleum on aquatic birds. Report: 14.

Stickley Jr AR, Mott DF, King JO (1995) Short-term effects of an inflatable effigy on cormorants at catfish farms. Wildlife Society (USA).

Stokstad E (2004) Songbirds Check Compass Against Sunset to Stay on Course. Science 304: 373–373.

Stone I, White J (1997) Case history: Bumblefoot in four oil-rehabilitated American Coots (*Fulica americana*). J Wildl Rehab 20: 9–13.

Stowe T, Underwood L (1984) Oil spillages affecting seabirds in the United Kingdom, 1966-1983. Mar Pollut Bull 15: 147–152.

Strategy CB (1995) Canadian Biodiversity Strategy: Canada's response to the convention on biological diversity. Biodiversity Convention Office, Environment Canada, Hull, Québec.

Strosher MT, Peake E (1976) The evaluation of wastewaters from an Oil Sands extraction plant.

Stubblefield W, McKee R, Kapp Jr R, Hinz J (1989) An evaluation of the acute toxic properties of liquids derived from oil sands. J Appl Toxicol 9: 59–65.

Stubblefield WA, Hancock GA, Ford WH, Ringer RK (1995) Acute and subchronic toxicity of naturally weathered Exxon Valdez crude oil in mallards and ferrets. Environ Toxicol Chem 14: 1941–1950.

Stubblefield WA, Hancock GA, Prince HH, Ringer RK (1995) Effects of naturally weathered Exxon Valdez crude oil on mallard reproduction. Environ Toxicol Chem 14: 1951–1960.

Stutchbury BJ, Gow EA, Done T, MacPherson M, Fox JW, et al. (2011) Effects of post-breeding moult and energetic condition on timing of songbird migration into the tropics. Proc R Soc B Biol Sci 278: 131–137.

Subcommittee. UNABCIM (2007) Opportunities for improving avian monitoring. US North American Bird Conservation Initiative Report. Available from the Division of Migratory Bird Management, US.

Sultana S, Hassan M, Choe HS, Ryu KS (2013) The effect of monochromatic and mixed LED light colour on the behaviour and fear responses of broiler chicken. Avian Biol Res 6: 207–214.

Summers R (1985) The effect of scarers on the presence of starlings (*Sturnus vulgaris*) in cherry orchards. Crop Prot 4: 520–528.

Sundar K, Kittur S (2013) Can wetlands maintained for human use also help conserve biodiversity? Landscape-scale patterns of bird use of wetlands in an agricultural landscape in north India. Biol Cons 168: 49–56.

Swaisgood RR (2010) The conservation-welfare nexus in reintroduction programs: a role for sensory ecology. Anim Welf 19: 125–137.

Swann DE, Hass CC, Dalton DC, Wolf SA (2004) Infrared-triggered cameras for detecting wildlife: an evaluation and review. Wildlife Soc B 32: 357–365.

Swift BL (2000) Suburban goose management: insights from New York State. Wildlife Damage Management Conferences-Proceedings. p. 44.

Syncrude Court Order 2011 No. 090157926P1 (2011).

Syncrude_Canada_Ltd. (1973) Migratory waterfowl and the Syncrude tar sands lease: a report. Environ Res Monogr1973-3.

Systad GH, Bustnes JO, Erikstad KE (2000) Behavioral responses to decreasing day length in wintering sea ducks. Auk 117: 33–40.

Szaro R, Albers P t (1977) Effects of external applications of No. 2 fuel oil on common eider eggs. Fate and Effects of Petroleum Hydrocarbons in Marine Organisms and Ecosystems D A Wolfe, Ed, Pergammon Press: 164–167.

Szaro RC (1977) Effects of petroleum on birds. Transactions of the Noth American Wildlife and Natural Resources Conferences. Vol. 42. pp. 374–381.

Szaro RC (1979) Bunker C fuel oil reduces mallard egg hatchability. Bull Environ Contam Toxicol 22: 731–732.

Szaro RC, Albers PH, Coon NC (1978) Petroleum: effects on mallard egg hatchability. The J Widlife Manag: 404–406.

Szaro RC, Coon NC, Stout W (1980) Weathered petroleum: Effects on mallard egg hatchability. The J Widlife Manag 44: 709–713.

Szaro RC, Dieter MP, Heinz GH, Ferrell JF (1978) Effects of chronic ingestion of South Louisiana crude oil on mallard ducklings. Environ Res 17: 426–436.

Szaro RC, Hensler G, Heinz GH (1981) Effects of chronic ingestion of no. 2 fuel oil on mallard ducklings. J Toxicol Environ Health 7: 789–799.

Taft OW, Haig SM, Kiilsgaard C (2003) Use of radar remote sensing (RADARSAT) to map winter wetland habitat for shorebirds in an agricultural landscape. Environ Manag 32: 268–281.

Tanaka KD, Morimoto G, Stevens M, Ueda K (2011) Rethinking visual supernormal stimuli in cuckoos: visual modeling of host and parasite signals. Behav Ecol 22: 1012–1019.

Tanis J, Mörzer Bruyns W (1969) The impact of oil-pollution on sea birds in Europe. International Conference on Oil Pollution of the Sea. Rome.

Tarshis IB, Rattner BA (1982) Accumulation of 14 C-naphthalene in the tissues of redhead ducks fed oil-contaminated crayfish. Arch Environ Contam Toxicol 11: 155–159.

Tasker ML, Jones PH, Dixon T, Blake BF (1984) Counting seabirds at sea from ships: a review of methods employed and a suggestion for a standardized approach. Auk: 567–577.

Taylor E, Stubblefield W (1997) Petroleum in the freshwater environment an annotated bibliography. Submitted to American Petroleum Institute Publication Number 4640.

Taylor PD, Brzustowski JM, Matkovich C, Peckford ML, Wilson D (2010) radR: an open-source platform for acquiring and analysing data on biological targets observed by surveillance radar. BMC Ecol 10: 22.

Taylor PD, Mackenzie SA, Thurber BG, Calvert AM, Mills AM, et al. (2011) Landscape movements of migratory birds and bats reveal an expanded scale of stopover. PLoS ONE 6: e27054.

Taylor SL, Pollard KS (2008) Evaluation of two methods to estimate and monitor bird populations. PLoS ONE 3: e3047.

Temple S, Pignatelli V, Cook T, How M, Chiou T-H, et al. (2012) High-resolution polarisation vision in a cuttlefish. Curr Biol 22: R121–R122.

Tenenbaum DJ (2009) Oil sands development: a health risk worth taking? Environmental health perspectives 117: A150.

Terzuoli A, McGowan J, Gustafson S, Jackson J, Martin R (2010) The effect of synthetic aperture radar image resolution on target discrimination. 38th COSPAR Scientific Assembly. Vol. 38. p. 391.

Tetreault GR, McMaster ME, Dixon DG, Parrott JL (2003) Physiological and biochemical responses of Ontario slimy sculpin (Cottus cognatus) to sediment from the Athabasca oil sands area. Water Qual Res J Canada 38: 361–377.

Tetreault GR, Mcmaster ME, Dixon DG, Parrott JL (2003) Using reproductive endpoints in small forage fish species to evaluate the effects of Athabasca oil sands activities. Environ Toxicol Chem 22: 2775–2782.

Thaxter CB, Burton NH (2009) High definition imagery for surveying seabirds and marine mammals: a review of recent trials and development of protocols. British Trust for Ornithology Report Commissioned by Cowrie Ltd.

Thomas L, Buckland ST, Burnham KP, Anderson DR, Laake JL, et al. (2006) Distance sampling. Encyclopedia of environmetrics. Wiley Online Library, Vol. 1. pp. 544–552.

Thomas R (2002) An updated, provisional bird inventory for the Peace-Athabasca Delta, northeastern Alberta.

Thompson SK, Ramsey FL (1987) Detectability functions in observing spatial point processes. Biometrics: 355–362.

Thompson WL (2001) Comparison of Three Plot Selection Methods for Estimating Change in Temporally Variable, Spatially Clustered Populations.

Thompson WL (2002) Towards reliable bird surveys: accounting for individuals present but not detected. Auk 119: 18–25.

Thorup K, Alerstam T, Hake M, Kjellén N (2006) Traveling or stopping of migrating birds in relation to wind: an illustration for the osprey. Behav Ecol 17: 497–502.

Thuppil V, Coss RG (2013) Wild Asian elephants distinguish aggressive tiger and leopard growls according to perceived danger. Biol Lett 9.

Tietje WD, Ruff RL (1983) Responses of black bears to oil development in Alberta. Wildlife Soc B: 99–112.

Timoney KP (2009) Three centuries of change in the Peace-Athabasca Delta, Canada. Clim Change 93: 485–515.

Timoney KP, Lee P (2009) Does the Alberta tar sands industry pollute? The scientific evidence. Open Conserv Biol J 3: 65–81.

Timoney KP, Lee P (2011) Polycyclic aromatic hydrocarbons increase in Athabasca River Delta sediment: Temporal trends and environmental correlates. Environ Sci Technol 45: 4278–4284.

Timoney KP, Ronconi RA (2010) Annual bird mortality in the bitumen tailings ponds in northeastern Alberta, Canada. Wilson J Ornithol 122: 569–576.

Tinbergen N, Perdeck AC (1950) On the stimulus situation releasing the begging response in the newly hatched herring gull chick (*Larus argentatus argentatus Pont.*). Behaviour: 1–39.

Tipton AR, Rappole JH, Kane AH, Flores RH, Johnson D, et al. (1989) Use of monofilament line, reflective tape, beach-balls, and pyrotechnics for controlling grackle damage to citrus. Great Plains Wildlife Damage Control Workshop Proceedings. p. 413.

Tisdale V, Fernández-Juricic E (2009) Vigilance and predator detection vary between avian species with different visual acuity and coverage. Behav Ecol 20: 936–945.

Toor NS, Franz ED, Fedorak PM, MacKinnon MD, Liber K (2012) Degradation and aquatic toxicity of naphthenic acids in oil sands process-affected waters using simulated wetlands. Chemosphere.

Trail PW (2006) Avian mortality at oil pits in the United States: a review of the problem and efforts for its solution. Environ Manag 38: 532–544.

Transport Canada (2002) Wildlife control procedures manual. TP11500E

Transport Canada (2010) Bird control at airports. Available: http://www.tc.gc.ca/eng/civilaviation/publications/tp13029-airports-115.htm.

Transport Canada (n.d.) Wildlife control: section e - active management using dispersal techniques. Available:

http://www.tc.gc.ca/civilaviation/aerodromeairnav/standards/wildlifecontrol/tp11500/sectione/sectione5.htm. Accessed 3 October 2009.

Trathan PN (2004) Image analysis of color aerial photography to estimate penguin population size. Wildlife Soc B 32: 332–343.

Trimper PG, Thomas PW, Chubbs TE (2008) Harlequin ducks in Labrador. Waterbirds 31: 32–43.

Trivelpiece WZ, Butler RG, Miller DS, Peakall DB (1984) Reduced survival of chicks of oil-dosed adult Leach's storm-petrels. Condor 86: 81–82.

Troisi G, Borjesson L (2005) Development of an immunoassay for the determination of polyaromatic hydrocarbons in plasma samples from oiled seabirds. Environ Sci Technol 39: 3748–3755.

Troisi G, Borjesson L, Bexton S, Robinson I (2007) Biomarkers of polycyclic aromatic hydrocarbon (PAH)-associated hemolytic anemia in oiled wildlife. Enviro Res 105: 324–329.

Troisi GM (2013) T-cell responses in oiled guillemots and swans in a rehabilitation setting. Arch Environ Contam Toxicol 65: 1–7.

Troisi GM, Bexton S, Robinson I (2006) Polyaromatic hydrocarbon and PAH metabolite burdens in oiled Common Guillemots (*Uria aalge*) stranded on the East Coast of England (2001-2002). Environ Sci Technol 40: 7938–7943.

Truett JC, Miller ME, Kertell K (1997) Effects of Arctic Alaska oil development on brant and snow geese. Arctic 50: 138–146.

Trust KA, Esler D, Woodin BR, Stegeman JJ (2000) Cytochrome P450 1A induction in sea ducks inhabiting nearshore areas of Prince William Sound, Alaska. Mar Pollut Bull 40: 397–403.

Tsai J-S, Venne LS, Smith LM, McMurry ST, Haukos DA (2012) Influence of local and landscape characteristics on avian richness and density in Wet Playas of the Southern Great Plains, USA. Wetlands 32: 605–618.

Tseng FS (1993) Care of oiled seabirds: A veterinary perspective. Abstract presented at Oil Spill Conference, ed. J. Ludwigson. American Petroleum Institute, Washington, DC. pp. 421–424.

Tseng FS (1999) Considerations in care for birds affected by oil spills. Seminars in Avian and Exotic Pet Medicine 8: 21–31.

Tubaro PL, Lijtmaer DA (2002) Hybridization patterns and the evolution of reproductive isolation in ducks. Biol J Linn Soc 77: 193–200.

Tupper SK, Cummings JL, Andelt WF, Werner SJ, Harness RE (2011) Evaluation of sonic dissuader to reduce damage by Pileated Woodpeckers. Wildlife Soc B 35: 40–44.

Twiss MP, Thomas VG (1998) Preventing fishing-sinker-induced lead poisoning of common loons through Canadian policy and regulative reform. J Environ Manag 53: 49–59.

Underhill LG, Bartlett PA, Baumann L, Crawford RJM, Dyer BM, et al. (1999) Mortality and survival of African Penguins *Spheniscus demersus* involved in the Apollo Seaoil spill: an evaluation of rehabilitation efforts. Ibis 141: 29–37.

Univeristy of British Columbia, City of Vancouver (2012) From sea to sky. In: 5th North American Ortnithological Conference, Vancouver.

Valera E, Powter CB (2012) Implications of changing environmental requirements on oil sands royalties. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-23. 21 pp.

Vallorttgara G, Zanforlin M (1988) Open-field behavior of young chicks (*Gallus gallus*): antipredatory responses, social reinstatement motivation, and gender effects. Anim Learn Behav 16: 359–362.

Van Belle J, Shamoun-Baranes J, Van Loon E, Bouten W (2007) An operational model predicting autumn bird migration intensities for flight safety. J Appl Ecol 44: 864–874.

Van den Heuvel M, Dixon DG (1998) Assessment of Fathead Minnow Population Health Parameters Associated with the Syncrude Fine Tailings Capping Demonstration Ponds.

Van den Heuvel M, Power M, MacKinnon M, Dixon D (1999) Effects of oil sands related aquatic reclamation on yellow perch (*Perca flavescens*). II. Chemical and biochemical indicators of exposure to oil sands related waters. Can J Fish Aquat Sci 56: 1226–1233.

Van den Heuvel M, Power M, MacKinnon M, Meer TV, Dobson E, et al. (1999) Effects of oil sands related aquatic reclamation on yellow perch (*Perca flavescens*). I. Water quality characteristics and yellow perch physiological and population responses. Can J Fish Aquat Sci 56: 1213–1225.

Van den Heuvel M, Power M, Richards J, MacKinnon M, Dixon D (2000) Disease and gill lesions in Yellow Perch (*Perca flavescens*) exposed to oil sands mining-associated waters. Ecotoxicol Environ Saf 46: 334–341.

Van den Heuvel MR, Hogan NS, MacDonald GZ, Berrue F, Young RF, et al. (2013) Assessing accumulation and biliary excretion of naphthenic acids in yellow perch exposed to oil sandsaffected waters. Chemosphere 95: 619–627.

Van Gasteren J, Holleman I, Bouten W, van Loon E, Baranes JS (2008) Extracting bird migration information from C-band Doppler weather radars. Ibis 150: 674–686.

Van Horne B (1983) Density as a misleading indicator of habitat quality. The J Widlife Manag: 893–901.

Van Kampen D (1971) Successful cleaning of oiled birds in Holland. Mar Pollut Bull 2: 140–142.

Van Meer T, Arner B (1985) Bird surveillance and protection programme, summary of 1984 and 1985 activities. Syncrude Canada Ltd, Calgary, Alberta, Canada

Van Wilgenburg SL, Hobson KA, Bayne EM, Koper N (2013) Estimated avian nest loss associated with oil and gas exploration and extraction in the western Canadian sedimentary basin. Avian Conserv Ecol 8: 9.

Vangilder LD, Peterle TJ (1980) South Louisiana crude oil and DDE in the diet of mallard hens: Effects on reproduction and duckling survival. Bull Environ Contam Toxicol 25: 23–28.

Vangilder LD, Peterle TJ (1981) South Louisiana crude oil or DDE in the diet of mallard hens: effects on egg quality. Bull Environ Contam Toxicol 26: 328–336.

Vangilder LD, Peterle TJ (1983) Mallard egg quality: enhancement by low levels of petroleum and chlorinated hydrocarbons. Bull Environ Contam Toxicol 30: 17–23.

Vangilder LD, Peterle TJ (1984) Impairment of mallard reproduction by dietary crude oil or DDE. Actas del XV Congreso Internacional de Fauna Cinegetica y Silvestre, Trujill (Cáceres), España, 1981. Transactions of the Congress of the International Union of Game Biologists. Estacion Biologica de Doñana, Federacion Española de Caza, Seville, Spain. p. 107–119.

Vaughn CR (1985) Birds and insects as radar targets: A review. Proceedings of the IEEE 73: 205–227.

Vauk G, Hartwig E, Reineking B, Schrey E, Vauk-Hentzelt E (1990) Extensive investigations into the effects on seabirds of the oil pollution on the German North Sea coast. The International Union of Game Biologists: the 19. IUGB Congress, September 1989, Trondheim, Norway: transactions. p. 288.

Velando A, Alvarez D, Mourino J, Arcos F, Barros A (2005) Population trends and reproductive success of the European shag *Phalacrocorax aristotelis* on the Iberian Peninsula following the Prestige oil spill. J Ornithol146: 116–120.

Velando A, Munilla I, Leyenda PM (2005) Short-term indirect effects of the Prestige' oil spill on European shags: changes in availability of prey. Mar Ecol Prog Ser 302: 263–274.

Verbeek A (1994) A toxicity assessment of oil sands wastewater: a toxic balance.

VerCauteren KC, Gilsdorf JM, Hygnstrom SE, Fioranelli PB, Wilson JA, et al. (2006) Green and blue lasers are ineffective for dispersing deer at night. Wildlife Soc B 34: 371–374.

Vermeer K (1976) Colonial auks and eiders as potential indicators of oil pollution. Mar Pollut Bull 7: 165–167.

Vermeer K, Anweiler GG (1975) Oil threat to aquatic birds along the Yukon coast. Wilson Bull: 467–480.

Verstraeten WW, Vermeulen B, Stuckens J, Lhermitte S, Van der Zande D, et al. (2010) Webcams for bird detection and monitoring: A demonstration study. Sensors 10: 3480–3503.

Vicen-Bueno R, Rosa-Zurera M, Jarabo-Amores M, Gil-Pita R (2010) Automatic target detection in simulated ground clutter (Weibull distributed) by multilayer perceptrons in a low-resolution coherent radar. Radar, Sonar \& Navigation, IET 4: 315–328.

Vickery JA, Summers RW (1992) Cost-effectiveness of scaring brent geese *Branta b. bernicla* from fields of arable crops by a human bird scarer. Crop Prot 11: 480–484.

Vidal M, Dominguez J, Luis A (2011) Spatial and temporal patterns of polycyclic aromatic hydrocarbons (PAHs) in eggs of a coastal bird from northwestern Iberia after a major oil spill. Sci Total Environ 409: 2668–2673.

Vikebø FB, Rønningen P, Lien VS, Meier S, Reed M, et al. (2013) Spatio-temporal overlap of oil spills and early life stages of fish. ICES J Mar Sci : 2-12.

Von Bayern AM, Emery NJ (2009) Jackdaws respond to human attentional states and communicative cues in different contexts. Curr Biol 19: 602–606.

Vorobyev M, Osorio D, Bennett A, Marshall N, Cuthill I (1998) Tetrachromacy, oil droplets and bird plumage colours. J Comp Physiol A Neuroethol Sens Neural Behav Physiol 183: 621–633.

Vorotkov M, Sinelschikova A, Griffiths M (2009) Optical matrix device: technical aspects of a new tool for the detection and recording of small nocturnal aerial targets. J Navig 62: 23-32.

Votier S, Birkhead T, Oro D, Trinder M, Grantham M, et al. (2008) Recruitment and survival of immature seabirds in relation to oil spills and climate variability. J Anim Ecol 77: 974–983.

Votier SC, Hatchwell BJ, Beckerman A, McCleery RH, Hunter FM, et al. (2005) Oil pollution and climate have wide-scale impacts on seabird demographics. Ecol Lett 8: 1157–1164.

Vymazal J (2008) Constructed wetlands for wastewater treatment: a review. The 12th World Lake Conference. pp. 965–980.

Wallace RR (2013) Geoscience of Climate and Energy 9. History and governance models as a blueprint for future federal-provincial co-operation on environmental monitoring in the Alberta Oil Sands region. GeoSci Can 40: 167–173.

Wallraff H (1991) Conceptual approaches to avian navigation systems. Experientia 46: 379–388.

Walsh DA, Otter KA (2011) A Technique for Avian Nocturnal Migration Research: Determining Flight Patterns of Local Nocturnal Migrants around Pre-operational Wind Turbines.

Walters CJ (1986) Adaptive Management of Renewable Resources. Vancouver, BC: MacMillan Publishing Company.

Walters CJ, Holling CS (1990) Large-scale management experiments and learning by doing. Ecology 71: 2060–2068.

Walters P, Khan S, O'Brien P, Payne J, Rahimtula A (1987) Effectiveness of a Prudhoe Bay crude oil and its aliphatic, aromatic and heterocyclic fractions in inducing mortality and aryl hydrocarbon hydroxylase in chick embryo in ovo. Arch Toxicol 60: 454–459.

Walton P, Turner CMR, Austin G, Burns MD, Monaghan P (1997) Sub-lethal effects of an oil pollution incident on breeding kittiwakes Rissa tridactyla. Mar Ecol Prog Ser 155: 261–268.

Wang N, Chelme-Ayala P, Perez-Estrada L, Garcia-Garcia E, Pun J, et al. (2013) Impact of ozonation on naphthenic acids speciation and toxicity of oil sands process-affected water to *Vibrio fischeri* and mammalian immune system. Environ Sci Technol 47: 6518–6526.

Wang Z, Yang C, Parrott JL, Frank RA, Yang Z, et al. (2014) Forensic source differentiation of petrogenic, pyrogenic, and biogenic hydrocarbons in Canadian oil sands environmental samples. J Hazard Mater 271C: 166–177.

Ward JG (1978) Tests of the Syncrude bird deterrent device for use on a tailings pond. Syncrude Canada Limited.

Warith MA, Yong RN (1994) Toxicity assessment of sludge fluid associated with tar sand tailings. Environ Technol 15: 381–387.

Warnken J, Buckley R (1998) Scientific quality of tourism environmental impact assessment. J Appl Ecol: 1–8.

Washburn BE, Chipman RB, Francoeur LC (2006) Evaluation of bird response to propane exploders in an airport environment. Proceedings of the Vertebrate Pest Conference. Vol. 22. pp. 212–215.

Wasser SK, Keim JL, Taper ML, Lele SR (2011) The influences of wolf predation, habitat loss, and human activity on caribou and moose in the Alberta oil sands. Front Ecol Environ 9: 546–551.

Watson BM, Putz G (2011) Preliminary watershed hydrology model for reclaimed oil sands sites. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-39. 193 pp.

Watson GPL (2011) Field birding and digital objects: immaterial technologies and their implications for one practice of coming to know the more-than-human. Environ Educ Res 17: 789–799.

Watve M, Thakar J, Kale A, Puntambekar S, Shaikh I, et al. (2002) Bee-eaters (*Merops orientalis*) respond to what a predator can see. Anim Cogn 5: 253–259.

Wayland M, Headley JV, Peru KM, Crosley R, Brownlee BG (2008) Levels of polycyclic aromatic hydrocarbons and dibenzothiophenes in wetland sediments and aquatic insects in the oil sands area of Northeastern Alberta, Canada. Environ Monit Assess 136: 167–182.

Wayland M, Smits JE (2004) The ecological viability of constructed wetlands at Suncor: population and health- related considerations in birds. Assessment of natural and anthropogenic

impacts of oil sands contaminants within the northern river basins Environment Canada, Saskatoon, Saskatchewan, Canada. pp. 48–57.

Webb EB, Smith LM, Vrtiska MP, Lagrange TG (2010) Community structure of wetland birds during spring migration through the rainwater basin. The J Widlife Manag 74: 765–777.

Webb SM (2009) Marten Fur Harvests and Landscape Change in West-Central Alberta. The J Widlife Manag 73: 894–903.

Weber LM, Haig SM (1997) Shorebird-prey interactions in South Carolina coastal soft sediments. Canadian J Zool 75: 245–252.

Welham C (2010) Oil sands terrestrial habitat and risk modeling for disturbance and reclamation – phase i report. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-8. 109 pp.

Wells J, Casey-Lefkowitz S, Dyer S, Chavarria G (2008) Danger in the Nursery: Impact on Birds of Tar Sands Oil Development in Canada's Boreal Forest. Natural Resources Defense Council.

Welsh P (n.d.) The toxicological responses of avian species to hydrocarbons: Implications for the management of oil sands process-affected water.

Werner SJ, Buchholz R, Tupper SK, Pettit SE, Ellis JW (2013) Functional Significance Of Ultraviolet Feeding Cues In Wild Turkeys. *Physiol Behav*.

Werner SJ, Clark L (2006) Effectiveness of a motion-activated laser hazing system for repelling captive Canada geese. Wildlife Soc B 34: 2–7.

Western EcoSystems Technology I (2014) Review of Potential Bird Deterrent Strategies for Large Scale Solar Facilities.

Whelly MP, Ciborowski JJH, Leonhardt C, Laing D (1998) Chironomidae as Indicators of Wetland Viability: Report on Field Work in Wetlands of the Fort McMurray, Alberta Area.

Whisson DA, Takekawa JY (2000) Testing the effectiveness of an aquatic hazing device on waterbirds in the San Francisco Bay estuary of California. Waterbirds 23: 56–63.

Whitby C (2010) Microbial naphthenic acid degradation. Adv Appl Microbiol 70: 93–125.

White DH, King KA, Coon NC (1979) Effects of No. 2 fuel oil on hatchability of marine and estuarine bird eggs. Bull Environ Contam Toxicol 21: 7–10.

Whitmore RC, Mosher JA, Frost HH (1977) Spring migrant mortality during unseasonable weather. Auk: 778–781.

Whittington PA (2003) Post-release survival of rehabilitated African Penguins. Rehabilitation of oiled African Penguins: a conservation success story. Bird Life South Africa and the Avian Demography Unit, Johannesburg and Cape Town: 8–10.

Wiens JA (1989) Spatial scaling in ecology. Funct Ecol 3: 385–397.

Wiens JA (1995) Recovery of seabirds following the Exxon Valdez oil spill: an overview. ASTM Special Technical Publication: 854–893.

Wiens JA (1996) Oil, seabirds, and science: the effects of the Exxon Valdez oil spill. Biosci 46: 587-597.

Wiens JA (2007) Applying ecological risk assessment to environmental accidents: Harlequin ducks and the Exxon Valdez oil spill. Biosci 57: 769–777.

Wiens JA, Chr N, Van Horne B, Ims RA (1993) Ecological mechanisms and landscape ecology. Oikos 66: 369–380.

Wiens JA, Crist TO, Day RH, Murphy SM, Hayward GD (1996) Effects of the Exxon Valdez oil spill on marine bird communities in Prince William Sound, Alaska. Ecol Appl 6: 828–841.

Wiens JA, Crist TO, Day RH, Murphy SM, Hayward GD (2001) A canonical correspondence analysis of the effects of the Exxon Valdez oil spill on marine birds. Ecol Appl 11: 828–839.

Wiens JA, Day RH, Murphy SM, Fraker MA (2010) Assessing Cause-Effect Relationships in Environmental Accidents: Harlequin Ducks and the Exxon Valdez Oil Spill. In: Thompson, C. F., editor. Current Ornithology. Vol. 17. pp. 131–189.

Wiens JA, Day RH, Murphy SM, Parker KR (2004) Changing habitat and habitat use by birds after the Exxon Valdez oil spill, 1989-2001. Ecol Appl 14: 1806–1825.

Wiens JA, Ford RG, Heinemann D (1984) Information needs and priorities for assessing the sensitivity of marine birds to oil spills. Biol Cons 28: 21–49.

Wiens JA, Hayward GD, Holthausen RS, Wisdom MJ (2008) Using surrogate species and groups for conservation planning and management. Biosci 58: 241–252.

Wiese FK, Jones IL, Nettleship D (2001) Experimental support for a new drift block design to assess seabird mortality from oil pollution. Auk 118: 1062–1068.

Wiese FK, Montevecchi W, Davoren G, Huettmann F, Diamond A, et al. (2001) Seabirds at risk around offshore oil platforms in the North-west Atlantic. Mar Pollut Bull 42: 1285–1290.

Wiese FK, Robertson GJ (2004) Assessing seabird mortality from chronic oil discharges at sea. The J Widlife Manag 68: 627–638.

Wiese FK, Robertson GJ, Gaston AJ (2004) Impacts of chronic marine oil pollution and the murre hunt in Newfoundland on thick-billed murre *Uria lomvia* populations in the eastern Canadian Arctic. Biol Cons 116: 205–216.

Wiese FK, Ryan PC (2003) The extent of chronic marine oil pollution in southeastern Newfoundland waters assessed through beached bird surveys 1984-1999. Mar Pollut Bull 46: 1090–1101.

Wiklund JA, Hall RI, Wolfe BB, Edwards TW, Farwell AJ, et al. (2012) Has Alberta oil sands development increased far-field delivery of airborne contaminants to the Peace-Athabasca Delta? Sci Total Environ 433: 379–382.

Wilhelm SI, Robertson GJ, Ryan PC, Schneider DC (2007) Comparing an estimate of seabirds at risk to a mortality estimate from the November 2004 Terra Nova FPSO oil spill. Mar Pollut Bull 54: 537–544.

Wilhelm SI, Robertson GJ, Ryan PC, Tobin SF, Elliot RD (2009) Re-evaluating the use of beached bird oiling rates to assess long-term trends in chronic oil pollution. Mar Pollut Bull 58: 249–255.

Williams JM, Tasker ML, Carter IC, Webb A (1995) A method of assessing seabird vulnerability to surface pollutants. Ibis 137: S147–S152.

Williams T, Williams J (1990) Open ocean bird migration. Radar and Signal Processing, IEE Proceedings F. Vol. 137. pp. 133–137.

Williams TC (1988) Bias in Radar Data Recording Techniques . J Field Ornithol 59: 63–66.

Williams TC, Marsden JE, Lloyd-Evans TL, Krauthamer V (1981) Spring migration studied by mist-netting, ceilometer, and radar. J Field Ornithol 52: 177–190.

Williams TC, Webb III T (1996) Neotropical bird migration during the ice ages: orientation and ecology. Auk: 105–118.

Williams TC, Williams JM (1988) Radar and visual observations of autumnal (southward) shorebird migration on Guam. Auk: 460–466.

Williams TC, Williams JM, Kloeckner PD (1986) Airspeed and heading of autumnal migrants over Hawaii. Auk 103: 634–635.

Williams TC, Williams JM, Williams PG, Stokstad P (2001) Bird migration through a mountain pass studied with high resolution radar, ceilometers, and census. Auk 118: 389–403.

Williams TC, Ying M (1990) A comparison of radar observations of bird migration at Haizhou Bay, China, and Guam, Marianas. Auk 107: 404–406.

Wiltschko R, Stapput K, Thalau P, Wiltschko W (2010) Directional orientation of birds by the magnetic field under different light conditions. J R Soc Interface 7: S163–S177.

Wiltschko R, Wiltschko W (2013) The magnetite-based receptors in the beak of birds and their role in avian navigation. J Comp Physiol A Neuroethol Sens Neural Behav Physiol 199: 89-98.

Wiltschko W, Munro U, Ford H, Wiltschko R (1993) Red light disrupts magnetic orientation of migratory birds. Nature 364: 525–527.

Wiltschko W, Munro U, Ford H, Wiltschko R (2006) Bird navigation: what type of information does the magnetite-based receptor provide? Proc R Soc B Biol Sci 273: 2815–2820.

Wiltschko W, Wiltschko R (1995) Migratory orientation of European robins is affected by the wavelength of light as well as by a magnetic pulse. J Comp Physiol A Neuroethol Sens Neural Behav Physiol 177: 363–369.

Wiseman S, Anderson J, Liber K, Giesy J (2013) Endocrine disruption and oxidative stress in larvae of *Chironomus dilutus* following short-term exposure to fresh or aged oil sands process-affected water. Aquat Toxicol 142: 414–421.

Witherington BE (1997) The problem of photopollution for sea turtles and other nocturnal animals. Behavioral approaches to conservation in the wild. Cambridge University Press Cambridge. pp. 303–328.

Wolfaardt AC, Underhill LG, Altwegg R, Visagie J, Williams AJ (2008) Impact of the treasure oil spill on African penguins *Spheniscus demersus* at Dassen Island: case study of a rescue operation. Afr J Mar Sci 30: 405–419.

Wolfaardt AC, Underhill LG, Nel DC, Williams AJ, Visagie J (2008) Breeding success of African penguins *Spheniscus demersus* at Dassen Island, especially after oiling following the Apollo Sea spill. Afr J Mar Sci 30: 565–580.

Wolfaardt AC, Williams AJ, Underhill LG, Crawford RJM, Whittington PA (2009) Review of the rescue, rehabilitation and restoration of oiled seabirds in South Africa, especially African penguins *Spheniscus demersus* and Cape gannets *Morus capensis*, 1983–2005. Afr J Mar Sci 31: 31–54.

Wood AW, Barjaktarovic L, Kennedy CJ, Farrell AP, Bendell-Young LI (1995) Fate of U-4C-palmitic acid in constructed wetlands: A microcosm study.

Wood MA, Heaphy N (1991) Rehabilitation of oiled seabirds and bald eagles following the Exxon Valdez oil spill. International Oil Spill Conference API Publication 4529: 235–239.

Wootton TA, Grau CR, Roudybush TE, Hahs ME, Hirsch KV (1979) Reproductive responses of quail to bunker-c oil fractions. Arch Environ Contam Toxicol 8: 457–463.

Woynillowicz D, Severson-Baker C, Raynolds M (2005) Oil sands fever: The environmental implications of Canada's oil sands rush. Pembina Institute Drayton Valley, Alberta, Canada.

Wu L-Q, Dickman JD (2012) Neural correlates of a magnetic sense. Science 336: 1054–1057.

Wyatt S, Fortier J-F, Natcher DC, Smith MAP, Hébert M (2013) Collaboration between Aboriginal peoples and the Canadian forest sector: A typology of arrangements for establishing control and determining benefits of forestlands. J Environ Manag 115: 21–31.

Xu JG, Johnson RL, Yeung PYP, Wu S (1995) Plant growth and metal uptake by plants from two oil sand fine tailings (A Progress Report on Greenhouse Experiment).

Xu JG, Johnson RL, Yeung PYP, Wu S (1996) Plant growth, dewatering and contaminant uptake from oil sands fine tails and tailings (A progress report on the 1995 field experiment).

Xu JG, Johnson RL, Yeung PYP, Wu S (1997) Plant Growth and Metal Uptake from Oil Sands Fine Tails and Tailings - A Progress Report on the 1996 Field Experiment. Alberta Research Council, Vegreville, AB.

Yamato O, Goto I, Maede Y (1996) Hemolytic anemia in wild seaducks caused by marine oil pollution. J Wildl Dis 32: 381–384.

Yaukey PH, Powell SC (2008) Numbers of migratory birds stopping over in New Orleans, Louisiana, USA in relation to weather. Wilson J Ornithol 120: 286–295.

Yoccoz NG, Nichols JD, Boulinier T (2001) Monitoring of biological diversity in space and time. Trends Ecol Evol 16: 446–453.

Yonge KS (1979) Development of a bird protection strategy for tar sands tailings ponds. Proceedings of the 8th Bird Control Seminar, Bowling Green, Ohio, Oct.

Yonge KS, Christiansen ML (1979) A review of bird migration patterns and techniques for monitoring migration. Professional Paper 1981-2. Syncrude Canada Ltd, Calgary, Alberta, Canada.

Yonge KS, Christiansen ML, Samoil HP, Smith JA, Van Meer T (1981) Syncrude bird surveillance program: A methodology for assessing bird activity and bird losses associated with the tailings pond.

Young Jr D, Erickson W, Strickland M, Good R, Sernka K (2003) Comparison of avian responses to UV-light-reflective paint on wind turbines. National Renewable Energy Laboratory, Golden, Colorado, USA. Subcontract report,

Young R, Orr E, Goss G, Fedorak P (2007) Detection of naphthenic acids in fish exposed to commercial naphthenic acids and oil sands process-affected water. Chemosphere 68: 518–527.

Young RF, Michel LM, Fedorak PM (2011) Distribution of naphthenic acids in tissues of laboratory-exposed fish and in wild fishes from near the Athabasca oil sands in Alberta, Canada. Ecotoxicol Environ Saf 74: 889–896.

Z. Hashisho CCS, Morshed G (2012) Review of technologies for the characterization and monitoring of vocs, reduced sulphur compounds and CH₄. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-19. 93 pp.

Zanette LY, White AF, Allen MC, Clinchy M (2011) Perceived predation risk reduces the number of offspring songbirds produce per year. Science 334: 1398–1401.

Zaugg S, Saporta G, Van Loon E, Schmaljohann H, Liechti F (2008) Automatic identification of bird targets with radar via patterns produced by wing flapping. J R Soc Interface 5: 1041–1053.

Zehnder S, Åkesson S, Liechti F, Bruderer B (2001) Nocturnal autumn bird migration at Falsterbo, South Sweden. J Avian Biol 32: 239–248.

Zhang J, Xu Q, Cao X, Yan P, Li X (2013) Hierarchical Incorporation of Shape and Shape Dynamics for Flying Bird Detection. Neurocomputing.

Zhang X, Wu X, Zhou X, Wang X, Zhang Y (2009) Birds' Flap Frequency Measure Based on Automatic Detection and Tracking in Captured Videos. 13th International Conference on Biomedical Engineering. pp. 1858–1861.

Zhang Y, Wu L, Wei G (2009) A new classifier for polarimetric SAR images. Progress In Electromagnetics Research 94: 83–104.

Zhao B, Currie R, Mian H (2012) Catalogue of analytical methods for naphthenic acids related to oil sands operations. Oil Sands Research and Information Network, University of Alberta, School of Energy and the Environment, Edmonton, Alberta. OSRIN Report No. TR-21. 65 pp.

Zhao S, Kotlyar LS, Woods JR, Sparks BD, Gao J, et al. (2002) A benchmark assessment of residues: comparison of Athabasca bitumen with conventional and heavy crudes. Fuel 81: 737–746.

Zimmer C, Boos M, Bertrand F, Robin J-P, Petit O (2011) Behavioural Adjustment in Response to Increased Predation Risk: A Study in Three Duck Species. PLoS ONE 6: e18977.

Zimmerling JR, Pomeroy AC, d' Entremont MV, Francis CM (2013) Canadian estimate of bird mortality due to collisions and direct habitat loss associated with wind turbine developments. Avian Conserv Ecol 8: 10.

Zock J-P, Rodríguez-Trigo G, Rodríguez-Rodríguez E, Souto-Alonso A, Espinosa A, et al. (2014) Evaluation of the persistence of functional and biological respiratory health effects in clean-up workers 6 years after the Prestige oil spill. Environ Int 62: 72–77.

Zuberogoitia I, Martínez JA, Iraeta A, Azkona A, Zabala J, et al. (2006) Short-term effects of the prestige oil spill on the peregrine falcon (*Falco peregrinus*). Mar Pollut Bull 52: 1176–1181.

Zuclich J, Connolly J (1976) Ocular damage induced by near-ultraviolet laser radiation. Invest Ophthalmol Vis Sci 15: 760–764.

Zuercher F, Thuer M (1978) Rapid weathering processes of fuel oil in natural waters: analyses and interpretations. Environ Sci Technol 12: 838–843.

Zyla W, Prepas E (1995) The effects of nutrient additions on phytoplankton from Syncrude Canada Ltd.'s water capped lake systems.